

1-1-1999

## Changes in the whole body mineral concentration of the Rainbow trout (*Oncorhynchus mykiss*Walbaum) yolk-sac fry exposed to various combinations of aluminium and calcium concentrations in two different acidic water

METİN ÇALTA

Follow this and additional works at: <https://journals.tubitak.gov.tr/zoology>



Part of the [Zoology Commons](#)

---

### Recommended Citation

ÇALTA, METİN (1999) "Changes in the whole body mineral concentration of the Rainbow trout (*Oncorhynchus mykiss*Walbaum) yolk-sac fry exposed to various combinations of aluminium and calcium concentrations in two different acidic water," *Turkish Journal of Zoology*. Vol. 23: No. 3, Article 13. Available at: <https://journals.tubitak.gov.tr/zoology/vol23/iss3/13>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Zoology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Changes in the whole body mineral concentration of the Rainbow trout (*Oncorhynchus mykiss* Walbaum) yolk-sac fry exposed to various combinations of aluminium and calcium concentrations in two different acidic water

Metin ÇALTA  
Firat Üniversitesi, Su Fakültesi, Temel Bilimler Bölümü, Elazığ-TURKEY

Received: 06.10.1998

**Abstract:** The effects of 12 combinations of pH (4.5 and 7.0), aluminium (4 and 8 µmol/L) and calcium (10 and 100 µmol/L) on survival and body mineral (Ca, Mg, K and Na) concentrations of rainbow trout (*Oncorhynchus mykiss* Walbaum) at the yolk-sac stage were investigated.

Mortality was low (< 17%) in most combinations and increased (> 27%) in the presence of aluminium and low pH (4.5).

Aluminium (4 and 8 µmol/L) impaired larval development, net uptake of calcium, potassium and sodium, and slightly increased the net loss of magnesium. It was found that these effects of aluminium were more severe at low calcium concentration (10 µmol/L) and low pH (4.5) than at high calcium concentration (100 µmol/L) and high pH (7.0). In the absence of aluminium, only the combination of low pH (4.5) and low calcium (10 µmol/L) significantly reduced body mineral concentrations.

**Key Words:** Aluminium, pH, calcium, rainbow trout, *Oncorhynchus mykiss*, body mineral concentration.

### Farklı İki Asidik Su Ortamında Değişik Aliminyum ve Kalsiyum Kombinasyonlarına Maruz Bırakılan Yumurta Kesesi Evresindeki Gökkuşuğu Alabalığı (*Oncorhynchus mykiss* Walbaum)'nın Vücudunda Mineral Konsantrasyonu Değişimi

**Özet:** Bu çalışmada, pH (7.0 ve 4.5), aliminyum (0,4 ve 8 µmol/L) and kalsiyum (10 ve 100 µmol/L)'un 12 kombinasyonunun yumurta kesesi evresindeki gökkuşuğu alabalığı (*Oncorhynchus mykiss* Walbaum)'nın yaşamı ve vücut mineral (Ca, Mg, K and Na) konsantrasyonuna etkisi araştırıldı.

Ölüm oranı çoğu kombinasyonlarda %17'den daha düşük olup, pH (4.5) ve aliminyumun varlığında bu oran %27'nin üzerine çıktı.

Aliminyum (4 ve 8 µmol/L) larvaların gelişimini, kalsiyum, potasyum ve sodyum alımını yavaşlattı. Ayrıca azda olsa larvanın magnezyum kaybını artırdı. Aliminyumun bu etkilerinin, pH(4.5) ve calcium (10 µmol/L)'da, pH (7.0) ve calcium (100 µmol/L)'da kine oranla daha şiddetli olduğu gözlemlendi.

**Anahtar Sözcükler:** Aliminyum, pH, kalsiyum, gökkuşuğu alabalığı, *Oncorhynchus mykiss*, vucut mineral konsantrasyonu.

### Introduction

The rainbow trout originally inhabited the Pacific coasts of North America. It was introduced Europe in 1875 and later to other continents. The rainbow trout is the most widespread member of the family Salmonidae in Europe. It occurs both in the wild and in trout farms. Its enormous adaptability to water conditions and its economical value have made it an important subject for scientific studies.

Recently, a decline in the trout population was observed in North America and Northern Europe. The factors usually implicated in the decline of fish populations are a combination of low pH at low water hardness and the presence of toxic trace metals such as aluminium (1).

It has been determined that the main target of aluminium is the fish gills which are the main site of ion exchange. Disturbed ion balance has been observed in salmonids exposed to aluminium at low pH and low calcium concentrations (2, 3).

The present study examined the effects of combinations of two pH values and two calcium concentrations on the calcium, magnesium, potassium and sodium content of yolk-sac rainbow trout in the absence and presence of aluminium frequently found in acidic waters.

### Material and Methods

The experimental animals (newly hatched rainbow trout) were obtained from Firat University, Fisheries Faculty, Cip Fish Reproduction and Research Station. The

experiment was conducted in small aquariums (20x15x10 cm) placed in a cold room (12-15 °C) in the Fisheries Faculty. The experimental media were prepared with deionized water by the addition of major ions in concentrations given in Table 1. Two aluminium concentrations (4 and 8 µmol/L) were provided by the addition of AlCl<sub>3</sub>. Two pH regimes were adjusted by the addition of 0.1N H<sub>2</sub>SO<sub>4</sub> solution. All aquariums were aerated and dechlorinated before and during the experiments.

Table 1. Nominal concentrations of major ions in experimental media (µmol/L).

Ions		
Ca <sup>+2</sup>	10	100
K <sup>+</sup>	5	5
Mg <sup>+2</sup>	20	20
Na <sup>+</sup>	50	50
Cl <sup>-</sup>	60	150
NO <sub>3</sub> <sup>-</sup>	5	5
SO <sub>4</sub> <sup>-2</sup>	45	65

At the beginning of the experiments, 10 newly hatched larvae were killed and treated as described below for the determination of dry weight and the whole body mineral content, and the results were compared with those obtained at the end of the experiments. A total of 50 newly hatched larvae were transferred to 12 aquariums containing different experimental media (Table 2).

During the experiments, all aquariums were checked

daily and any dead animals were removed. At the same time, pH, conductivity and temperature measurements were taken in all aquariums. At 3 day intervals, water samples were taken from each aquarium for analysis of total aluminium and calcium. The concentrations of total aluminium and calcium were determined with an atomic absorption spectrophotometer (Perkin-Elmer model 370).

Experiments continued the yolk-sac phase of the larvae (approximately 30 days). After absorption of yolk by the control animals, 10 of the surviving larvae were removed from each aquarium and they were killed in anaesthetic solution (Sandoz, MS 222) and rinsed twice deionized water. In order to determine the dry weight and the whole body mineral content in dry weight, the animals were transferred individually into 4 ml preweighed vials. They were dried in an oven (60-65 °C) for 24 h and then transferred to a low pressure desiccator for several days until constant mass was obtained. After weighing for dry weight; the larvae were digested in HNO<sub>3</sub> 16 µmol/L for 24 h at room temperature, and then the digests were diluted with LaCl<sub>3</sub> in deionized water to give [La]=1% w/v. Ca, K, Mg and Na concentrations were determined in the whole body of the larvae with an atomic absorption spectrophotometer. A blank was prepared for comparison, following the same procedure used for the digests.

The results were expressed as mean ± standard error. Statistical analyses were performed by using Analysis of Variance (ANOVA) and Kruskal-Wallis Test.

Aquarium	pH		Total aluminium (µmol/L)		Calcium (µmol/L)		Temperature (°C)	Conductivity (µS/cm)
	Nominal	Actual	Nominal	Actual*	Nominal	Actual*	Actual	Actual
No			0	0	10	9±0.4	13±1.5	17.6±0.62
1			0	0	100	92±6	13±2.0	38.8±1.2
2			4	3.65±0.20	10	8±0.6	14±1.0	18.0±0.40
3	7.0	6.7-7.4	4	3.82±0.18	100	90±5	13±1.5	37.6±0.9
4			8	7.60±0.51	10	8±0.5	14±1.0	16.8±0.54
5			8	7.45±0.60	100	93±4	13±2.0	39.2±1.1
6			0	0	10	9±0.6	14±1.5	17.4±0.33
7			0	0	100	95±4	13±2.0	38.5±0.8
8			4	3.78±0.17	10	9±0.4	13±1.5	17.6±0.40
9	4.5	4.2-4.6	4	3.92±0.18	100	93±6	14±1.0	37.8±1.3
10			8	7.68±0.42	10	8±0.7	13±2.0	16.7±0.28
11			8	7.70±0.52	100	91±5	13±1.5	39.0±0.8
12								

Table 2. Water quality data for experimental media. Mean ± standard error (n=30; \*n=15).

## Results

### Water Quality

Nominal and actual quality data for the experimental media are given in Table 2. The actual concentrations of Al and Ca were slightly lower than the nominal values. The actual pH values in all aquariums were quite close to the nominal values during the experiments.

### Mortality, Behaviour and Appearance of the Larvae

At the end of the experiments, mortality was less than 17 % in all animals except in those exposed to combinations of aluminium (8  $\mu\text{mol/L}$ ) and pH (4.5 and 7.0) at low calcium (10  $\mu\text{mol/L}$ ) (mortality higher than 27%).

In the absence of aluminium, larvae had nearly absorbed the yolk after 30 days. They were also very active and reached the swim-up stage. However, in the presence of aluminium, larvae still possessed substantial yolk reserves as a result of retardation of the resorption of yolk. They were also less active and never reached the swim-up stage, especially at low pH (4.5) and low calcium (10  $\mu\text{mol/L}$ ).

### Dry Weight and Whole Body Mineral Content of the Larvae

In general, calcium, potassium and sodium concentrations in the larvae were higher at the end of the experiment (especially in the control) than at the beginning (see Figures 1, 3 and 4), while magnesium concentration remained approximately constant or slightly lower (see Figure 2).

Whole body calcium content declined significantly ( $P < 0.001$ ) with increasing aluminium concentration at all combinations of pH and low calcium concentration (10  $\mu\text{mol/L}$ ). The effect of aluminium was less severe ( $P < 0.01$ ) at the higher calcium concentration (100  $\mu\text{mol/L}$ ) (see Figure 1). There was little difference in the effect of aluminium between the two pH regimes. In the absence of aluminium, body calcium content was significantly ( $P < 0.001$ ) lower at pH 4.5 at both calcium concentrations.

Whole body magnesium content declined as aluminium concentration increased (Figure 2). The analysis of variance indicated that the effect of aluminium was significant ( $P < 0.001$ ). However, there were no significant effects of pH or calcium concentration.




Figure 1. Whole body calcium concentration of the larvae at the end of the experiment. (a) pH 7.0; (b) pH 4.5. ● whole body Ca concentration of the newly hatched yolk-sac fry




Figure 2. Whole body magnesium concentration of the larvae at the end of the experiment. (a) pH 7.0; (b) pH 4.5. ● whole body Mg concentration of the newly hatched yolk-sac fry

Whole body potassium content declined significantly ( $P < 0.001$ ) with increasing aluminium concentration at all combinations of pH and low calcium concentration (10 µmol/L). The effect of aluminium was less severe ( $P < 0.01$ ) at the higher calcium concentration (100 µmol/L) (Figure 3). There was little difference in the effect of aluminium between the two pH regimes. In the absence of aluminium, there was no significant difference attributable to pH or to calcium concentrations.

Whole body sodium content declined significantly ( $P < 0.001$ ) as the aluminium concentration increased (Figure 4). Analysis of variance of the results from treatments without aluminium indicated that the effect of pH was less significant ( $P < 0.01$ ). Body sodium content was lower at the lower calcium concentration (10 µmol/L) at each combination of pH and aluminium concentration. This effect was more obvious at lower pH (4.5).




Figure 3. Whole body potassium concentration of the larvae at the end of the experiment. (a) pH 7.0; (b) pH 4.5. ● whole body K concentration of the newly hatched yolk-sac fry




Figure 4. Whole body sodium concentration of the larvae at the end of the experiment. (a) pH 7.0; (b) pH 4.5. ● whole body Na concentration of the newly hatched yolk-sac fry

## Discussion

The recently hatched brown trout fry in this study supplemented their calcium, potassium and sodium content by net uptake from the water, whereas there was no net magnesium uptake. A relationship was observed between whole body calcium content and the net uptake of calcium, potassium and sodium (but not magnesium) in Atlantic salmon, *Salmo salar*, alevins (4) and in brown trout alevins (5, 6).

The results of the present study showed that the effects of low pH alone were less than the effects of aluminium and pH combinations, whereas a number of reports of laboratory experiments with salmonid alevins indicated severe effect at pH levels relevant to the present study (7-10).

It was observed that water calcium concentration plays an important role in the effects of pH and aluminium. The deleterious effects of combinations of pH and aluminium increased with decreasing water calcium level. In previous reports, impaired skeletal calcium deposition (10) and impaired otolith growth (11) were found in salmonids at low pH and low calcium levels.

The form in which aluminium is present in water is highly pH-dependent. Its solubility decreases to a minimum at around pH 5.5-6.0 and increases towards both extremes of the pH scale. Soluble, monomeric, cationic species, e.g.  $\text{Al}^{3+}$ ,  $\text{Al}(\text{OH})^{+2}$  and  $\text{Al}(\text{OH})_2^+$ , are formed at pH levels lower than about 5.5, and soluble aluminate species, e.g.  $\text{Al}(\text{OH})_4^-$ , predominate at pH levels

greater than about pH 5.5. Between pH 5.5 and 6.0 most aluminium is present in the form of insoluble  $\text{Al}(\text{OH})_3$  (12, 13).

Aluminium toxicity is generally regarded as an important factor in the decline of fish population in soft acidic water. Toxicity is modified by pH and tends to be greater at lower calcium concentrations (14). The deleterious effects of aluminium observed as mortality, impaired potassium, sodium and calcium uptake in the present study were all ameliorated by high calcium concentration (100 compared with 10  $\mu\text{mol/L}$ ) and the retarded gross development and impaired uptake of potassium were ameliorated by high pH (7.0) compared with 4.5. The slight increase in the net loss of magnesium caused by exposure to aluminium was not significantly modified either by calcium concentration or by pH.

High mortality (> 27%) occurred only at aluminium concentration greater than 4  $\mu\text{mol/L}$ , in the low calcium concentration (10  $\mu\text{mol/L}$ ). Lower concentrations of aluminium are lethal to older juvenile brown trout (15) and rainbow trout, *Salmo gairdneri* (3). However, yolk-sac fry are less sensitive to aluminium than older animals (14, 16). Neville (3) concluded that the cause of death in aluminium exposed rainbow trout was predominantly disturbance of the sodium and chloride balance below pH 5 and hypoxia at pH 6.1, with a transition between two mechanisms around pH 5.0-5.5. Active uptake of sodium from the medium is inhibited by aluminium in yearling brown trout below pH 5.0 (17). At this pH, aluminium is predominantly in the form of monomeric cations, and

these cations are presumably responsible for the impairment of the sodium and chloride balance. This sort of toxicity is similar to that of hydrogen ions alone at very low pH, in which death probably results from circulation failure brought about by ionoregulatory disturbances (18). Net sodium uptake in the present study was more severely impaired at pH 4.5 than at pH 7.0. Net uptake of calcium was impaired by aluminium, whereas active uptake of calcium by brown trout fingerlings is not inhibited under similar conditions in short-term experiments (19).

## References

1. Sayer, M. D. J., Reader, J. P. and Morris, R., Embryonic and larval development of brown trout exposed to aluminium, copper, lead and zinc in soft acid water. *J. Fish Biol.* 38: 431-455, 1991.
2. Muniz, I. P. and Leivestad, H., Toxic effects of aluminium on the brown trout, *Salmo trutta* L. In Proc. Int. Conf. Ecological Impact of Acid Precipitation (eds. D. Drablos and A. Tollan), pp. 320-321, Oslo-As, Norway: SNSF Project, 1980.
3. Neville, C. M., Physiological response of juvenile rainbow trout, *Salmo gairdneri*, to acid and aluminium-prediction of field responses from laboratory data. *Can. J. Fish. Aquat. Sci.* 42: 2004-2019, 1985.
4. Rombough, P. J. and Garside, E. T., Disturbed ion balance in alevins of Atlantic salmon, *Salmo salar*, chronically exposed to sublethal concentration of cadmium. *Can. J. Zool.* 62: 1443-1450, 1984.
5. Runn, P. and Sohtell, M., Sodium influx and transepithelial potential in brown trout (*Salmo trutta*) embryos and yolk sac fry neutral and low pH. Acta Univ. Uppsala, Abstract of Uppsala Dissertations from the Faculty of Science 647 (IV), 1982a.
6. Runn, P. and Sohtell, M., Sodium and potassium accumulation in brown trout (*Salmo trutta* L.) embryos and yolk sac fry incubated at neutral and low pH. Acta Univ. Uppsala, Abstract of Uppsala Dissertations from the Faculty of Science 647 (V), 1982b.
7. Daye, P. G. and Garside, E. T., Development and survival of embryos and alevins of the Atlantic salmon, *Salmo salar* L. induced by continuous or short-term exposure to acidic levels of pH, from fertilisation. *Can. J. Zool.* 58: 27-43, 1979.
8. Johansson, N., Runn, P. and Milbrink, G., Early development of three salmonid species in acidified water. *Zoon*, 5: 127-132, 1977.
9. Brown, D. J. A., Influence of calcium on the survival of eggs and fry of brown trout (*Salmo trutta*) at pH 4.5. *Bull. Envir. Contam. Toxicol.* 28: 664-668, 1982.
10. Nelson, J. A., Physiological observations on developing rainbow trout, *Salmo gairdneri* (Richardson), exposed to low pH and varied calcium ion concentrations. *J. Fish Biol.* 20: 359-372, 1982.
11. Green, G. H., Neilson, J. D. and Bradford, M., Effects of pH on the early development and growth and otolith microstructure of chinook salmon, *Oncorhynchus tshawytscha*. *Can. J. Zool.* 63: 22-27, 1985.
12. Burrows, W. D., Aquatic aluminium: chemistry, toxicology and environmental prevalence. *Crit. Rev. Envir. Control* 7: 167-216, 1977.
13. Spry, D. J., Wood, C. M. and Hodson, P. B., The effects of environmental acid on freshwater fish with particular reference to soft water lakes in Ontario and the modifying effects of heavy metals. A literature review. *Can. Tech. Rep. Fish. Aquat. Sci.* 999, 145 pp, 1981.
14. Brown, D. J. A., Effect of calcium and aluminium concentrations on the survival of brown trout (*Salmo trutta*) at low pH. *Bull. Envir. Contam. Toxicol.* 30: 582-587, 1983.
15. Sadler, K. and Lynam, S., Some effects on the growth of brown trout from exposure to aluminium at different pH levels. *J. Fish Biol.* 31: 209-219, 1987.
16. Baker, J. P. and Schofield, C. L., Aluminium toxicity to fish in acidic waters. *Water Air and Soil Pollution*. 18: 289-309, 1982.
17. Dalziel, T. R. K., Morris, R. and Brown, D. J. A., The effects of low pH, low calcium concentrations and elevated aluminium concentrations on sodium fluxes in brown trout, *Salmo trutta* L. *Water, Air and Soil Pollution* 30: 569-577, 1986.
18. Milligan, C. L. and Wood, C. M., Disturbances in haematology, fluid volume distribution and circulatory function associated with low environmental pH in the rainbow trout, *Salmo gairdneri*. *J. Exp. Biol.* 99: 397-415, 1982.
19. Reader, J. P., Effects of cadmium, manganese and aluminium in soft acid water on ion regulation in *Salmo trutta* L. Ph. D. thesis, University of Nottingham, 1986.