

1-1-2003

On Some Properties of Szasz-Mirakyan Operators in Hölder Spaces

ZBIGNIEW WALCZAK

LUCYNA REMPULSKA

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

WALCZAK, ZBIGNIEW and REMPULSKA, LUCYNA (2003) "On Some Properties of Szasz-Mirakyan Operators in Hölder Spaces," *Turkish Journal of Mathematics*: Vol. 27: No. 3, Article 7. Available at: <https://journals.tubitak.gov.tr/math/vol27/iss3/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

On Some Properties of Szasz-Mirakyan Operators in Hölder Spaces

L. Rempulska, Z. Walczak

Abstract

We study some properties of modified Szasz-Mirakyan operators in Hölder exponential weighted spaces. We give theorems on the degree of approximation of functions by these operators.

Key Words: Szasz-Mirakyan operator, Hölder space, modulus of smoothness, degree of approximation.

1. Introduction

1.1. Paper [1] examined approximation properties of Szasz-Mirakyan operators

$$S_n(f; x) := \sum_{k=0}^{\infty} p_k(nx) f\left(\frac{k}{n}\right), \quad (1)$$

$x \in R_0 = [0, +\infty)$, $n \in N := \{1, 2, \dots\}$, where

$$p_k(t) := e^{-t} \frac{t^k}{k!} \quad \text{for } t \in R_0, k \in N_0 := N \cup \{0\}, \quad (2)$$

in exponential weighted spaces C_q . The space C_q , with a fixed $q > 0$, is related with the weighted function $v_q(x) := e^{-qx}$, $x \in R_0$, and C_q is the set of all real-valued functions f

A.M.S. Subject Classification: 41A36

continuous on R_0 for which $v_q f$ is uniformly continuous and bounded on R_0 . The norm in C_q is defined by

$$\|f\|_q \equiv \|f(\cdot)\|_q := \sup_{x \in R_0} v_q(x)|f(x)|. \tag{3}$$

It is obvious that $C_q \subset C_p$ if $0 < q < p < +\infty$.

In [1] was proved that S_n defined by (1) is a positive linear operator from the space C_q into C_p provided that $p > q > 0$ and $n > q/\ln(p/q)$.

Recently in many papers were introduced various modifications of operators S_n (see e.g. [3, 4, 5, 8, 10]).

In the paper [8] were introduced for $f \in C_q$, $q > 0$, the following modified Szasz-Mirakyan operators

$$S_{n,q}(f; x) := \sum_{k=0}^{\infty} p_k(nx) f\left(\frac{k}{n+q}\right), \quad x \in R_0, \quad n \in N, \tag{4}$$

where $p_k(\cdot)$ is defined by (2). Also in [8] was proved that, for every $n \in N$ and $q > 0$, $S_{n,q}$ is positive linear operator from the space C_q into C_q and

$$\|S_{n,q}(f; \cdot)\|_q \leq \|f\|_q, \quad n \in N, \tag{5}$$

for every $f \in C_q$. Moreover, in [8] were given approximation theorems for $f \in C_q$ and $S_{n,q}(f)$.

1.2. The purpose of this paper is the examination of approximation properties of operators $S_{n,q}$ in Hölder spaces related with exponential weighted space C_q , $q > 0$.

Approximation of 2π -periodic functions in the Hölder spaces first was considered by S. Prössdorf and J. Prestin in the papers [6] and [7].

The results given in [6] and [7] were extend by many authors.

Since the Szasz-Mirakyan operators are important in approximation theory and $S_{n,q}$ are operators from the space C_q into C_q , we shall consider the properties of these operators in the Hölder spaces.

In this paper, as had been similarly done in [6] and [7], we shall apply the modulus of smoothness of the order $r \in N$ ([2], [9]) of function $f \in C_q$:

$$\omega_r(f; q; t) := \sup_{0 \leq h \leq t} \|\Delta_h^r f(\cdot)\|_q, \quad t \geq 0, \tag{6}$$

where

$$\Delta_h^1 f(x) := f(x+h) - f(x), \quad \Delta_h^r f(x) := \Delta_h^1 (\Delta_h^{r-1} f(x)) \quad \text{if } r \geq 2, \quad (7)$$

for $x, h \in R_0$. It is known ([9]) that

$$\Delta_h^r f(x) = \sum_{j=0}^r \binom{r}{j} (-1)^{r-j} f(x+jh) \quad \text{for } x, h \in R_0 \text{ and } r \in N. \quad (8)$$

Moreover from (6) – (8) we deduce that if $f \in C_q$, $q > 0$, and $r \in N$, then $\omega_r(f, q; \cdot)$ is non-negative and non-decreasing function and $\lim_{t \rightarrow 0+} \omega_r(f; q; t) = 0$.

1.3. Let $r \in N$ and let Ω_r be the set of functions of $\omega_r(f; q; \cdot)$ type, i.e. Ω_r is the set of all functions ω satisfying the following conditions:

- (i) ω is defined, non-negative and increasing on R_0 ;
- (ii) $\omega(t) \rightarrow 0$ as $t \rightarrow 0+$;
- (iii) $\omega(t)t^{-r}$ is decreasing for $t > 0$.

Similarly as in [6] and [7], for given $r \in N$, $\omega \in \Omega_r$ and $q > 0$, we define the generalized Hölder spaces $H_q^{r,\omega}$ and $\tilde{H}_q^{r,\omega}$. The space $H_q^{r,\omega}$ is the set of all functions $f \in C_q$ for which

$$\|f\|_q^{*r,\omega} := \sup_{h>0} \frac{\|\Delta_h^r f(\cdot)\|_q}{\omega(h)} < +\infty \quad (9)$$

and the norm is defined by the formula

$$\|f\|_{H_q^{r,\omega}} := \|f\|_q + \|f\|_q^{*r,\omega}. \quad (10)$$

The space $\tilde{H}_q^{r,\omega}$ is the set of all functions $f \in H_q^{r,\omega}$ for which

$$\lim_{t \rightarrow 0+} \frac{\omega_r(f; q; t)}{\omega(t)} = 0 \quad (11)$$

and the norm is defined by (10).

From definition of $H_q^{r,\omega}$ we deduce that $f \in H_q^{r,\omega}$ if and only if there exists a positive constant $M_1(f)$ depending only on f and such that

$$\omega_r(f, q; t) \leq M_1(f)\omega(t) \quad \text{for } t \geq 0. \tag{12}$$

The spaces $H_q^{r,\omega}$ and $\tilde{H}_q^{r,\omega}$, with fixed $q > 0$, $r \in N$ and $\omega(t) = t^\alpha$, $0 < \alpha \leq r$, are classical Hölder - Lipschitz - Zygmund spaces. Moreover, we observe that if $\omega, \mu \in \Omega_r$, $r \in N$, and

$$\lambda(t) := \frac{\omega(t)}{\mu(t)}, \quad t > 0, \tag{13}$$

is increasing function, then for every $q > 0$ we have

$$H_q^{r,\omega} \subset H_q^{r,\mu}, \quad \tilde{H}_q^{r,\omega} \subset \tilde{H}_q^{r,\mu} \tag{14}$$

for every $q > 0$.

2. Main Results

2.1. First we shall give certain inequalities for operators $S_{n,q}$ and $f \in C_q$.

Denote by C_q^p , $p \in N$, $q > 0$, the class of all functions $f \in C_q$ which have derivatives $f^{(k)}$, $k = 1, \dots, p$, on R_0 and these $f^{(k)}$ belong also to C_q .

Theorem 1 *Let $n \in N$ and $q > 0$ be fixed numbers. Then $S_{n,q}$ defined by (4) is an operator from the space C_q into C_q^∞ . Moreover, for every $p \in N$ and $f \in C_q$, we have*

$$\| [S_{n,q}(f)]^{(p)} \|_q \leq (1 + e)^p n^p \|f\|_q. \tag{15}$$

If $f \in C_q^p$, $p \in N$, $q > 0$, then

$$\| [S_{n,q}(f)]^{(p)} \|_q \leq e^p \|f^{(p)}\|_q, \quad n \in N. \tag{16}$$

Proof. From (4), (2) and (7) we derive the formula

$$[S_{n,q}(f; x)]^{(p)} = n^p S_{n,q} \left(\Delta_{1/(n+q)}^p f(t); x \right),$$

for every $f \in C_q$ and $n, p \in N$. By (3) and (5) we have

$$\| [S_{n,q}(f)]^{(p)} \|_q \leq n^p \| \Delta_{1/(n+q)}^p f(\cdot) \|_q, \quad n, p \in N. \tag{17}$$

Applying (8) and (3), we get

$$\begin{aligned} \|\Delta_{1/(n+q)}^p f(\cdot)\|_q &\leq \|f\|_q \sum_{k=0}^p \binom{p}{k} e^{kq/(n+q)} \leq \\ &\leq \|f\|_q (1+e)^p, \quad n, p \in N. \end{aligned}$$

From the above follows (15).

It is known ([2], [9]) that if $f \in C_q^p$ with fixed $p \in N$ and $q > 0$, then

$$\Delta_h^p f(x) = \int_0^h \dots \int_0^h f^{(p)}(x + u_1 + \dots + u_p) du_1 \dots du_p, \quad h > 0,$$

which by (3) implies

$$\|\Delta_h^p f(\cdot)\|_q \leq \|f^{(p)}\|_q e^{hpq} h^p, \quad h > 0.$$

From this and by (17) we get

$$\|[S_{n,q}(f)]^{(p)}\|_q \leq \left(\frac{n}{n+q}\right)^p e^{pq/(n+q)} \|f^{(p)}\|_q \leq e^p \|f^{(p)}\|_q, \quad n, p \in N.$$

Thus the proof is completed. □

2.2. Let $f \in C_q$, $q > 0$, and let $r \in N$. It is known ([2], Section 6.1) that the modulus of smoothness $\omega_r(f; q; \cdot)$ defined by (6) is equivalent to the weighted K -functional

$$K_r(f; q; t^r) := \inf_{\varphi \in C_q^r} \left\{ \|f - \varphi\|_q + t^r \|\varphi^{(r)}\|_q \right\}, \quad (18)$$

i.e. there exists a positive constant M independent on f and t such that

$$M^{-1} \omega_r(f; q; t) \leq K_r(f; q; t^r) \leq M \omega_r(f; q; t) \quad \text{for } t > 0. \quad (19)$$

Using this equivalence, we shall prove theorem on modulus of smoothness of $S_{n,q}(f)$.

Theorem 2 *Let $q > 0$ and $r \in N$ be fixed numbers. Then there exists the suitable positive constant $M_2(r)$ depending only on r and such that for every $f \in C_q$ and $n \in N$ we have*

$$\omega_r(S_{n,q}(f); q; t) \leq M_2(r)\omega_r(f; q; t), \quad t \geq 0. \quad (20)$$

Proof. By Theorem 1 we have $S_{n,q}(f) \in C_q^\infty$ if $f \in C_q$. Hence for $S_{n,q}(f)$ we can apply K -functional (18) and the inequality (19). For given $f \in C_q$ and $n, r \in N$ we have

$$\begin{aligned} \omega_r(S_{n,q}(f); q; t) &\leq MK_r(S_{n,q}(f); q; t^r) = \\ &= M \inf_{\varphi \in C_q^r} \left\{ \|S_{n,q}(f) - \varphi\|_q + t^r \|\varphi^{(r)}\|_q \right\} \leq \\ &\leq M \left\{ \|S_{n,q}(f) - S_{n,q}(\Psi)\|_q + t^r \| [S_{n,q}(\Psi)]^{(r)} \|_q \right\} \end{aligned}$$

for $t \geq 0$, where Ψ is arbitrary fixed function in C_q^r .

By (4) and (5) we have

$$\|S_{n,q}(f) - S_{n,q}(\Psi)\|_q = \|S_{n,q}(f - \Psi; \cdot)\|_q \leq \|f - \Psi\|_q$$

for all $n \in N$. Moreover by Theorem 1, we have

$$\| [S_{n,q}(\Psi)]^{(r)} \|_q \leq e^r \|\Psi^{(r)}\|_q, \quad n \in N.$$

From the above we get

$$\omega_r(S_{n,q}(f); q; t) \leq Me^r \left\{ \|f - \Psi\|_q + t^r \|\Psi^{(r)}\|_q \right\}, \quad t \geq 0,$$

for every fixed $f \in C_q$, $n, r \in N$ and for every $\Psi \in C_q^r$, which by (18) and (19) yields

$$\begin{aligned} \omega_r(S_{n,q}(f); q; t) &\leq Me^r \inf_{\Psi \in C_q^r} \left\{ \|f - \Psi\|_q + t^r \|\Psi^{(r)}\|_q \right\} \leq \\ &\leq Me^r K_r(f; q; t^r) \leq M^2 e^r \omega_r(f; q; t) \end{aligned}$$

for $t \geq 0$, $f \in C_q$ and $n, r \in N$. Thus the inequality (20) was proved. \square

2.3. Now we shall give the main property of $S_{n,q}$ in Hölder spaces.

Theorem 3 *We assume that $q > 0$ and $r \in N$ are fixed numbers and $\omega \in \Omega_r$ is a given function. Then $S_{n,q}$, $n \in N$, defined by (4) is a positive linear operator from the space $H_q^{r,\omega}(\tilde{H}_q^{r,\omega})$ into $H_q^{r,\omega}(\tilde{H}_q^{r,\omega})$.*

Proof. Let $f \in H_q^{r,\omega}$ and $n \in N$. By Theorem 1, (20) and (12) we have $S_{n,q}(f) \in C_q^\infty$ and

$$\omega_r(S_{n,q}(f); q; t) \leq M_3(f, r)\omega(t), \quad t \geq 0, \quad (21)$$

where $M_3(f, r) = \text{const.} > 0$ depending only on f and r . The inequality (21) and Section 1.3 show that if $f \in H_q^{r,\omega}$ then also $S_{n,q}(f) \in H_q^{r,\omega}$.

Now, assume that $f \in \widetilde{H}_q^{r,\omega}$. The condition (11) and the inequality (20) imply that

$$\lim_{t \rightarrow 0^+} \frac{\omega_r(S_{n,q}(f); q; t)}{\omega(t)} = 0, \quad n \in N,$$

which proves that $S_{n,q}(f) \in \widetilde{H}_q^{r,\omega}$, $n \in N$. □

2.4. In [8] (p.127) was proved the following theorem.

Theorem 4 *If $f \in C_q^2$, $q > 0$, then*

$$\begin{aligned} v_q(x)|S_{n,q}(f; x) - f(x)| &\leq \|f'\|_q \frac{qx}{n+q} + \\ &+ \|f''\|_q \left(\frac{(4e^2 + 1)q^2x^2}{(n+q)^2} + \frac{4x}{n+q} \right) \end{aligned} \quad (22)$$

for $x \in R_0$ and $n \in N$.

Arguing as in [1] and [8], we shall prove the following theorem.

Theorem 5 *Suppose that $a > 0$ and $q > 0$ are fixed numbers. Then there exists a positive constant $M_4(a, q)$ (depending only on a and q) such that*

$$\sup_{0 \leq x \leq a} v_q(x)|S_{n,q}(f; x) - f(x)| \leq M_4(a, q)\omega_2(f; q; 1/\sqrt{n+q}) \quad (23)$$

for every $f \in C_q$ and $n \in N$.

Proof. Analogously as [1] and [8] we use the Steklov function f_h of $f \in C_q$:

$$f_h(x) := \frac{4}{h^2} \int_0^{\frac{h}{2}} \int_0^{\frac{h}{2}} [2f(x+s+t) - f(x+2(s+t))] ds dt$$

for $x \in R_0$ and $h > 0$. It is known ([1], [8]) that $f_h \in C_q^2$ if $f \in C_q$ and

$$\|f_h - f\|_q \leq \omega_2(f; q; h); \tag{24}$$

$$\|f'_h\|_q \leq 5e^{qh}h^{-1}\omega_1(f; q; h); \tag{25}$$

$$\|f''_h\|_q \leq 9h^{-2}\omega_2(f; q; h); \tag{26}$$

for $h > 0$. Applying (5), (22), and (24)–(26), we get

$$\begin{aligned} & v_q(x) |S_{n,q}(f; x) - f(x)| \leq v_q(x) \{ |S_{n,q}(f - f_h; x)| + \\ & + |S_{n,q}(f_h; x) - f_h(x)| + |f_h(x) - f(x)| \} \leq 2\|f_h - f\|_q + \\ & + \|f'_h\|_q \frac{qx}{n+q} + \|f''_h\|_q \left(\frac{(4e^2 + 1)q^2x^2}{(n+q)^2} + \frac{4x}{n+q} \right) \leq \\ & \leq \frac{5qe^{qh}x}{n+q}\omega_1(f; q; h) + \left\{ 2 + 9h^{-2} \left(\frac{(4e^2 + 1)q^2x^2}{(n+q)^2} + \frac{4x}{n+q} \right) \right\} \omega_2(f; q; h), \end{aligned}$$

for $x \in R_0$, $n \in N$ and $h > 0$. Hence, for fixed $a > 0$, $n \in N$ and $h = 1/\sqrt{n+q}$, we obtain

$$\begin{aligned} \sup_{0 \leq x \leq a} v_q(x) |S_{n,q}(f; x) - f(x)| & \leq M_5(a, q) \left\{ \frac{1}{\sqrt{n+q}} \omega_1 \left(f; q; \frac{1}{\sqrt{n+q}} \right) + \right. \\ & \left. + \omega_2 \left(f; q; \frac{1}{\sqrt{n+q}} \right) \right\}, \end{aligned}$$

which implies the desired estimation (23). □

2.5. Applying Theorem 5 we shall give two approximation theorems for function f belonging to Hölder spaces $H_q^{2,\omega}$ and $\tilde{H}_q^{2,\omega}$. For $f \in H_q^{2,\omega}$ with fixed $q > 0$, $\omega \in \Omega_2$ and for a fixed $a > 0$ we write

$$\|f\|_{q,a} := \sup_{0 \leq x \leq a} v_q(x) |f(x)|, \tag{27}$$

$$\|f\|_{q,a}^{*2,\omega} := \sup_{0 < h \leq 1} \frac{\|\Delta_h^2 f(\cdot)\|_{q,a}}{\omega(h)}, \tag{28}$$

$$\|f\|_{q,a}^{2,\omega} := \|f\|_{q,a} + \|f\|_{q,a}^{*2,\omega}. \quad (29)$$

Theorem 6 *Suppose that $a, q > 0$ are fixed numbers, $\omega, \mu \in \Omega_2$ are given functions such that $\lambda(\cdot)$ defined by (13) is increasing and $f \in H_q^{2,\omega}$. Then there exists a positive constant $M_6 \equiv M_6(a, q, f, \mu(1))$ (depending only on indicated parameters) such that*

$$\|S_{n,q}(f; \cdot) - f(\cdot)\|_{q,a}^{2,\mu} \leq M_6 \lambda(1/\sqrt{n+q}), \quad n \in N. \quad (30)$$

Proof. By our assumptions follows (14) and by Theorem 3 and (29) we have

$$\|S_{n,q}(f; \cdot) - f(\cdot)\|_{q,a}^{2,\mu} = \|S_{n,q}(f; \cdot) - f(\cdot)\|_{q,a} + \|S_{n,q}(f; \cdot) - f(\cdot)\|_{q,a}^{*2,\mu}$$

for $n \in N$. But by (27) and (23) and (12) we get

$$\begin{aligned} \|S_{n,q}(f; \cdot) - f(\cdot)\|_{q,a} &\leq M_4(a, q) \omega_2(f; q; 1/\sqrt{n+q}) \leq \\ &\leq M_7(a, q, f) \omega(1/\sqrt{n+q}) \leq M_7(a, q, f) \mu(1) \lambda(1/\sqrt{n+q}), \quad n \in N. \end{aligned}$$

Using definition (28) and denoting by

$$A = \{h : 0 < h \leq 1/\sqrt{n+q}\}, \quad B = \{h : 1/\sqrt{n+q} < h \leq 1\},$$

we can write

$$\|S_{n,q}(f; \cdot) - f(\cdot)\|_{q,a}^{*2,\mu} \leq \left(\sup_{h \in A} + \sup_{h \in B} \right) \frac{\|\Delta_h^2 [S_{n,q}(f; \cdot) - f(\cdot)]\|_{q,a}}{\mu(h)} := W_1 + W_2.$$

By (27), (3), (7), (6) and Theorem 2 we deduce that

$$\begin{aligned} \|\Delta_h^2 [S_{n,q}(f; \cdot) - f(\cdot)]\|_{q,a} &\leq \|\Delta_h^2 [S_{n,q}(f; \cdot) - f(\cdot)]\|_q \leq \\ &\leq \|\Delta_h^2 S_{n,q}(f; \cdot)\|_q + \|\Delta_h^2 f(\cdot)\|_q \leq \\ &\leq \omega_2(S_{n,q}(f); q; h) + \omega_2(f; q; h) \leq M_8 \omega_2(f; q; h) \end{aligned}$$

for $h > 0$. From this and by (12) we get

$$W_1 \leq M_9(f) \sup_{h \in A} \frac{\omega(h)}{\mu(h)} \leq M_9(f) \lambda(1/\sqrt{n+q}), \quad n \in N.$$

Applying (8) and (27) we get $\|\Delta_h^2 f(\cdot)\|_{q,a} \leq M_{10}(a, q)\|f\|_{q,a}$ and

$$W_2 \leq M_{10}(a, q) \sup_{h \in B} \frac{\|S_{n,q}(f; \cdot) - f(\cdot)\|_{q,a}}{\mu(h)},$$

which by Theorem 5 and (12) implies

$$\begin{aligned} W_2 &\leq M_{11}(a, q)\omega_2(f; q; 1/\sqrt{n+q}) / \mu(1/\sqrt{n+q}) \leq \\ &\leq M_{12}(a, q, f)\lambda(1/\sqrt{n+q}), \quad n \in N. \end{aligned}$$

Combining these, we obtain estimation (30). □

Analogously we can prove the following theorem

Theorem 7 *Suppose that a, q, ω, μ and λ satisfy the assumptions of Theorem 6. If $f \in \tilde{H}_q^{2,\omega}$, then*

$$\|S_{n,q}(f; \cdot) - f(\cdot)\|_{q,a}^{2,\mu} = o(\lambda(1/\sqrt{n+q})) \quad \text{as } n \rightarrow \infty.$$

From Theorem 6 and Theorem 7 we derive the following corollary.

Corollary. Let $a, q > 0$ be fixed numbers and let $\omega(t) = t^\alpha$, $\mu(t) = t^\beta$ for $t \geq 0$ and for fixed $0 < \beta < \alpha \leq 2$.

1. If $f \in H_q^{2,\omega}$, then

$$\|S_{n,q}(f; \cdot) - f(\cdot)\|_{q,a}^{2,\mu} = O\left((n+q)^{(\beta-\alpha)/2}\right) \quad \text{as } n \rightarrow \infty.$$

2. If $f \in \tilde{H}_q^{2,\omega}$, then

$$\|S_{n,q}(f; \cdot) - f(\cdot)\|_{q,a}^{2,\mu} = o\left((n+q)^{(\beta-\alpha)/2}\right) \quad \text{as } n \rightarrow \infty.$$

Remark. Analogous theorems we can obtain for operators of Kantorovitch type:

$$T_{n,q}(f; x) := \sum_{k=0}^{\infty} p_k(nx)(n+q) \int_{k/(n+q)}^{(k+1)/(n+q)} f(t)dt, \quad x \in R_0, \quad n \in N,$$

which are positive linear operators from the space C_q into C_q and $\|T_{n,q}\|_q \leq e\|f\|_q$ for every $f \in C_q$ and $n \in N$.

Similar theorems can also be obtained for Szasz-Mirakyan operators (1) in the Hölder polynomial weighted spaces related with the polynomial weighted spaces C_p , $p \in N_0$, and the weighted function $w_p(x) := 1/(1+x^p)$ if $p \geq 1$, and $w_0(x) := 1$ for $x \in R_0$.

Cleary, approximation properties of other operators (e.g. operators considered in [3, 4, 10]) can be examined in suitable Hölder spaces.

References

- [1] M. Becker, D. Kucharski, R.J. Nessel, Global approximation theorems for the Szasz-Mirakjan operators in exponential weight spaces, In: Linear Spaces and Approximation (Proc. Conf. Oberwolfach, 1977), Birkhuser Verlag, Basel ISNM, 40(1978), 319 - 333.
- [2] Z. Ditzian, V. Totik, Moduli of Smoothness, Springer Series in Computational Mathematics, Vol. 9, Springer-Verlag, New York Inc. 1987.
- [3] V. Gupta, R.P. Pant, Rate of convergence of the modified Szasz-Mirakyan operators on functions of bounded variation, J. Math. Anal. Appl., 233(199), 476-483.
- [4] V. Gupta, G.S. Srivastava, Approximation by Durrmeyer type operators, Annal. Polonici Math., LXIV(1996), 153-159.
- [5] B. Firlej, L. Rempulska, Approximation of functions of several variables by some operators of Szasz-Mirakyan type, Fasciculi Mate., 27(1997), 15-28.
- [6] J. Prestin, S. Prössdorf, Error estimates in generalized trigonometric Hölder norms, Z. Anal. Anwendungen, 9(1990), 343-349.
- [7] S. Prössdorf, Zur konvergenz der Fourierreihen hölderstetiger funktionen, Math. Nachr., 69(1975), 7-14.
- [8] L. Rempulska, Z. Walczak, Approximation properties of certain modified Szasz-Mirakyan operators, Le Matematiche 55(2001)1, 121-132.
- [9] A.F. Timann, Theory of Approximation of Functions of a Real Variable, New York 1963.
- [10] [10] Z. Walczak, Approximation of functions of two variables by modified Szasz-Mirakyan operators, Fasciculi Math., 31(2001), 117-128.

REMPULSKA, WALCZAK

L. REMPULSKA, Z. WALCZAK
Institute of Mathematics
Poznań University of Technology
Piotrowo 3A
60-965 Poznań-POLAND
e-mail: zwalczak@math.put.poznan.pl

Received 10.12.2002