

1-1-2003

## On Pseudohyperbolic Curves in Minkowski Space-Time

ÇETİN CAMCI

KAZIM İLARSLAN

EMILIJA SUCUROVIC

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

CAMCI, ÇETİN; İLARSLAN, KAZIM; and SUCUROVIC, EMILIJA (2003) "On Pseudohyperbolic Curves in Minkowski Space-Time," *Turkish Journal of Mathematics*: Vol. 27: No. 2, Article 5. Available at: <https://journals.tubitak.gov.tr/math/vol27/iss2/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## On Pseudohyperbolic Curves in Minkowski Space-Time

*Çetin Camcı, Kazım İlarıslan and Emilija Šučurović*

### Abstract

In this paper, we characterize all spacelike, timelike and null curves lying on the pseudohyperbolic space  $H_0^3$  in the Minkowski space-time  $E_1^4$ . Moreover, we prove that there are no timelike and no null curves lying on the pseudohyperbolic space  $H_0^3$  in Minkowski space-time  $E_1^4$ .

**Key words and phrases:** Minkowski space-time, pseudohyperbolic space, curvature.

### 1. Introduction

A necessary and sufficient conditions for a curve to be a spherical curve in Euclidean space  $E^3$  are given in [6] and [7]. On the other hand, a similar characterizations of a spacelike, a timelike and a null curves lying on the pseudohyperbolic space  $H_0^2$  in the Minkowski space  $E_1^3$  are obtained in [4]. The corresponding Frenet's equations for an arbitrary curve in the Minkowski space-time  $E_1^4$ , are given in [5]. By using these equations, in this paper we give some necessary and sufficient conditions for a spacelike curve to lie on the pseudohyperbolic space  $H_0^3$  in Minkowski space-time.

## 2. Preliminaries

Minkowski space–time  $E_1^4$  is a Euclidean space  $E^4$  provided with the standard flat metric given by

$$g = -dx_1^2 + dx_2^2 + dx_3^2 + dx_4^2,$$

where  $(x_1, x_2, x_3, x_4)$  is a rectangular coordinate system in  $E_1^4$ .

Since  $g$  is an indefinite metric, recall that a vector  $v \in E_1^4$  can have one of three causal characters: it can be *spacelike* if  $g(v, v) > 0$  or  $v = 0$ , *timelike* if  $g(v, v) < 0$  and *null (lightlike)* if  $g(v, v) = 0$  and  $v \neq 0$ . Similarly, an arbitrary curve  $\alpha = \alpha(s)$  in  $E_1^4$  can locally be *spacelike*, *timelike* or *null (lightlike)*, if all of its velocity vectors  $\alpha'(s)$  are respectively *spacelike*, *timelike* or *null*. Also, recall that the norm of a vector  $v$  is given by  $\|v\| = \sqrt{|g(v, v)|}$ . Therefore,  $v$  is a unit vector if  $g(v, v) = \pm 1$ . Next, vectors  $v, w$  in  $E_1^4$  are said to be orthogonal if  $g(v, w) = 0$ . The velocity of the curve  $\alpha(s)$  is given by  $\|\alpha'(s)\|$ .

The pseudohyperbolic space with center  $m = (m_1, m_2, m_3, m_4) \in E_1^4$  and radius  $r \in R^+$  in the space–time  $E_1^4$  is the hyperquadric

$$H_0^3(r) = \{a = (a_1, a_2, a_3, a_4) \in E_1^4 \mid g(a - m, a - m) = -r^2\}.$$

with dimension 3 and index 0.

Denote by  $\{T(s), N(s), B_1(s), B_2(s)\}$  the moving Frenet frame along the curve  $\alpha(s)$  in the space  $E_1^4$ . Then  $T, N, B_1, B_2$  are, respectively, the tangent, the principal normal, the first binormal and the second binormal vector fields. Spacelike or timelike curve  $\alpha(s)$  is said to be parameterized by arclength function  $s$ , if  $g(\alpha'(s), \alpha'(s)) = \pm 1$ . In particular, a null curve  $\alpha(s)$  is said to be parameterized by a pseudo–arclength function  $s$  if  $g(\alpha''(s), \alpha''(s)) = 1$ , where pseudo–arclength function  $s$  is defined in [1] by  $s = \int_0^t (g(\alpha''(t), \alpha''(t)))^{\frac{1}{4}}$ .

Let  $\alpha(s)$  be a curve in the space–time  $E_1^4$ , parameterized by arclength function  $s$ . Then for the curve  $\alpha$  the following Frenet equations are given in [5] :

### Case 1. $\alpha$ is spacelike curve:

Then  $T$  is spacelike vector, so depending on the causal character of the principal normal vector  $N$ , we distinguish subcases 1.1, 1.2 and 1.3 as follows.

Case 1.1.  $N$  is spacelike:

Then the first binormal  $B_1$  can have all three causal characters, so we distinguish subcases 1.1.1, 1.1.2 and 1.1.3.

Case 1.1.1.  $B_1$  is spacelike:

In this case, the Frenet formulae is read

$$\begin{bmatrix} \dot{T} \\ \dot{N} \\ \dot{B}_1 \\ \dot{B}_2 \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ -k_1 & 0 & k_2 & 0 \\ 0 & -k_2 & 0 & k_3 \\ 0 & 0 & k_3 & 0 \end{bmatrix} \begin{bmatrix} T \\ N \\ B_1 \\ B_2 \end{bmatrix},$$

where  $T, N, B_1, B_2$  are mutually orthogonal vectors satisfying equations

$$g(T, T) = g(N, N) = g(B_1, B_1) = 1, g(B_2, B_2) = -1.$$

Case 1.1.2.  $B_1$  is timelike:

The Frenet formulae has the form

$$\begin{bmatrix} \dot{T} \\ \dot{N} \\ \dot{B}_1 \\ \dot{B}_2 \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ -k_1 & 0 & k_2 & 0 \\ 0 & k_2 & 0 & k_3 \\ 0 & 0 & k_3 & 0 \end{bmatrix} \begin{bmatrix} T \\ N \\ B_1 \\ B_2 \end{bmatrix},$$

where  $T, N, B_1, B_2$  are mutually orthogonal vectors satisfying equations

$$g(T, T) = g(N, N) = g(B_2, B_2) = 1, g(B_1, B_1) = -1.$$

Case 1.1.3.  $B_1$  is a null:

The Frenet formulae are given by

$$\begin{bmatrix} \dot{T} \\ \dot{N} \\ \dot{B}_1 \\ \dot{B}_2 \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ -k_1 & 0 & k_2 & 0 \\ 0 & 0 & k_3 & 0 \\ 0 & -k_2 & 0 & -k_3 \end{bmatrix} \begin{bmatrix} T \\ N \\ B_1 \\ B_2 \end{bmatrix},$$

where  $T, N, B_1, B_2$  satisfy equations

$$g(T, T) = g(N, N) = 1, g(B_1, B_1) = g(B_2, B_2) = 0, \\ g(T, N) = g(T, B_1) = g(T, B_2) = g(N, B_1) = g(N, B_2) = 0, g(B_1, B_2) = 1.$$

Case 1.2.  $N$  is timelike:

In this case, the Frenet equations has the form

$$\begin{bmatrix} \dot{T} \\ \dot{N} \\ \dot{B}_1 \\ \dot{B}_2 \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ k_1 & 0 & k_2 & 0 \\ 0 & k_2 & 0 & k_3 \\ 0 & 0 & -k_3 & 0 \end{bmatrix} \begin{bmatrix} T \\ N \\ B_1 \\ B_2 \end{bmatrix},$$

where  $T, N, B_1, B_2$  are mutually orthogonal vectors satisfying equations

$$g(T, T) = g(B_1, B_1) = g(B_2, B_2) = 1, g(N, N) = -1.$$

Case 1.3.  $N$  is a null:

Now the Frenet formulae read

$$\begin{bmatrix} \dot{T} \\ \dot{N} \\ \dot{B}_1 \\ \dot{B}_2 \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ 0 & 0 & k_2 & 0 \\ 0 & k_3 & 0 & -k_2 \\ -k_1 & 0 & -k_3 & 0 \end{bmatrix} \begin{bmatrix} T \\ N \\ B_1 \\ B_2 \end{bmatrix},$$

where the first curvature  $k_1$  can take only two values: 0 when  $\alpha$  is a straight line or 1 in all other cases. In this case, the vectors  $T, N, B_1, B_2$  satisfy equations

$$g(T, T) = g(B_1, B_1) = 1, g(N, N) = g(B_2, B_2) = 0, \\ g(T, N) = g(T, B_1) = g(T, B_2) = g(N, B_1) = g(B_1, B_2) = 0, g(N, B_2) = 1.$$

**Case 2.  $\alpha$  is a timelike curve:**

Then  $T$  is timelike vector, so the Frenet formulae has the form

$$\begin{bmatrix} \dot{T} \\ \dot{N} \\ \dot{B}_1 \\ \dot{B}_2 \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ k_1 & 0 & k_2 & 0 \\ 0 & -k_2 & 0 & k_3 \\ 0 & 0 & -k_3 & 0 \end{bmatrix} \begin{bmatrix} T \\ N \\ B_1 \\ B_2 \end{bmatrix},$$

where  $T, N, B_1, B_2$  are mutually orthogonal vectors satisfying equations

$$g(T, T) = -1, g(N, N) = g(B_1, B_1) = g(B_2, B_2) = 1.$$

**Case 3.  $\alpha$  is a null curve:**

Then  $T$  is null vector, so the Frenet equations are given by

$$\begin{bmatrix} \dot{T} \\ \dot{N} \\ \dot{B}_1 \\ \dot{B}_2 \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ k_2 & 0 & -k_1 & 0 \\ 0 & -k_2 & 0 & k_3 \\ -k_3 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} T \\ N \\ B_1 \\ B_2 \end{bmatrix},$$

where the first curvature  $k_1$  can take only two values: 0 when  $\alpha$  is a straight null line or 1 in all other cases. In this case, the vectors  $T, N, B_1, B_2$  satisfy the equations

$$\begin{aligned} g(T, T) &= g(N, N) = g(B_1, B_1) = 0, g(B_2, B_2) = 1, \\ g(T, N) &= g(T, B_2) = g(N, B_1) = g(N, B_2) = g(B_1, B_2) = 0, g(T, B_1) = 1. \end{aligned}$$

Recall that the functions  $k_1 = k_1(s)$ ,  $k_2 = k_2(s)$  and  $k_3 = k_3(s)$  are called respectively the first, the second and the third curvature of curve  $\alpha(s)$ .

**3. The spacelike, timelike and null curves lying on  $H_0^3$**

In this section, under the assumption that spacelike, timelike and null curves lie on the pseudohyperbolic space  $H_0^3(r)$ , we give some theorems characterizing these curves in terms of their three curvatures  $k_1(s)$ ,  $k_2(s)$  and  $k_3(s)$ .

**Theorem 3.1** *Let  $\alpha(s)$  be a unit speed spacelike curve in  $E_1^4$  with spacelike  $N, B_1$  and with curvatures  $k_1(s) \neq 0, k_2(s) \neq 0, k_3(s) \neq 0$  for each  $s \in I \subset R$ . Then  $\alpha$  lies on  $H_0^3(r)$  if and only if*

$$(1/k_1)^2 + ((1/k_2)(1/k_1)')^2 - [(1/k_3)(k_2/k_1 + ((1/k_2)(1/k_1)'))]^2 = -r^2. \tag{1}$$

**Proof.** Let us first suppose that  $\alpha$  lies on  $H_0^3(r)$  with center  $m$ . Then  $g(\alpha - m, \alpha - m) = -r^2$ , for each  $s \in I \subset R$ . Differentiating the previous equation four times with respect to

$s$  and by applying Frenet's equations, we get

$$\begin{aligned} g(T, \alpha - m) = 0, g(N, \alpha - m) = -1/k_1, g(B_1, \alpha - m) = -(1/k_2)(1/k_1)', \\ g(B_2, \alpha - m) = -(1/k_3)[k_2/k_1 + ((1/k_2)(1/k_1)')']. \end{aligned} \quad (2)$$

Let us decompose the vector  $\alpha - m$  with respect to the pseudo-orthonormal basis  $\{T, N, B_1, B_2\}$  by

$$\alpha(s) - m = a(s)T(s) + b(s)N(s) + c(s)B_1(s) + d(s)B_2(s), \quad (3)$$

where  $a(s), b(s), c(s), d(s)$  are arbitrary functions. Then we easily find that

$$g(T, \alpha - m) = a, g(N, \alpha - m) = b, g(B_1, \alpha - m) = c, g(B_2, \alpha - m) = -d. \quad (4)$$

Therefore, substituting (2) and (4) into (3) yields (1).

Conversely, let us assume that the relation (1) hold. By taking the derivative of (1) with respect to  $s$ , we find

$$\begin{aligned} (1/k_3)[k_2/k_1 + ((1/k_2)(1/k_1)')][(k_3/k_2)(1/k_1)' - ((1/k_3)(k_2/k_1) \\ + ((1/k_2)(1/k_1)')')] = 0. \end{aligned} \quad (5)$$

If there holds  $(1/k_3)[k_2/k_1 + ((1/k_2)(1/k_1)')'] = 0$ , then

substituting this into (1) yields a contradiction. Therefore, it follows that

$$(k_3/k_2)(1/k_1)' - [(1/k_3)(k_2/k_1 + ((1/k_2)(1/k_1)')')] = 0. \quad (6)$$

Next, we may consider the vector  $m \in E_1^4$  given by

$$m = \alpha + (1/k_1)N + (1/k_2)(1/k_1)'B_1 - (1/k_3)[k_2/k_1 + ((1/k_2)(1/k_1)')']B_2. \quad (7)$$

Differentiating (7) with respect to  $s$  and by using Frenet formulae, we obtain

$$m' = [(k_3/k_2)(1/k_1)' - ((1/k_3)(k_2/k_1 + ((1/k_2)(1/k_1)')')')]B_2. \quad (8)$$

Then substitution of (6) into (8) implies  $m' = 0$  and thus  $m = \text{constant}$ . Finally, from (7) we find  $g(\alpha - m, \alpha - m) = -r^2$ , so  $\alpha$  lies on  $H_0^3(r)$ .  $\square$

**Theorem 3.2** *Let  $\alpha(s)$  be a unit speed spacelike curve in  $E_1^4$ , with spacelike  $N$ ,  $B_1$  and with curvatures  $k_1(s) \neq 0$ ,  $k_2(s) \neq 0$ ,  $k_3(s) \neq 0$  for each  $s \in I \subset R$ . Then  $\alpha$  lies on  $H_0^3(r)$  if and only if*

$$\begin{aligned} (k_3/k_2)(1/k_1)' &= [(1/k_3)(k_2/k_1 + ((1/k_2)(1/k_1)'))']' \\ &[(1/k_3)(k_2/k_1 + (1/k_2)(1/k_1)')]^2 > (1/k_1)^2 + ((1/k_2)(1/k_1)')^2. \end{aligned} \quad (9)$$

**Proof.** Let us first assume that  $\alpha$  lies on  $H_0^3(r)$ . Then by Theorem 3.1. it follows that the relation (1) holds. Differentiating (1) with respect to  $s$ , we find that the equation in (9) is satisfied. Moreover, by using (1) we immediately get that the inequality in (9) holds.

Conversely, let us suppose that (9) holds. It can be easily seen that (9) is the differential of the equation

$$(1/k_1)^2 + ((1/k_2)(1/k_1)')^2 - [(1/k_3)(k_2/k_1 + ((1/k_2)(1/k_1)'))']^2 = \text{constant} = c < 0.$$

Finally, we may take  $c = -r^2$ ,  $r \in R^+$ , so Theorem 3.1 implies that  $\alpha$  lies on  $H_0^3(r)$ .  $\square$

**Theorem 3.3** *A unit speed spacelike curve  $\alpha(s)$  in  $E_1^4$  with spacelike  $N$ ,  $B_1$  and with curvatures  $k_1(s) \neq 0$ ,  $k_2(s) \neq 0$ ,  $k_3(s) \neq 0$  for each  $s \in I \subset R$  lies on  $H_0^3(r)$  if and only if there exist a differentiable function  $f(s)$  such that*

$$fk_3 = k_2/k_1 + ((1/k_2)(1/k_1)')', f' = (k_3/k_2)(1/k_1)', f^2 - (f'/k_3)^2 > (1/k_1)^2. \quad (10)$$

**Proof.** Let us first assume that  $\alpha(s)$  lies on  $H_0^3(r)$ . Then Theorems 3.1. and 3.2. imply that respectively relations (1) and (9) hold. Next, let us define the differentiable function  $f(s)$  by

$$f(s) = (1/k_3)[k_2/k_1 + (1/k_2)(1/k_1)']'. \quad (11)$$

Consequently, by using (1), (9) and (11) we easily find that the relations in (10) are satisfied.



Conversely, let us assume that there exist a differentiable function  $f(s)$  such that the relations in (10) hold for each  $s \in I \subset R$ . By using relations in (10), we easily find that both relations in (9) are satisfied. Finally, by Theorem 3.2. it follows that  $\alpha$  lies on  $H_0^3(r)$ .  $\square$

**Theorem 3.4** *A unit speed spacelike curve  $\alpha(s)$  in  $E_1^4$  with spacelike  $N$ ,  $B_1$  and with curvatures  $k_1(s) \neq 0$ ,  $k_2(s) \neq 0$ ,  $k_3(s) \neq 0$  for each  $s \in I \subset R$  lies on  $H_0^3(r)$  if and only if there exist constants  $A, B \in R$  such that the following relations hold:*

$$\begin{aligned} (1/k_2)(1/k_1)' &= [A + \int_0^s (k_2/k_1) \sinh(\int_0^s k_3 ds) ds] \sinh(\int_0^s k_3 ds) \\ &\quad - [B + \int_0^s (k_2/k_1) \cosh(\int_0^s k_3 ds) ds] \cosh(\int_0^s k_3 ds), \\ [A + \int_0^s (k_2/k_1) \sinh(\int_0^s k_3 ds) ds]^2 &> [B + \int_0^s (k_2/k_1) \cosh(\int_0^s k_3 ds) ds]^2 + (1/k_1)^2. \end{aligned} \tag{12}$$

**Proof.** Let us first suppose that  $\alpha$  lies on  $H_0^3(r)$ . By Theorem 3.3. there exist a differentiable function  $f(s)$  such that (10) hold. Next, let us define the  $C^2$  function  $\theta(s)$  by  $\theta(s) = \int_0^s k_3 ds$ . Moreover, let us define the  $C^1$  functions  $g(s)$  and  $h(s)$  by

$$\begin{aligned} g(s) &= -(1/k_2)(1/k_1)' \sinh \theta + f(s) \cosh \theta - \int_0^s (k_2/k_1) \sinh \theta ds, \\ h(s) &= -(1/k_2)(1/k_1)' \cosh \theta + f(s) \sinh \theta - \int_0^s (k_2/k_1) \cosh \theta ds. \end{aligned} \tag{13}$$

Differentiating functions  $\theta(s)$ ,  $g(s)$  and  $h(s)$  with respect to  $s$ , we find  $\theta'(s) = k_3$ ,  $g'(s) = h'(s) = 0$ . Hence  $g(s) = A$ ,  $h(s) = B$ ,  $A, B \in R$ . Substituting this into (13) yields

$$\begin{aligned} -(1/k_2)(1/k_1)' \sinh \theta + f(s) \cosh \theta - \int_0^s (k_2/k_1) \sinh \theta ds &= A, \\ -(1/k_2)(1/k_1)' \cosh \theta + f(s) \sinh \theta - \int_0^s (k_2/k_1) \cosh \theta ds &= B. \end{aligned} \tag{14}$$

By multiplying the first of the equations in (14) with  $\sinh \theta$  and the second with  $-\cosh \theta$  and adding, we find that the equation in (12) holds. Next, by multiplying the first of the equations in (14) with  $\cosh \theta$  and the second with  $-\sinh \theta$  and adding, we get

$$f(s) = (A + \int_0^s (k_2/k_1) \sinh \theta ds) \cosh \theta - (B + \int_0^s (k_2/k_1) \cosh \theta ds) \sinh \theta. \quad (15)$$

Then the relations (10) and (15) imply that inequality in (12) holds.  $\square$

Conversely, let us suppose that there exist constants  $A, B \in R$  such that (12) holds for each  $s \in I \subset R$ . Differentiating the equation in (12) with respect to  $s$  yields

$$\begin{aligned} ((1/k_2)(1/k_1)')' &= k_3[(A + \int_0^s (k_2/k_1) \sinh(\int_0^s k_3 ds) ds) \cosh(\int_0^s k_3 ds) \\ &\quad - (B + \int_0^s (k_2/k_1) \cosh(\int_0^s k_3 ds) ds) \sinh(\int_0^s k_3 ds)] - k_2/k_1. \end{aligned} \quad (16)$$

Let us define the differentiable function  $f(s)$  by (11). Then by (11) and (16) it follows that (15) holds. Differentiating (15) with respect to  $s$  and using (12) we obtain  $f' = (k_3/k_2)(1/k_1)'$ . Moreover, by using (15), inequality in (12) and by taking the derivative of (15) with respect to  $s$ , we find  $f^2 - (f'/k_3)^2 > (1/k_1)^2$ . Finally, Theorem 3.3. implies that  $\alpha$  lies on  $H_0^3(r)$ .

In the sequel, recall that spacelike curve with spacelike principal normal  $N$  and a null binormal  $B_1$  is called a *partially null curve* [5].

**Remark 3.1.** In the case when  $\alpha$  is spacelike curve with timelike principal normal  $N$  in the space  $E_1^4$ , there holds theorems which are analogous with Theorems 3.1, 3.2, 3.3 and 3.4.

**Theorem 3.5** *A partially null unit speed curve  $\alpha(s)$  in  $E_1^4$  with curvatures  $k_1(s) \neq 0$ ,  $k_2(s) \neq 0$  for each  $s \in I \subset R$  lies fully in a lightlike hyperplane of  $E_1^4$  and has  $k_3(s) = 0$  for each  $s$ .*

**Proof.** By using the Frenet equations, we easily get  $\dot{\alpha} = T$ ,  $\ddot{\alpha} = k_1 N$ ,  $\ddot{\ddot{\alpha}} = -k_1 T + \dot{k}_1 N + k_1 k_2 B_1$ ,  $\ddot{\ddot{\ddot{\alpha}}} = -3k_1 \dot{k}_1 T + (\ddot{k}_1 - k_1^3) N + (2\dot{k}_1 k_1 + k_1 \dot{k}_2 + k_1 k_2 k_3) B_1$ . It follows that  $\dot{\alpha}$ ,  $\ddot{\alpha}$ ,  $\ddot{\ddot{\alpha}}$  are linearly independent vectors and that  $\dot{\alpha}$ ,  $\ddot{\alpha}$ ,  $\ddot{\ddot{\alpha}}$  are linearly dependent vectors. Moreover, by using the MacLaurin expansion for  $\alpha$  given by

$$\alpha(s) = \alpha(0) + \dot{\alpha}(0) \frac{s}{1!} + \ddot{\alpha}(0) \frac{s^2}{2!} + \ddot{\ddot{\alpha}}(0) \frac{s^3}{3!} + \dots,$$

we find that  $\alpha$  lies fully in a lightlike hyperplane  $\pi$  of the space  $E_1^4$  spanned by  $\{\dot{\alpha}(0), \ddot{\alpha}(0), \ddot{\alpha}(0)\}$ . Therefore,  $\alpha$  satisfies the equation of  $\pi$ , namely  $g(\alpha(s) - p, q) = 0$ , where  $p, q \in E_1^4$  and  $q$  is a constant null vector. Differentiating the previous equation three times with respect to  $s$  and by using Frenet equations, we find  $g(B_1, q) = 0$ . Hence  $q = \lambda B_1$ ,  $\lambda \in R$ . Then  $\dot{q} = \lambda k_3 B_1 = 0$  and therefore  $k_3(s) = 0$  for each  $s$ .  $\square$

**Theorem 3.6** *A partially null unit speed curve  $\alpha$  in  $E_1^4$  with curvatures  $k_1(s) \neq 0$ ,  $k_2(s) \neq 0$  for each  $s \in I \subset R$  lies on  $H_0^3(r)$  if and only if  $(1/k_2)(1/k_1)' = \text{constant} \neq 0$ .*

**Proof.** Let us first suppose that  $\alpha$  lies on  $H_0^3(r)$  with center  $m$ . By Theorem 3.5. it follows that  $k_3 = 0$  for each  $s$  and by definition there holds  $g(\alpha - m, \alpha - m) = -r^2$ . Differentiating the previous equation three times with respect to  $s$  and by using Frenet equations, we get  $g(B_1, \alpha - m) = (-1/k_2)(1/k_1)'$ . By taking the derivative of the last equation with respect to  $s$ , we find  $((1/k_2)(1/k_1)')' = 0$  and thus  $(1/k_2)(1/k_1)' = \text{constant} = c_0$ . If  $c_0 = 0$ , then  $g(B_1, \alpha - m) = 0$ . This means that null vector  $B_1$  and timelike vector  $\alpha - m$  are orthogonal in  $E_1^4$ , which is a contradiction. Thus  $c_0 \neq 0$ .

Conversely, if  $(1/k_2)(1/k_1)' = \text{constant} \neq 0$  let us consider the vector

$$m = \alpha + (1/k_1)N - [((1/k_1)^2 + r^2)/((2/k_2)(1/k_1)')]B_1 + (1/k_2)(1/k_1)'B_2, \quad (17)$$

where  $r \in R^+$ . Differentiating the previous equation with respect to  $s$  yields  $m' = 0$ , so  $m = \text{constant}$ . Finally, (17) implies that  $g(\alpha - m, \alpha - m) = -r^2$  and hence  $\alpha$  lies on  $H_0^3(r)$ .  $\square$

**Theorem 3.7** *Let  $\alpha(s)$  be a partially null unit speed curve in  $E_1^4$  with curvatures  $k_1(s) \neq 0$ ,  $k_2(s) \neq 0$  for each  $s \in I \subset R$ . Then  $\alpha$  lies on  $H_0^3(r)$  if and only if*

$$\sinh\left(\int_0^s k_2 ds\right) \int_0^s (1/k_1)' \sinh\left(\int_0^s k_2 ds\right) ds - \cosh\left(\int_0^s k_2 ds\right) \int_0^s (1/k_1)' \cosh\left(\int_0^s k_2 ds\right) ds = 0. \quad (18)$$

**Proof.** Let us first assume that  $\alpha$  lies on  $H_0^3(r)$ . By Theorem 3.6. it follows that  $(1/k_2)(1/k_1)' = \text{constant} \neq 0$ . Let us put  $c_0 = (1/k_2)(1/k_1)'$  and  $\theta(s) = \int_0^s k_2(s)ds$ . Then we have

$$\sinh \theta \int_0^s (1/k_1)' \sinh \theta ds - \cosh \theta \int_0^s (1/k_1)' \cosh \theta ds = 0.$$

Conversely, let us assume that (18) holds. Let us put  $\theta(s) = \int_0^s k_2(s)ds$ . By taking the derivative of (18) with respect to  $s$ , we find

$$\cosh \theta \int_0^s (1/k_1)' \sinh \theta ds - \sinh \theta \int_0^s (1/k_1)' \cosh \theta ds = (1/k_2)(1/k_1)'. \quad (19)$$

Differentiating (19) with respect to  $s$  yields  $((1/k_2)(1/k_1)')' = 0$ . Hence  $(1/k_2)(1/k_1)' = \text{constant} = c_0$ ,  $c_0 \in R$ . If  $c_0 = 0$ , then subtracting (19) of (18) implies a contradiction. Therefore,  $c_0 \neq 0$  so by Theorem 3.6. curve  $\alpha$  lies on  $H_0^3(r)$ .

Recall that a spacelike curve with a null principal normal  $N$  in the space  $E_1^4$  is called a *pseudo null curve* [5]. For such curves, we have the following two theorems.  $\square$

**Theorem 3.8** *Let  $\alpha(s)$  be a pseudo null unit speed curve with curvatures  $k_1(s) = 1$ ,  $k_2(s) \neq 0$ ,  $k_3(s) \neq 0$  for each  $s \in I \subset R$ . Then  $\alpha$  lies on  $H_0^3(r)$  if and only if  $k_3/k_2 = \text{constant} < 0$ .*

**Proof.** Let us first suppose that  $\alpha$  lies on  $H_0^3(r)$  with center  $m$ . By definition we have  $g(\alpha - m, \alpha - m) = -r^2$ . Differentiating the previous equation four times with respect to  $s$  and by using Frenet equations, we get

$$g(\alpha - m, T) = 0, g(N, \alpha - m) = -1, g(B_1, \alpha - m) = 0, g(B_2, \alpha - m) = -k_3/k_2. \quad (20)$$

Moreover, differentiating the last equation in (20) with respect to  $s$ , we find

$$-g(T, \alpha - m) - k_3 g(B_1, \alpha - m) = -(k_3/k_2)'. \quad (21)$$

Therefore, substituting (20) into (21) we obtain  $(k_3/k_2)' = 0$ . Thus  $k_3/k_2 = \text{constant} = c_0$ ,  $c_0 \in R$ . We shall prove that  $c_0 < 0$ . Let us decompose the vector  $\alpha - m$  by

$$\alpha(s) - m = a(s)T(s) + b(s)N(s) + c(s)B_1(s) + d(s)B_2(s), \quad (22)$$

where  $a(s), b(s), c(s), d(s)$  are arbitrary functions. Then by (20) and (22) it follows that  $g(T, \alpha - m) = a = 0$ ,  $g(N, \alpha - m) = d = -1$ ,  $g(B_1, \alpha - m) = c = 0$ ,  $g(B_2, \alpha - m) = b = -c_0$ . Substituting this into (22) we find  $\alpha - m = -c_0N - B_2$  and thus  $g(\alpha - m, \alpha - m) = 2c_0 = -r^2$ . Hence  $c_0 < 0$ .

Conversely, let us suppose that  $k_3/k_2 = \text{constant} < 0$ . Then we may put  $k_3/k_2 = -r^2/2, r \in R^+$ . Let us consider vector  $m = \alpha - (r^2/2)N + B_2$ . Differentiating the previous equation with respect to  $s$ , we find  $m' = 0$  and thus  $m = \text{constant}$ . Moreover,  $g(\alpha - m, \alpha - m) = -r^2$ , so  $\alpha$  lies on  $H_0^3(r)$ .  $\square$

**Theorem 3.9** *Let  $\alpha(s)$  be a pseudo null unit speed curve with curvatures  $k_1(s) = 1$ ,  $k_2(s) \neq 0$ ,  $k_3(s) \neq 0$  for each  $s \in I \subset R$ . Then  $\alpha$  lies on  $H_0^3(r)$  if and only if the following two relations hold:*

$$\sinh\left(\int_0^s k_3 ds\right) \int_0^s k_2 \sinh\left(\int_0^s k_3 ds\right) ds - \cosh\left(\int_0^s k_3 ds\right) \int_0^s k_2 \cosh\left(\int_0^s k_3 ds\right) ds = 0, \quad (23)$$

$$\cosh\left(\int_0^s k_3 ds\right) \int_0^s k_2 \sinh\left(\int_0^s k_3 ds\right) ds < \sinh\left(\int_0^s k_3 ds\right) \int_0^s k_2 \cosh\left(\int_0^s k_3 ds\right) ds.$$

**Proof.** Let us first assume that  $\alpha$  lies on  $H_0^3$ . By Theorem 3.8 we have  $k_2/k_3 = \text{constant} < 0$ . Let us put  $k_2/k_3 = -c_0^2$ ,  $c_0 \in R$  and  $\theta(s) = \int_0^s k_3(s) ds$ . Then we easily find

$$\sinh \theta \int_0^s k_2 \sinh \theta ds - \cosh \theta \int_0^s k_2 ds = 0.$$

Differentiating the previous equation with respect to  $s$ , we get

$$\cosh \theta \int_0^s k_2 \sinh \theta ds - \sinh \theta \int_0^s k_2 \cosh \theta ds = -c_0^2 = k_2/k_3, \quad (24)$$

and hence

$$\cosh \theta \int_0^s k_2 \sinh \theta ds < \sinh \theta \int_0^s k_2 \cosh \theta ds.$$

Conversely, if (23) holds, let us put  $\theta(s) = \int_0^s k_3(s) ds$ . By taking the derivative of (23) with respect to  $s$ , we find

$$\cosh \theta \int_0^s k_2 \sinh \theta ds - \sinh \theta \int_0^s k_2 \cosh \theta ds = k_2/k_3. \quad (25)$$

Differentiating (25) with respect to  $s$  yields  $(k_2/k_3)' = 0$ . Hence  $k_2/k_3 = \text{constant} = c_0$ ,  $c_0 \in \mathbb{R}$ . Moreover, (23) and (25) imply  $c_0 < 0$ . Finally, by Theorem 3.8 it follows that  $\alpha$  lies on  $H_0^3(r)$ .  $\square$

**Theorem 3.10** *There are no timelike and no null curves  $\alpha(s)$  lying on the pseudohyperbolic space  $H_0^3(r)$  in  $E_1^4$ .*

**Proof.** If  $\alpha(s)$  is timelike unit speed curve lying on  $H_0^3(r)$  with center  $m$ , then  $g(\alpha - m, \alpha - m) = -r^2$ . Differentiating the previous equation with respect to  $s$ , we get  $g(T, \alpha - m) = 0$ . It follows that  $T$  and  $\alpha - m$  are two timelike mutually orthogonal vectors in  $E_1^4$ , which is a contradiction. If  $\alpha(s)$  is a null curve lying on  $H_0^3(r)$ , then in a similar way it follows that null vector  $T$  and timelike vector  $\alpha - m$  are orthogonal vectors in  $E_1^4$ , which is a contradiction.  $\square$

## References

- [1] Bonnor, W. B.: Null curves in a Minkowski space–time, *Tensor*, 20 1969, 229–242.
- [2] Milman, R. S. and Parker G. D.: *Elements of Differential Geometry*, Prentice–Hall Inc., Englewood Cliffs, New Jersey, 1977.
- [3] O’Neill, B.: *Semi–Riemannian Geometry*, Academic Press, New York, 1983.

- [4] Petrović-Torgašev, M. and Šučurović, E.: Some characterizations of curves lying on the pseudohyperbolic space  $H_0^2$  in the Minkowski space  $E_1^3$ , Kragujevac J. Math. 22 2000, 71–82.
- [5] Walrave, J.: Curves and surfaces in Minkowski space, Doctoral thesis, K. U. Leuven, Fac. of Science, Leuven, 1995.
- [6] Wong, Y. C.: A global formulation of the condition for a curve to lie in a sphere, Monatshefte für Mathematik, 67, 1963, 363–365.
- [7] Wong, Y. C.: On an explicit characterization of spherical curves, Proceedings of the American Mathematical Society 34, 1972, 239–242.

Çetin CAMCI

Received 08.08.2002

Department of Mathematics,  
Faculty of Science,  
Ankara University,  
06100 Ankara-TURKEY

Kazım İLARSLAN

Department of Mathematics,  
Faculty of Science and Arts,  
Kocatepe University,  
Afyon-TURKEY

e-mail: kilarslan@aku.edu.tr

Emilija ŠUĆUROVIĆ

Department of Mathematics,  
Faculty of Science,  
Kragujevac University,  
Radoja Domanovića 12, 34000  
Yugoslavia

e-mail: emilija@knez.uis.kg.ac.yu