

1-1-2015

## Pandemic H1N1 IgG levels in healthcare workers

VIROJ WIWANITKIT

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

---

### Recommended Citation

WIWANITKIT, VIROJ (2015) "Pandemic H1N1 IgG levels in healthcare workers," *Turkish Journal of Medical Sciences*: Vol. 45: No. 2, Article 33. <https://doi.org/10.3906/sag-1501-22>

Available at: <https://journals.tubitak.gov.tr/medical/vol45/iss2/33>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Pandemic H1N1 IgG levels in healthcare workers

Viroj WIWANITKIT

Surin Rajabhat University, Surin Thailand

E-mail: wviroj@yahoo.com

Received: 07.01.2015 • Accepted: 07.02.2015 • Published Online: 01.04.2015 • Printed: 30.04.2015

Editor, the recent report on pandemic H1N1 IgG levels in healthcare workers is interesting (1). Çıfci et al. found that “the IgG antibody level was higher in the vaccinated healthcare workers than among the unvaccinated healthcare workers (1).” The implication of this important finding should be discussed. Whether this finding shows the effectiveness and usefulness of vaccination among healthcare workers or not should be further considered. First, IgG antibody might be due to natural infection or vaccination. Since there are no data for the pre-vaccination stage of vaccinated workers, this

point is still a myth. A recent report by Khaokam et al. showed that “approximately half of H1N1 infections were asymptomatic (2).” In addition, the sensitivity of the IgG detection should be discussed. In a recent report by Li et al., it was shown that “the IgG antibody response to A(H1N1)pdm09 HA was cross-reactive with HAs from H1, H3, H5, and H13 subtypes (3).” Finally, the reason for no vaccination should be further discussed. Indeed, different kinds of healthcare workers might have different chances to be exposed to influenza. This might be a possible reason why some workers might not have vaccination.

### References

1. Çıfci A, Erol Ö, Cesur S, Aksoy N, Kisa Ü. The comparison of pandemic H1N1 IgG levels between H1N1 influenza-vaccinated healthcare workers and unvaccinated healthcare workers. *Turk J Med Sci* 2014; 44: 1047–1050.
2. Khaokam CB, Selent M, Loustalot FV, Zarecki SM, Harrington D, Hoke E, Faix DJ, Ortiguerra R, Alvarez B, Almond N et al. Seroepidemiologic investigation of an outbreak of pandemic influenza A H1N1 2009 aboard a US Navy vessel—San Diego, 2009. *Influenza Other Respir Viruses* 2013; 7: 791–798.
3. Li ZN, Lin SC, Carney PJ, Li J, Liu F, Lu X, Liu M, Stevens J, Levine M, Katz JM, Hancock K. IgM, IgG, and IgA antibody responses to influenza A(H1N1)pdm09 hemagglutinin in infected persons during the first wave of the 2009 pandemic in the United States. *Clin Vaccine Immunol* 2014; 21: 1054–1060.

### Reply to Letter to the Editor:

I would like to thank for your important contributions to our study. We will consider your suggestions in our succeeding studies.

### Aydın ÇİFCİ

Department of Internal Disease, Faculty of Medicine,

Kırıkkale University,

Kırıkkale, Turkey

E-mail: dr.aydin.71@hotmail.com