

1-1-1999

On The Artin Conductor $f_{\text{Artin}}(\chi)$ of a Character χ of Gal (E/K) I: Basic Definitions

KAZIM İLHAN İKEDA

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

İKEDA, KAZIM İLHAN (1999) "On The Artin Conductor $f_{\text{Artin}}(\chi)$ of a Character χ of Gal (E/K) I: Basic Definitions," *Turkish Journal of Mathematics*: Vol. 23: No. 4, Article 4. Available at: <https://journals.tubitak.gov.tr/math/vol23/iss4/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

ON THE ARTIN CONDUCTOR $f_{\text{Artin}}(\chi)$ OF A CHARACTER χ of $\text{Gal}(E/K)$ I: BASIC DEFINITIONS

*Kâzım İlhan İkedâ**

Dedicated to Professor Masatoshi Ikeda on the occasion of his 70th birthday

Abstract

Let K be a local field with finite residue class field and E a finite Galois extension over K . In this paper, we study the Artin conductor $f_{\text{Artin}}(\chi_\rho)$ of a character χ_ρ associated to a representation $\rho : \text{Gal}(E/K) \rightarrow GL(V)$ of $\text{Gal}(E/K)$ with metabelian kernel $\ker(\rho)$. In order to do so, we first review the Artin character $a_{\text{Gal}(E/K)}$ of $\text{Gal}(E/K)$ and review the metabelian local class field theory. We finally propose the definition of the conductor $f(E/K)$ of a metabelian extension E/K in the sense of Koch-de Shalit local class field theory, and compute $f_{\text{Artin}}(\chi_\rho)$ under a suitable assumption.

§1. Introduction

Let K be a local field with finite residue class field $O_K/\mathfrak{p}_K =: \kappa_K$ of q_K elements, where as usual, O_K stands for the ring of integers in K with the unique maximal ideal \mathfrak{p}_K . Let ν denote the corresponding normalized valuation on K (normalized by $\nu(K^\times) = \mathbb{Z}^\times$), and $\tilde{\nu}$ the unique extension of ν to a fixed separable closure K^{sep} of K .

Let E be a finite Galois extension over K , and let $\rho : \text{Gal}(E/K) \rightarrow GL(V)$ be an irreducible (finite-dimensional) representation of the Galois group $\text{Gal}(E/K)$ of E/K over \mathbb{C} . Furthermore, setting $E^{\ker(\rho)}$ to denote the fixed field of $\ker(\rho)$, suppose that $E^{\ker(\rho)}/K$

Key words and phrases. Local fields, higher-ramification groups, Artin conductor, metabelian local class field theory

*This work has been carried out under the frame of the Project “ $GL(n)$ ” at Feza Gürsey Institute of Theoretical Physics, İstanbul-Turkey.

is an abelian extension. It is then well-known that

$$(1.1) \quad \mathfrak{f}(E^{\ker(\rho)}/K) = \mathfrak{f}_{\text{Artin}}(\chi_\rho),$$

where $\mathfrak{f}(E^{\ker(\rho)}/K)$ is the conductor of the abelian extension $E^{\ker(\rho)}/K$, which is defined in the sense of abelian local class field theory (cf. chapter 5, paragraph 1.6 of [5]), and $\mathfrak{f}_{\text{Artin}}(\chi_\rho)$ is the Artin conductor of the character $\chi_\rho : \text{Gal}(E/K) \rightarrow \mathbb{C}$ associated to the representation (ρ, V) of $\text{Gal}(E/K)$ (which will be reviewed in the next section).

It is then natural to ask more generally the behaviour of $\mathfrak{f}_{\text{Artin}}(\chi_\rho)$ without any assumption on the kernel $\ker(\rho)$ of the representation (ρ, V) . It seems that, this general problem is closely related with the 2-abelian local class field theory of Koch and de Shalit, and more generally with the n -abelian local class field theory.

In this paper, we start our investigation on this problem for the metabelian kernel $\ker(\rho)$ case. In particular, in this paper, we will collect the necessary basic definitions, propose the definition of the conductor $\mathfrak{f}(E/K)$ of a finite metabelian (=2-abelian) extension E/K in the sense of metabelian local class field theory (following the main theorem of higher ramification theory in metabelian local class field theory), and under a suitable assumption, compute $\mathfrak{f}_{\text{Artin}}(\chi_\rho)$.

Our aim, which is the subject of our next paper is to study the following problem (which should be viewed as the metabelian generalization of eq. no. (1.1)):

Problem 1.1 *Let E/K be a finite Galois extension, $\rho : \text{Gal}(E/K) \rightarrow \text{GL}(V)$ an irreducible (finite-dimensional) representation of the Galois group of E/K over \mathbb{C} . Suppose that $E^{\ker(\rho)}/K$ is a 2-abelian extension. Then,*

$$\mathfrak{f}(E^{\ker(\rho)}/K) = \mathfrak{f}_{\text{Artin}}(\chi_\rho).$$

§2. Artin representation $A_{\text{Gal}(E/K)}$ of $\text{Gal}(E/K)$

For a finite separable extension L/K , and for any $\sigma \in \text{Hom}_K(L, K^{\text{sep}})$, introduce

$$i_{L/K}(\sigma) := \min_{x \in O_L} \{\tilde{\nu}(\sigma(x) - x)\},$$

put

$$\gamma_t := \#\{\sigma \in \text{Hom}_K(L, K^{\text{sep}}) : i_{L/K}(\sigma) \geq t\}$$

for $-1 \leq t \in \mathbb{R}$, and define the function $\varphi_{L/K} : \mathbb{R}_{\geq -1} \rightarrow \mathbb{R}_{\geq -1}$ (the Hasse-Herbrand

transition function of the extension L/K) by

$$\varphi_{L/K}(u) := \begin{cases} t_0^u \frac{\gamma_t}{\gamma_0} dt, & 0 \leq u \in \mathbb{R}, \\ u, & -1 \leq u \leq 0. \end{cases}$$

It is well-known that, $\varphi_{L/K} : \mathbb{R}_{\geq -1} \rightarrow \mathbb{R}_{\geq -1}$ is a monotone-increasing (piecewise linear) function, and induces a homeomorphism $\mathbb{R}_{\geq -1} \rightarrow \mathbb{R}_{\geq -1}$. Let $\psi_{L/K} : \mathbb{R}_{\geq -1} \rightarrow \mathbb{R}_{\geq -1}$ be the mapping inverse to the function $\varphi_{L/K} : \mathbb{R}_{\geq -1} \rightarrow \mathbb{R}_{\geq -1}$.

Now, assume that E is a finite Galois extension over K with Galois group $\text{Gal}(E/K) =: G$. The subgroup G_u of G defined by

$$G_u = \{\sigma \in G : i_{E/K}(\sigma) \geq u + 1\}$$

for $-1 \leq u \in \mathbb{R}$ is called the u^{th} ramification group of G (in the lower numbering), and $\{G_u\}_{u \in \mathbb{R}_{\geq -1}}$ induces a filtration on G , called the lower ramification filtration of G . A break in the filtration $\{G_u\}_{u \in \mathbb{R}_{\geq -1}}$ of G is defined to be any number $u \in \mathbb{R}_{\geq -1}$ satisfying $G_u \neq G_{u+\varepsilon}$ for every $0 < \varepsilon \in \mathbb{R}$. The function $\psi_{L/K} : \mathbb{R}_{\geq -1} \rightarrow \mathbb{R}_{\geq -1}$ induces the upper ramification filtration $\{G^v\}_{v \in \mathbb{R}_{\geq -1}}$ on G by setting

$$G^v := G_{\psi_{L/K}(v)}$$

for $-1 \leq v \in \mathbb{R}$, where G^v is called the v^{th} upper ramification group of G . A break in the upper filtration $\{G^v\}_{v \in \mathbb{R}_{\geq -1}}$ of G is defined to be any number $v \in \mathbb{R}_{\geq -1}$ satisfying $G^v \neq G^{v+\varepsilon}$ for every $0 < \varepsilon \in \mathbb{R}$.

The basic properties of lower and upper ramification filtrations on G are as follows: Suppose that $K \subseteq F \subseteq E$ is a sub-extension of E/K , let $\text{Gal}(E/F) = H$.

(i) The lower numbering on G passess well to the subgroup H of G in the sense that:

$$H_u = H \cap G_u$$

for $-1 \leq u \in \mathbb{R}$;

(ii) and if furthermore, $H \triangleleft G$, the upper numbering on G passess well to the quotient G/H as:

$$(G/H)^v = G^v H/H$$

for $-1 \leq v \in \mathbb{R}$;

(iii) (transitivity of the Hasse-Herbrand function) $\varphi_{E/K} = \varphi_{F/K} \circ \varphi_{E/F}$.

Let $\mathcal{B}_{\mathcal{E}/\mathcal{K}}$ be the set of all numbers $v \in \mathbb{R}_{\geq -1}$ which occur as breaks in the upper ramification filtration of G . Then,

(iv) (Hasse-Arf theorem) If E/K is an abelian extension, then $\mathcal{B}_{\mathcal{E}/\mathcal{K}}$ is a finite subset of $\mathbb{Z} \cap \mathbb{R}_{\geq -1}$.

Remark 2.1 If E/K is an infinite Galois extension with Galois group $\text{Gal}(E/K) = G$ (which is a topological group under the Krull topology), define the upper ramification filtration $\{G^v\}_{v \in \mathbb{R}_{\geq -1}}$ on G by the projective limit

$$G^v := \varprojlim_{K \subseteq F \subseteq E} \text{Gal}(F/K)^v$$

defined over the transition morphisms

$$\begin{array}{ccc} (G/\text{Gal}(E/F))^v & \xleftarrow{t_F^{F'}} & (G/\text{Gal}(E/F'))^v \\ \parallel & & \parallel \\ G^v \text{Gal}(E/F)/\text{Gal}(E/F) & \xleftarrow{\text{can.}} & G^v \text{Gal}(E/F')/\text{Gal}(E/F') \end{array}$$

induced from (ii), as $K \subseteq F \subseteq F' \subseteq E$ runs over all finite Galois extensions F and F' over K inside E . Observe that:

- (v) $G^{-1} = G$ and G^0 is the inertia group of G ;
- (vi) $\bigcap_{v \in \mathbb{R}_{\geq -1}} G^v = \langle 1 \rangle$;
- (vii) G^v is a closed subgroup of G (with respect to the Krull topology) for $-1 \leq v \in \mathbb{R}$.

In this setting, a number $-1 \leq v \in \mathbb{R}$ is said to be a break in the upper ramification filtration $\{G^v\}_{v \in \mathbb{R}_{\geq -1}}$ of G , if v is a break in the upper filtration of some finite quotient G/H for some $H \triangleleft G$. As introduced previously, let $\mathcal{B}_{\mathcal{E}/\mathcal{K}}$ be the set of all numbers $v \in \mathbb{R}_{\geq -1}$, which occur as breaks in the upper ramification filtration of G . Then,

(viii) (Hasse-Arf theorem.) $\mathcal{B}_{\mathcal{K}^{-1}/\mathcal{K}} = \mathbb{Z} \cap \mathbb{R}_{\geq -\infty}$;

and the final important result, in the spirit of Koch-de Shalit local class field theory, is

(ix) $\mathcal{B}_{\mathcal{K}^{\vee}/\mathcal{K}} = \mathbb{Q} \cap \mathbb{R}_{\geq -\infty}$. \square

Again assume that E/K is a finite Galois extension. Introduce the class function $a_{\text{Gal}(E/K)} : \text{Gal}(E/K) \rightarrow \mathbb{C}$ on the Galois group $\text{Gal}(E/K)$ of the extension E/K by

$$a_{\text{Gal}(E/K)}(\sigma) = \begin{cases} -f(E/K)i_{E/K}(\sigma), & \sigma \neq 1; \\ f(E/K)\sum_{\omega \neq 1} i_{E/K}(\omega), & \sigma = 1. \end{cases}$$

As the set of irreducible representations χ on $\text{Gal}(E/K)$ forms an orthonormal basis of the vector space $C(\text{Gal}(E/K))$ of all class functions on $\text{Gal}(E/K)$ over \mathbb{C} equipped with the hermitian scalar-product

$$(\ , \) : C(\text{Gal}(E/K)) \times C(\text{Gal}(E/K)) \rightarrow \mathbb{C}$$

defined by

$$(f, g) = \frac{1}{[E : K]} \sum_{\sigma} f(\sigma)\overline{g(\sigma)}$$

on the \mathbb{C} -vector space $C(\text{Gal}(E/K))$, the function $a_{\text{Gal}(E/K)} : \text{Gal}(E/K) \rightarrow \mathbb{C}$ is a \mathbb{C} -linear combination

$$a_{\text{Gal}(E/K)} = \sum_{\chi} f(\chi)\chi$$

of irreducible characters χ of $\text{Gal}(E/K)$, where the uniquely defined χ -coordinate of $a_{\text{Gal}(E/K)}$ is $f(\chi) = (a_{\text{Gal}(E/K)}, \chi) \in \mathbb{C}$.

Now, applying Hasse-Arf theorem for the characters χ of $\text{Gal}(E/K)$ of degree 1, and then using Brauer induction theory for an arbitrary character χ of $\text{Gal}(E/K)$, it follows that $f(\chi) \in \mathbb{Z}_{\geq 0}$. Hence the class function $a_{\text{Gal}(E/K)} : \text{Gal}(E/K) \rightarrow \mathbb{C}$ is in fact a character (called the Artin character of $\text{Gal}(E/K)$) of a representation $A_{\text{Gal}(E/K)} : \text{Gal}(E/K) \rightarrow GL(V)$, the Artin representation of $\text{Gal}(E/K)$. The explicit construction of this representation is not known.

Let $\phi \in C(\text{Gal}(E/K))$ and put $f_{\text{Artin}}(\phi) = (\phi, a_{\text{Gal}(E/K)}) \in \mathbb{C}$. If χ is a character of $\text{Gal}(E/K)$, define the Artin conductor $\mathfrak{f}_{\text{Artin}}(\chi)$ of χ to be the ideal in O_K defined by

$$\mathfrak{f}_{\text{Artin}}(\chi) = \mathfrak{p}_K^{f_{\text{Artin}}(\chi)}.$$

§3. Metabelian local class field theory

Recall that, a group G is called n -abelian, if the n^{th} -commutator subgroup $G^{(n)}$ of G is $\langle 1_G \rangle$. Now, an extension E over any field K is called n -abelian, if it is a Galois extension with an n -abelian Galois group $\text{Gal}(E/K)$. Fix a seperable closure K^{sep} of K (when it is convenient, instead of K^{sep} , sometimes the notation \overline{K} will be used to denote the seperable closure of K), and let $K^{(ab)^n}$ denote the maximal n -abelian extension of K inside K^{sep} (cf. part 2 of [1]).

In [4], Koch and de Shalit have constructed class field theory for metabelian (that is, 2-abelian) extensions of a local field K , which induces the abelian local class field theory when restricted to the abelian extensions of K . The main idea in [4] is the following: let K be a local field with finite residue field κ_K of q_K elements. Let $\overline{\kappa}_K$ be a fixed seperable closure of κ_K . Fix an extension $\phi \in \text{Gal}(K^{\text{sep}}/K)$ of the Frobenius automorphism ϕ_K over K , that is fix a Lubin-Tate splitting ϕ over K . It is then well-known that, depending on the choice of ϕ , there exists a unique norm-compatible set of primes

$$\mathcal{L}_\phi^o = \{\pi_L \in L : K \subseteq L \subset K_\phi \text{ s.t } [L : K] < \infty\},$$

(where K_ϕ denotes the fixed field of ϕ in K^{sep}), which has a canonical extension to a norm-compatible set of primes

$$\mathcal{L}_\phi = \{\pi_L \in \widetilde{L}^{nr} : K \subseteq L \subset K^{\text{sep}} \text{ s.t } [L : K] < \infty\},$$

called the Lubin-Tate labelling over K attached to the Lubin-Tate splitting ϕ over K . For a finite extension L over K which is pointwise fixed by ϕ , there exists a unique Lubin-Tate formal power series $f_L \in O_L[[X]]$ (belonging to π_L for \widetilde{L}^{nr} , where π_L chosen from \mathcal{L}_ϕ^o) satisfying certain properties (cf. 0.3 in [4]), and a unique Lubin-Tate formal group law $F_L \in O_{\widetilde{L}^{nr}}[[X, Y]]$ attached to f_L . Let $[u]_{f_L} : F_L \rightarrow F_L$ be the unique endomorphism of F_L over $O_{\widetilde{L}^{nr}}$ of the form $[u]_{f_L} = uX + (\text{higher-degree terms}) \in XO_{\widetilde{L}^{nr}}[[X]]$ for $u \in O_L$, and let

$$\{u\}_{f_L} = [u]_{f_L} \pmod{\pi_L}.$$

Let $\widehat{K^\times}$ denote the profinite completion of K^\times , and fix the isomorphism $\widehat{K^\times} \xrightarrow{\sim} U(K) \times \widehat{\mathbb{Z}}$ defined by $a = u_a \pi_K^{\nu_a} \mapsto (u_a, \nu_a)$ for $a \in \widehat{K^\times}$, where π_K is the prime element in K chosen uniquely from \mathcal{L}_ϕ^o . For $1 \leq d \in \mathbb{Z}$, consider the topological group

$$\mathfrak{G}_d^{[2]}(K, \phi) := \left\{ (a, \xi) \in \widehat{K^\times} \times \overline{\kappa}_K[[X]]^\times : \frac{\xi^{\phi^d}}{\xi} = \frac{\{u_a\}_{f_K}}{X} \right\}$$

under the law of composition defined by

$$(a, \xi)(b, \psi) = \left(ab, \xi(\psi^{\phi^{-\nu_a}} \circ \{u_a\}_{f_K}) \right)$$

for $(a, \xi), (b, \psi) \in \mathfrak{G}_d^{[2]}(K, \phi)$, and with a basis of neighborhoods of the underlying topology

$$\mathfrak{G}_d^{[2]}(K, \phi)^{(i,j)} = \left\{ (a, \xi) \in \mathfrak{G}_d^{[2]}(K, \phi) : a \in U^i(K), \quad \xi \equiv 1 \pmod{X^j} \right\}$$

for $0 \leq i, j \in \mathbb{Z}$. Let $\mathfrak{G}^{[2]}(K, \phi) := \varprojlim_d \mathfrak{G}_d^{[2]}(K, \phi)$, where the projective limit is defined over the transition morphisms $\mathfrak{G}_{d'}^{[2]}(K, \phi) \rightarrow \mathfrak{G}_d^{[2]}(K, \phi)$ for every $1 \leq d, d' \in \mathbb{Z}$ with $d \mid d'$, which is defined by

$$(a, \xi) \mapsto \left(a, \prod_{0 \leq i \leq \frac{d'}{d}} \xi^{\phi^{di}} \right)$$

for $(a, \xi) \in \mathfrak{G}_{d'}^{[2]}(K, \phi)$.

Let L be a ϕ -compatible extension over K , $\phi' = \phi^{f(L/K)}$ (cf. 0.4 in [4]), and

$$\mathfrak{G}^{[2]}(L, \phi') \xrightarrow{M_{\phi, L/K}} \mathfrak{G}^{[2]}(K, \phi)$$


the ‘‘2-abelian’’ norm map which is a canonical morphism defined via Coleman theory (cf. 1.5 in [4]). As a notation, put $M_{\phi}(L/K) := M_{\phi, L/K}(\mathfrak{G}^{[2]}(L, \phi'))$, and for an infinite extension E/K which is a union of ϕ -compatible extensions L over K (such an E will be called as an infinite ϕ -compatible extension over K), define $M_{\phi}(E/K) := \bigcap_{K \subseteq L \subseteq E} M_{\phi}(L/K)$ where L runs over all ϕ -compatible sub-extensions in E/K . The map

$$\left\{ \begin{array}{l} \phi\text{-compatible} \\ \text{extensions over } K \end{array} \right\} \xrightarrow{M_{\phi}} \left\{ \begin{array}{l} \text{subgroups of} \\ \mathfrak{G}^{[2]}(K, \phi) \end{array} \right\}$$

defined by

$$L/K \mapsto M_{\phi}(L/K)$$

for every ϕ -compatible extension L/K , has a canonical extension



The metabelian local class field theory of Koch-de Shalit states the following.

Theorem 3.1 (Metabelian local class field theory) *Let K be a local field with finite residue field κ_K of q_K elements. Fix an extension $\phi \in \text{Gal}(K^{sep}/K)$ of the Frobenius automorphism $\phi_K \in \text{Gal}(K^{nr}/K)$.*

(i) *There exists an order-preserving bijection*

$$\left\{ \begin{array}{c} \text{2-abelian extensions} \\ \text{over } K \end{array} \right\} \xrightarrow{M_\phi} \left\{ \begin{array}{c} \text{closed subgroups} \\ \text{of } \mathfrak{G}^{[2]}(K, \phi) \end{array} \right\}$$

defined by

$$L/K \mapsto M_\phi(L/K)$$

for any 2-abelian extension L/K satisfying

$$[L : K] = \left(\mathfrak{G}^{[2]}(K, \phi) : M_\phi(L/K) \right);$$

(ii) *there exists a topological isomorphism*

$$\mathfrak{G}^{[2]}(K, \phi) \xrightarrow[\sim]{(? , K)_\phi} \text{Gal}(K^{(ab)^2}/K)$$

called the “2-abelian” local Artin map, which depends only on the choice of ϕ ;

(iii) for any 2-abelian extension L over K , the surjective morphism

$$\begin{array}{ccc} \mathfrak{G}^{[2]}(K, \phi) & \xrightarrow[\sim]{(? , K)_\phi} & \text{Gal}(K^{(ab)^2}/K) \xrightarrow{\text{res}_i^{K^{(ab)^2}}} \text{Gal}(L/K) \\ & \searrow & \nearrow \\ & & (? , K)_{\phi L} \end{array}$$

induces the canonical topological isomorphism

$$\mathfrak{G}^{[2]}(K, \phi)/M_\phi(L/K) \xrightarrow[\sim]{(? , L/K)_\phi} \text{Gal}(L/K),$$

where $M_\phi(L/K)$ is the closed normal subgroup $M_{\phi, L/K}(\mathfrak{G}^{[2]}(L, \phi'))$ of $\mathfrak{G}^{[2]}(K, \phi)$;

(iv) if K' is ϕ -compatible over K , then the square

$$\begin{array}{ccc} \mathfrak{G}^{[2]}(K', \phi') & \xrightarrow{(? , K')_{\phi'}} & \text{Gal}(K'^{(ab)^2}/K') \\ M_{\phi, K'/K} \downarrow & & \downarrow \text{res}_K \\ \mathfrak{G}^{[2]}(K, \phi) & \xrightarrow{(? , K)_\phi} & \text{Gal}(K^{(ab)^2}/K) \end{array}$$

is commutative;

(v) the breaks in the upper-numbering of the ramification groups for 2-abelian extensions over K occur at $r = 0$ and at rational numbers of the form

$$u_{i,j} = i - \frac{q_K^i - 1 - j}{q_K^i - q_K^{i-1}}$$

with $1 \leq i \in \mathbb{Z}$ and $j \in \mathbb{Z}$ such that $q_K^{i-1} \leq j < q_K^i$. If E/K is a finite 2-abelian extension, and

$$i - 1 \leq u_{i,j-1} < r \leq u_{i,j} \leq i,$$

then

$$\alpha \in M_\phi(E/K)\mathfrak{G}^{[2]}(K, \phi)^{(i,j)} \iff (\alpha, E/K)_\phi \in \text{Gal}(E/K)^{(r)}.$$

The 2-abelian local class field theory is related with the abelian local class field theory as follows: if E/K is a 2-abelian extension, then

$$\text{pr}_1(M_\phi(E/K)) = N(E/K).$$

The abelian extension over K which is class field to $N(E/K)$ is the maximal abelian sub-extension of E/K .

§4. Conductor of a metabelian extension E/K

In this section, let E be a finite 2-abelian extension over K .

Observe that, if $0 \leq r \in \mathbb{R}$, then there exists a unique $1 \leq i(r) = i \in \mathbb{Z}$ and a unique $j(r) = j \in \{q_K^{i-1}, \dots, q_K^{i-1} + (q_K^i - q_K^{i-1} - 1)\}$ such that

$$i(r) - 1 \leq u_{i(r),j(r)-1} < r \leq u_{i(r),j(r)} \leq i(r).$$

In fact, $i(r) - 1 = \llbracket r \rrbracket$, where $0 \leq \llbracket r \rrbracket \in \mathbb{Z}$ is the “integer part” of the number r .

Define the conductor $\mathfrak{f}(E/K)$ of the extension E/K (in the sense of metabelian local class field theory) to be the ideal

$$\mathfrak{f}(E/K) = \mathfrak{p}_K^i$$

in O_K , where $1 \leq i \in \mathbb{Z}$ is the smallest integer defined by the condition

$$\mathfrak{G}^{[2]}(K, \phi)^{(i, q_K^i - 1)} \subseteq M_\phi(E/K).$$

Observe that, by the metabelian local class field theory, the conductor $\mathfrak{f}(E/K)$ of the 2-abelian extension E/K is defined by $\mathfrak{f}(E/K) = \mathfrak{p}_K^i$, where $1 \leq i \in \mathbb{Z}$ is the smallest integer such that $\text{Gal}(E/K)^{(r)} = 1$ for every $i - 1 \leq i - \frac{1}{q_K^i - q_K^{i-1}} < r \leq i$.

§5. Computation of $f_{\text{Artin}}(\chi_\rho)$

Let $\rho : \text{Gal}(E/K) \rightarrow GL(V)$ be an irreducible representation of the Galois group $\text{Gal}(E/K)$ of a finite Galois extension E/K . Let $\ker(\rho)$ be the kernel of the representation (ρ, V) of $\text{Gal}(E/K)$. Let $E^{\ker(\rho)}$ be the fixed field of $\ker(\rho)$. Note that, the representation $\tilde{\rho} : \text{Gal}(E^{\ker(\rho)}/K) \xrightarrow{\sim} \text{Gal}(E/K)/\ker(\rho) \rightarrow GL(V)$ is an irreducible representation of $\text{Gal}(E^{\ker(\rho)}/K)$.

Now, if $\phi : \text{Gal}(E^{\ker(\rho)}/K) \rightarrow \mathbb{C}$ is any central function on $\text{Gal}(E^{\ker(\rho)}/K)$, and ϕ' is the corresponding central function on $\text{Gal}(E/K)$ defined by

$$\phi' : \text{Gal}(E/K) \xrightarrow{\text{canonical map}} \text{Gal}(E^{\ker(\rho)}/K) \xrightarrow{\phi} \mathbb{C},$$

then $f_{\text{Artin}}(\phi) = f_{\text{Artin}}(\phi')$. In particular, for the case $\phi' = \chi_\rho$ and $\phi = \chi_{\tilde{\rho}}$, we get $f_{\text{Artin}}(\chi_\rho) = f_{\text{Artin}}(\chi_{\tilde{\rho}})$.

We will now explicitly compute $(\chi_{\tilde{\rho}}, a_{\text{Gal}(E^{\ker(\rho)}/K)})$ when $E^{\ker(\rho)}/K$ is a metabelian extension. For simplicity, fix the following notation: $G = \text{Gal}(E^{\ker(\rho)}/K)$ and $H = \text{Gal}(E^{\ker(\rho)}/E_1)$, where $E_1 = E \cap K^{ab}$ is the maximal abelian sub-extension in E/K . We assume that H is not contained in the center $Z(G)$ of G . Then it is well-known that there exists a subgroup $H \subseteq H_o \subsetneq G$ and an irreducible representation ψ of H_o such that

$$\rho = \text{Ind}_{H_o}^G(\psi).$$

We further assume that $H = H_o$. Then, ψ is a 1-dimensional representation of H , since H is abelian and by the Frobenius reciprocity:

$$f_{\text{Artin}}(\chi_\rho) = (\text{Ind}_H^G(\psi), a_G) = (\psi, \text{Res}_H(a_G)).$$

But $\text{Res}_H(a_G) = \text{ord}_{\mathfrak{p}_K}(\mathfrak{d}_{E_1/K})r_H + [\kappa_{E_1} : \kappa_K]a_H$, where $\mathfrak{d}_{E_1/K}$ is the discriminant of E_1/K and r_H is the character of the regular representation of H . Hence,

$$f_{\text{Artin}}(\chi_\rho) = \text{ord}_{\mathfrak{p}_K}(\mathfrak{d}_{E_1/K}) + [\kappa_{E_1} : \kappa_K]f_{\text{Artin}}(\psi),$$

which yields

$$f_{\text{Artin}}(\chi_\rho) = \mathfrak{d}_{E_1/K} f_{\text{Artin}}(\psi)^{[\kappa_{E_1} : \kappa_K]} = \mathfrak{d}_{E_1/K} \mathfrak{f}(E_1^{\ker(\psi)}/E_1)^{[\kappa_{E_1} : \kappa_K]}.$$

References

- [1] K. I. Ikeda, *A note on totally-ramified, n-abelian extensions of a local field*, preprint
- [2] K. I. Ikeda and M. G. Ikeda, *Two lemmas on formal power series*, Tr. J. of Math. (to appear).
- [3] K. Iwasawa *Local Class Field Theory*, Oxford University Press, Oxford-New York, 1986
- [4] H. Koch and E. de Shalit, *Metabelian local class field theory*, J. reine angew. Math. **478** (1996), 85-106.
- [5] J. Neukirch, *Algebraic Number Theory*, Springer-Verlag, Berlin, Heidelberg, New York 1999.
- [6] J.-P. Serre, *Corps Locaux*, Hermann, Paris, 1962
- [7] ———, *Représentations Linéaires des Groupes Finis*, Hermann, Paris, 1971

İKEDA

- [8] J. Tate *Number-theoretic background*, Automorphic Forms, Representations, *L*-functions, Proc. Sympos. Pure Math. vo. 33 (part 2) Amer. Math. Soc. Providence, Rhode Island 1979 1-XX.

Kazım İlhan İKEDA
Feza Gürsey Institute, P.O Box 6, 81220
Çengelköy, İstanbul, TURKEY

Received 17.07.2000

email: ilhan@gursey.gov.tr
Institut de Mathématiques,
“Théorie des Groupes,
Représentations et Applications”,
Université Paris 7, Case 7012, 2, place
Jussieu, 75251 Paris
Cedex 05, FRANCE
email: ilhan@math.jussieu.fr