

1-1-1999

Left Adjoint of Pullback Cat^1 -Groups

MURAT ALP

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

ALP, MURAT (1999) "Left Adjoint of Pullback Cat^1 -Groups," *Turkish Journal of Mathematics*: Vol. 23: No. 2, Article 2. Available at: <https://journals.tubitak.gov.tr/math/vol23/iss2/2>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

LEFT ADJOINT OF PULLBACK Cat^1 -GROUPS

Murat Alp

Abstract

In [1] we define the pullback Cat^1 -groups and showed that the category of pullback Cat^1 -groups is equivalent to the category of pullback crossed modules. In this paper we proved that the pullback Cat^1 -group has a left adjoint which is the induced Cat^1 -group. We also give the left adjoint construction.

1991 A. M. S. C.: 13D99, 16A99, 17B99, 17D99, 18D35.

Key words: Crossed modules, cat^1 -groups, pullback, Cocomplete category, Adjoint.

1. Introduction

Crossed modules are usefully regarded as 2-dimensional forms of groups. They were introduced by J. H. C. Whitehead in [13], and have powerful topological applications [5, 6, 7, 12]. Loday in [8] showed that the category of crossed modules is equivalent to that of cat^1 -groups. We implemented crossed modules and cat^1 -groups structures to the computed using the group theory language GAP [10] as a package in [11]. We also enumerated cat^1 -groups of low order and group order 41-47 in [2] and [1] using this program package XMOD.

Our aim is to define pullback cat^1 -groups and to show that the equivalence between cat^1 -groups and crossed modules due to Loday [5] takes pullback cat^1 -groups to the pullback crossed modules defined by Brown and Higgins in [3].

2. Pre-cat¹-groups and Pullback cat¹-groups

A crossed module $\chi = (\partial : S \rightarrow R)$ consists of a group homomorphism ∂ , called the *boundary of χ* , together with an action $\alpha : R \rightarrow \text{Aut}(S)$ satisfying, for all $s, s' \in S$ and $r \in R$,

$$\mathbf{XM1} : \partial(s^r) = r^{-1}(\partial s)r$$

$$\mathbf{XM2} : s^{\partial s'} = s'^{-1}ss'.$$

The standard examples of crossed modules are:

1. Any homomorphism $\partial : S \rightarrow R$ of abelian groups with R acting trivially on S may be regarded as a crossed module.
2. A conjugation crossed module is an inclusion of a normal subgroup $S \trianglelefteq R$, where R acts on S by conjugation.
3. A central extension crossed module has as boundary a surjection $\partial : S \rightarrow R$ with central kernel, where $r \in R$ acts on S by conjugation with $\partial^{-1}r$.
4. An automorphism crossed module has as its range a subgroup R of the automorphism group $\text{Aut}(S)$ of S which contains the inner automorphism group of S . The boundary maps $s \in S$ to the inner automorphism of S by s .
5. An R -module crossed module has an R -module as source and ∂ as the zero map.
6. The direct product $\chi_1 \times \chi_2$ of two crossed modules has source $S_1 \times S_2$, range $R_1 \times R_2$ and boundary $\partial_1 \times \partial_2$, with R_1, R_2 acting trivially on S_2, S_1 respectively.
7. An important motivating topological example of crossed module due to Whitehead [12] is the boundary $\partial : \pi_2(X, A, x) \rightarrow \pi_1(A, x)$ from the second relative homotopy group of a based pair (X, A, x) of topological spaces, with the usual action of the fundamental group $\pi_1(A, x)$.

A morphism between two crossed modules $\chi = (\partial : S \rightarrow R)$ and $\chi' = (\partial' : S' \rightarrow R')$ is a pair (σ, ρ) , where $\sigma : S \rightarrow S'$ and $\rho : R \rightarrow R'$ are homomorphisms satisfying

ALP

$$\partial' \sigma = \rho \partial, \quad \sigma(s^r) = (\sigma s)^{\rho r}.$$

In [8], Loday reformulated the notion of a crossed modules as a cat^1 -group, namely a group G with a pair of homomorphisms $t, h : G \rightarrow G$ having a common image R and satisfying certain axioms. We find it convenient to define a pre- cat^1 -group $\mathcal{C} = (e; t, h : G \rightarrow R)$ as a group G with two surjections $t, h : G \rightarrow R$ and an embedding $e : R \rightarrow G$ satisfying:

$$\mathbf{CAT1} : \quad te = he = id_R.$$

The pre- cat^1 -group $\mathcal{C} = (e, t, h : G \rightarrow R)$ is a cat^1 -group if it also satisfies

$$\mathbf{CAT2} : \quad [kert, kerh] = \{1_G\}.$$

The maps t, h are often called the source and target, but we choose to call them tail and head of \mathcal{C} , because source is the GAP term for the domain of a function.

A morphism $\mathcal{C} \rightarrow \mathcal{C}'$ of cat^1 -groups is a pair (γ, ρ) where $\gamma : G \rightarrow G'$ and $\rho : R \rightarrow R'$ are homomorphisms satisfying

$$h' \gamma = \rho h, \quad t' \gamma = \rho t, \quad e' \rho = \gamma e.$$

To any pre- cat^1 -group \mathcal{P} there is a canonically associated a cat^1 -group \mathcal{C} , obtained by quotienting the source group by the Peiffer subgroup $[ker t, ker h]$.


The corresponding functor is denoted

$$\mathbf{ass} : (\text{pre-}\text{cat}^1\text{-groups}) \rightarrow (\text{cat}^1\text{-groups}), \quad (0.1)$$

and is clearly the identity when restricted to cat^1 -group [6]

A pullback cat^1 -group is defined in [1] as follows.

ALP



Let $\mathcal{C} = (e; t, h : G \rightarrow R)$ be a cat^1 -group and let $\iota : Q \rightarrow R$ be a group homomorphism. Define $e^{**}; t^{**}, h^{**} : i^{**}G \rightarrow Q$ to be the pullback of G where

$$i^{**}G = \{(q_1, g, q_2) \in Q \times G \times Q \mid \iota q_1 = tg, \iota q_2 = hg\},$$

$t^{**}(q_1, g, q_2) = q_1, h^{**}(q_1, g, q_2) = q_2$ and $e^{**}(q) = (q, eiq, q)$. Multiplication in $i^{**}G$ is componentwise. The pair (π, ι) is a morphism of cat^1 -groups where $\pi : i^{**}G \rightarrow G, (q_1, g, q_2) \mapsto g$.

Proposition 2.1 [1] If $i^*\chi$ is the pullback of the crossed module χ over $\iota : Q \rightarrow R$ and if \mathcal{C}, \mathcal{D} are the cat^1 -groups obtained from $\chi, i^*\chi$ respectively, then $\mathcal{D} \cong i^{**}\mathcal{C}$.

3. Construction of the left adjoint

Proposition 3.1. The category of cat^1 -groups is co-complete.

Proof. Let $F : \mathcal{C} \rightarrow (\text{cat}^1\text{-groups})$. We wish to construct $\text{colim } F$. For each object c of \mathcal{C} , we write

$$F(c) = (e_c; t_c, h_c : F_1(c) \rightarrow F_0(c)).$$

Then we form

$$\begin{aligned} F' &= (i'; t', h' : \text{colim}_c F_1(c) \rightarrow \text{colim}_c F_0(c)) \\ &= (e'; t', h' : F'_1 \rightarrow F'_0), \end{aligned}$$

where F'_1, F'_0 are the colimits in the category of groups, so that $t'e' = h'e' = 1$. So F' is a pre- cat^1 -groups. The required colimit is the the associated cat^1 -group $\mathbf{ass} F'$ (see (1) on page 4),

$$\mathbf{ass}F' = (e''; t'', h'' : F'_1/[\ker t', \ker h'] \rightarrow F'_0).$$

□

We recall the definition of pushouts in a general category. Suppose we are given a commutative diagram of morphisms in a category \mathbf{C} :

$$\begin{array}{ccc} X_0 & \xrightarrow{i_1} & X_1 \\ \downarrow i_2 & & \downarrow v_1 \\ X_2 & \xrightarrow{v_2} & X \end{array}$$

Recall [9] that (v_1, v_2) is pushout of (i_1, i_2) , and also that the above square is a pushout square, if the following property holds: if $f_1 : X_1 \rightarrow H$, $f_2 : X_2 \rightarrow H$ are morphisms such that $f_1 i_1 = f_2 i_2$ then there is a unique map $f : X \rightarrow H$ such that $f v_1 = f_1$, $f v_2 = f_2$.

As usual, this property characterizes the pair (v_1, v_2) up to an automorphism of X . For this reason, it is common to make an abuse of language and refer to X as the pushout of (i_1, i_2) . In this case, we write

$$X = X_2 *_{X_0} X_1,$$

where $*_{X_0}$ is used to suggest a free product.


Proposition 3.2. *The functor $\iota^{**} : \text{Cat}^1 \text{ Grp}/U \rightarrow \text{Cat}^1 \text{ Grp}/R$ has a left adjoint $\iota_{**} : \text{Cat}^1 \text{ grp}/R \rightarrow \text{Cat}^1 \text{ Grp}/U$.*

Proof. We can give the left adjoint construction as follows. Let $\mathcal{C} = (e; t, h : G \rightarrow R)$ be at cat^1 -group over R and $\iota : R \rightarrow U$ is a morphism of groups. Then the induced cat^1 -group is $\iota_{**}\mathcal{C} = (e_{**}; t_{**}.h_{**} : \iota_{**}G \rightarrow U)$ is given by the pushout

ALP

$$\begin{array}{ccc}
 (1;1,1:R \rightarrow R) & \longrightarrow & (1;1,1:U \rightarrow U) \\
 \downarrow & & \downarrow \\
 (e;t,h:G \rightarrow R) & \longrightarrow & (e_{**}, \iota_{**}, h_{**}: \iota_{**} G \rightarrow U)
 \end{array}$$

we draw the above diagram as a three dimensional diagram as follows



in the category of cat^1 -groups. For computational purposes note that by the previous proposition 3.1, $\iota_{**}G = (G *_R U)/[\ker t_{**}, \ker h_{**}]$, where $*_R$ denotes coproduct of groups, that is, a free product with amalgamation over R . \square

References

- [1] Alp, M., Pullback of crossed modules and cat^1 -groups, *Turkish Journal of Mathematics* (to appear) 1998, 3 Vol 22.
- [2] Alp, M. Special case of Cat^1 -groups, *Algebras, Groups and Geometries*, Vol.15 (1998) (to appear).
- [3] Alp, M., Enumeration of Cat^1 -groups of group order 41-47, *Anadolu Üniversitesi Fen Fakültesi Dergisi*, 1997, Sayı 3 79-90. 4
- [4] Alp, M. and Wensley, C. D., Enumeration of Cat^1 -groups of low order, U. W. B. preprint, 96.05, 15pp *International Journal of algebra and Computation* (republished). 5

ALP

- [5] Brown, R. and Higgins, P. J., On the connection between the second relative homotopy groups of some related space, *Proc. London Math. Soc.*, (3) 36 (1978) 193-212.
- [6] Brown, R. and Loday, J. L., Homotopical excision, and Hurewicz theorems, for n-cubes of spaces, *Proc. London Math. Soc.*, (3) 54 (1987) 176-192.
- [7] Brown, R. and Wensley, C. D. On finite induced crossed modules. and the homotopy 2-type of mapping cones, *Theory and applications of Categories*, 1 (3) (1995)54-71.
- [8] Loday, J. L., Spaces with finitely many non-trivial homotopy groups, *J. App. Algebra*, 24 (1982) 179-202.
- [9] MacLane, S., *Categories for the Working Mathematician*, Graduate Texts in Maths., Springer Verlag, Berlin 5 (1971).
- [10] Schönert, M. et al, GAP: groups, Algorithms, and Programming, Lehrstuhl D für mathematik, Rheinisch Westfälische Technische Hochschule, Aachen, Germany, third edition, 1993.
- [11] Alp, M., and Wensley, C. D., XMOD, crossed modules and cat^1 -groups in GAP, version 1.3 *Manual for the XMOD share package*, (1996) 1-80.
- [12] Whitehead, J. H. C., on adding relations to homotopy groups, *Ann. Math.*,47 (1946) 806-810.
- [13] Whitehead, J. H. C., Combinatorial homotopy II, *Bull. a. M. S.*, 55 (1949) 453-496.

Murat ALP

Received 19.11.1997

Dumyupınar Üniversitesi Fen-Edebiyat Fakültesi
Matematik Bölümü Kütahya-TURKEY
email: malp@dumlupinar.edu.tr