

1-1-1999

Normal Subgroups and Elements of $H^{\{\}}(\lambda_q)$

İSMAİL NACİ CANGÜL

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

CANGÜL, İSMAİL NACİ (1999) "Normal Subgroups and Elements of $H^{\{\}}(\lambda_q)$," *Turkish Journal of Mathematics*: Vol. 23: No. 2, Article 3. Available at: <https://journals.tubitak.gov.tr/math/vol23/iss2/3>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

NORMAL SUBGROUPS AND ELEMENTS OF $H'(\lambda_q)$

İsmail Naci Cangül

Abstract

In this study, we consider the normal subgroups of $H'(\lambda_q)$, where $H(\lambda_q)$ denotes the Hecke groups. After recalling some results from [2], particularly on the group structure and on the relations with the power subgroups of $H(\lambda_q)$, the even subgroup $H_e(\lambda_q)$ of $H(\lambda_q)$ is discussed. It is shown that $H'(\lambda_q)$ is a normal subgroup of $H_e(\lambda_q)$ with index q . For this reason each subgroup of $H'(\lambda_q)$ consists of only even elements. $H''(\lambda_q)$ is also considered and it is concluded that it is the normal subgroup of $H'(\lambda_q)$ generated by all commutators of the elements of $H'(\lambda_q)$. Using the Kurosh subgroup theorem, the group structure of normal subgroups of $H(\lambda_q)$ can be found to be free groups. Their ranks are given in terms of the index.

1. Introduction

Let Γ be the classical modular group. Let R and S be its generators of order 2 and 3, respectively, defined by

$$R(z) = -\frac{1}{z} \quad \text{and} \quad S(z) = -\frac{1}{z+1}, \quad (1)$$

A generalisation of Γ is known as Hecke groups denoted by $H(\lambda_q)$ and generated by two elements

$$R(z) = -\frac{1}{z} \quad \text{and} \quad S(z) = -\frac{1}{z+\lambda_q}, \quad (2)$$

of order 2 and q , respectively, where $\lambda_q = 2 \cos \pi/q$, $q \in N$, $q \geq 3$.

In [2], the group structure of $H'(\lambda_q)$, the commutator subgroup of $H(\lambda_q)$, is obtained and some of its properties are given. In particular it is shown that $H'(\lambda_q)$ is a free group of rank $q-1$ and also normal with index $2q$ in $H(\lambda_q)$. This coincides with the result given for the modular group where $q = 3$, [4]. We define $H^m(\lambda_q)$ to be the subgroup of $H(\lambda_q)$ generated by the m -th powers of the elements of $H(\lambda_q)$. $H^m(\lambda_q)$ is called the m -th power subgroup of $H(\lambda_q)$. It is also normal in $H(\lambda_q)$.

Relations between the modular group Γ , its commutator subgroup Γ' and power subgroups of Γ are discussed in [5]. These are generalised to some Hecke groups in [2].

2. Elements of $H'(\lambda_q)$ and its Relation with the Even Subgroup

We now study the place of $H'(\lambda_q)$ amongst the normal subgroups of $H(\lambda_q)$. First we look at the elements of $H(\lambda_q)$. They form two classes:

$$(a) \quad \begin{pmatrix} a & b\lambda_q \\ c\lambda_q & d \end{pmatrix} ad - bc\lambda_q^2 = 1, \tag{3}$$

$$(b) \quad \begin{pmatrix} a\lambda_q & b \\ c & d\lambda_q \end{pmatrix} ad\lambda_q^2 - bc = 1, \tag{4}$$

where a, b, c, d are all polynomials of λ_q^2 with rational integer coefficients. The elements of type (a) are called even while those of type (b) are called odd. It is easy to see that

$$odd \cdot odd = even \cdot even = even \tag{5}$$

$$even \cdot odd = odd \cdot even = odd.$$

When q is even, the even elements form a subgroup of $H(\lambda_q)$ of index 2 called the even subgroup, denoted by $H_e(\lambda_q)$:

$$H_e(\lambda_q) = \left\{ M = \begin{pmatrix} a & b\lambda_q \\ c\lambda_q & d \end{pmatrix} : M \in H(\lambda_q) \right\}. \tag{6}$$

It is generated by $T = R \cdot S$ and $T \cdot U = RS^2R$ and in fact

CANGÜL

$$H_e(\lambda_q) \cong \langle T \rangle * \langle T \cdot U \rangle \quad (7)$$

Being of index 2, $H_e(\lambda_q)$ is normal in $H(\lambda_q)$. The set of odd elements form the other coset of $H_e(\lambda_q)$ in $H(\lambda_q)$. Actually

$$H(\lambda_q) = H_e(\lambda_q) \cup R \cdot H_e(\lambda_q) \quad (8)$$

as $R \notin H_e(\lambda_q)$, (see [3]).

When q is odd, $H(\lambda_q)$ does not have an even subgroup.

We now see the connection between the even subgroup and $H'(\lambda_q)$ for even q :

Theorem 2.1. *Let q be even. Then the commutator subgroup $H'(\lambda_q)$ of $H(\lambda_q)$ is a normal subgroup of the even subgroup $H_e(\lambda_q)$ with index q .*

Proof. Recall that $H'(\lambda_q)$ is a normal subgroup of $H(\lambda_q)$ with index $2q$. The even subgroup $H_e(\lambda_q)$, having index 2, is also normal in $H(\lambda_q)$. Therefore the required index is q .

Let us take two elements A, B of $H(\lambda_q)$. Note that whatever A and B are, their commutator $[A, B] = ABA^{-1}B^{-1}$ is always even. Hence for every pair of elements

$$[A, B] \in H_e(\lambda_q). \quad (9)$$

That is

$$H'(\lambda_q) \triangleleft H_e(\lambda_q). \quad (10)$$

□

Corollary 2.1. *A subgroup of $H'(\lambda_q)$ consists of only even elements.*

3. Second Commutator Subgroup

Lemma 3.1. *Let G be a finitely generated group and H be the subgroup of G generated by all the commutators of the generators of G . Then G/H is abelian.*

Proof. Let aH, bH be two elements of G/H . Then as $b^{-1}a^{-1}ba \in H$ we have

$$aH \cdot bH = abH = ab(b^{-1}a^{-1}ba)H = baH = bH \cdot aH$$

which proves the lemma. □

Theorem 3.1. *Let $H'(\lambda_q)$ be generated by the $q-1$ elements a_1, a_2, \dots, a_{q-1} . Let N be the normal subgroup of $H'(\lambda_q)$ generated by all the commutators $[a_i, a_j]$ of the generators. Then*

$$N = H''(\lambda_q).$$

Proof. By Lemma 3.1, $H'(\lambda_q)/N$ is abelian. But it is well-known that $H'(\lambda_q)/H''(\lambda_q)$ is the largest abelian quotient group of $H'(\lambda_q)$. As $N < H''(\lambda_q)$, the result follows. □

4. The Structure of Normal Subgroups of $H'(\lambda_q)$

Recall that $H'(\lambda_q)$ is a free group of rank $q-1$. By the Kurosh subgroup theorem, all of its subgroups will be free. Therefore we only need to find the rank of a normal subgroup. We use the following result:

Theorem 4.1. *Let H be a subgroup of finite index μ in $H'(\lambda_q)$. Then the rank r of H will also be finite and can be found by the formula*

$$r = 1 + \mu \cdot (q - 2). \tag{11}$$

CANGÜL

Proof. We need two results. First we use the fact that the rank of a free group of genus g having t parabolic classes is equal to $2g + t - 1$, (see [1]). We also need the Riemann-Hurwitz formula

$$\mu = \frac{\mu(H)}{\mu(H'(\lambda_q))}, \quad (12)$$

where $2\pi \cdot \mu(H)$ denotes the hyperbolic area of a fundamental region for H given by

$$\mu(H) = 2g - 2 + \sum_{i=1}^r \left(1 - \frac{1}{m_i} \right) + t. \quad (13)$$

Here m_i are the orders of finite ordered elements in H . Using these two results, one can easily prove the theorem. \square

References

- [1] Cangül, İ. N.: Determining Isomorphism class of a Fuchsian Group from its Signature , (to be printed).
- [2] Cangül, İ.N.: Commutator Subgroups of Hecke Groups, (to be printed).
- [3] Cangül, İ.N., Singerman, D.: Normal Subgroups of Hecke Groups and Regular Maps, Math. Proc. Cambridge Phil. Soc., 123 (1998), 59-74.
- [4] Newman, M.: A note on the Modular Groups, Proc. A.M.S. 14 (1963), 124-125.
- [5] Newman, M.: The Structure of some Subgroups of the Modular Group, Illinois J. of Math. 6 (1962), 480-487.

İsmail Naci CANGÜL
Uludağ Üniversitesi, Fen Fakültesi,
Matematik Bölümü, Görükle
16059 Bursa-TÜRKİYE
cangul@uludag.edu.tr

Received 25.11.1997