

1-1-2014

Detection the frequency of PER-1 type extended-spectrum β -lactamase-producing *Acinetobacter baumannii* clinical isolates in Turkey: a multicenter study

GÜLŞAH AŞIK

MEHMET ÖZDEMİR

MUHAMMET GÜZEL KURTOĞLU

SERVER YAĞCI

LÜTFİYE ÖKSÜZ

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>

 Part of the [Medical Sciences Commons](#)

Recommended Citation

AŞIK, GÜLŞAH; ÖZDEMİR, MEHMET; KURTOĞLU, MUHAMMET GÜZEL; YAĞCI, SERVER; ÖKSÜZ, LÜTFİYE; GÜL, MUSTAFA; KOÇOĞLU, MÜCAHİDE ESRA; ÇETİN, EMEL SESLİ; SEYREK, ADNAN; BERKTAŞ, MUSTAFA; AYYILDIZ, AHMET; and ÇİFTÇİ, İHSAN HAKKI (2014) "Detection the frequency of PER-1 type extended-spectrum β -lactamase-producing *Acinetobacter baumannii* clinical isolates in Turkey: a multicenter study," *Turkish Journal of Medical Sciences*: Vol. 44: No. 6, Article 19. <https://doi.org/10.3906/sag-1309-126>
Available at: <https://journals.tubitak.gov.tr/medical/vol44/iss6/19>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Detection the frequency of PER-1 type extended-spectrum β -lactamase-producing *Acinetobacter baumannii* clinical isolates in Turkey: a multicenter study

Authors

GÜLŞAH AŞIK, MEHMET ÖZDEMİR, MUHAMMET GÜZEL KURTOĞLU, SERVER YAĞCI, LÜTFİYE ÖKSÜZ, MUSTAFA GÜL, MÜCAHİDE ESRA KOÇOĞLU, EMEL SESLİ ÇETİN, ADNAN SEYREK, MUSTAFA BERKTAŞ, AHMET AYYILDIZ, and İHSAN HAKKI ÇİFTÇİ

Detection of the frequency of PER-1 type extended-spectrum β -lactamase-producing *Acinetobacter baumannii* clinical isolates in Turkey: a multicenter study

Gülşah AŞIK^{1*}, Mehmet ÖZDEMİR², Muhammet Güzel KURTOĞLU³, Server YAĞCI⁴, Lütfiye ÖKSÜZ⁵, Mustafa GÜL⁶, Mücahide Esra KOÇOĞLU⁷, Emel SESLİ ÇETİN⁸, Adnan SEYREK⁹, Mustafa BERKTAŞ¹⁰, Ahmet AYYILDIZ¹¹, İhsan Hakkı ÇİFTÇİ¹²

¹Department of Microbiology, Faculty of Medicine, Afyon Kocatepe University, Afyonkarahisar, Turkey

²Department of Microbiology, Faculty of Medicine, Necmettin Erbakan University, Konya, Turkey

³Department of Microbiology, Konya Training and Research Hospital, Konya, Turkey

⁴Department of Infectious Diseases and Clinical Microbiology, Ministry of Health, Ankara Training and Research Hospital, Ankara, Turkey

⁵Department of Microbiology, Faculty of Medicine, İstanbul University, İstanbul, Turkey

⁶Department of Microbiology, Faculty of Medicine, Sütçü İmam University, Kahramanmaraş, Turkey

⁷Department of Microbiology, Faculty of Medicine, Abant İzzet Baysal University, Bolu, Turkey

⁸Department of Microbiology, Faculty of Medicine, Süleyman Demirel University, Isparta, Turkey

⁹Department of Microbiology, Faculty of Medicine, Fırat University, Elazığ, Turkey

¹⁰Department of Microbiology, Faculty of Medicine, Yüzüncü Yıl University, Van, Turkey

¹¹Department of Microbiology, Faculty of Medicine, Atatürk University, Erzurum, Turkey

¹²Department of Microbiology, Faculty of Medicine, Sakarya University, Sakarya, Turkey

Received: 27.09.2013 • Accepted: 03.02.2014 • Published Online: 24.10.2014 • Printed: 21.11.2014

Background/aim: β -Lactamases are an important resistance mechanism in *Acinetobacter baumannii*. Pseudomonas extended-resistance (PER-1) type β -lactamase-producing strains have been reported from various geographic locations; however, PER-1 type β -lactamases from Turkish hospitals have not been investigated extensively. The aim of this study was to determine the prevalence of PER-1 type β -lactamases in *A. baumannii* isolates in various regions of Turkey.

Materials and methods: A total of 763 clinical *A. baumannii* isolates were collected from 9 university hospitals and 2 state hospitals between 2008 and 2011. Molecular amplification of the OXA-51 gene from the *A. baumannii* genome was performed in order to verify identification of the species. Real-time polymerase chain reaction was used to detect *bla*_{PER-1} genes.

Results: PER-1 was detected in 24.6% of the isolates. The annual frequencies of the PER-1 enzyme were detected as 52.2%, 35.9%, and 8.3% in 2008, 2009, and 2010, respectively. PER-1 prevalence decreased gradually over time. The differences observed in PER-1 prevalence among the regions of Turkey were statistically significant (chi-square test; $P < 0.001$).

Conclusion: These data demonstrate that the frequency of detection of PER-1 type β -lactamases in *A. baumannii* species has decreased in Turkey. However, the increased carbapenem resistance, together with multidrug resistance, has created a worrisome situation regarding this pathogen.

Key words: *Acinetobacter baumannii*, β -lactamases, PER-1

1. Introduction

Since members of the genus *Acinetobacter* were realized to be significant nosocomial pathogens, much information has been learned. In the first in vitro studies, most clinical isolates were susceptible to generally used antimicrobial agents so that infections caused by these organisms could be treated relatively easily (1). However, successive surveys have shown increasing resistance among clinical isolates, particularly those belonging to the *Acinetobacter*

baumannii complex, and high proportions of isolates are now resistant to the most commonly used antimicrobial agents.

Over the past decade, nosocomial outbreaks of *A. baumannii* have been described with increasing frequency, mostly in association with intensive care units, burn units, or surgical wards (2). Resistance to β -lactam antibiotics in *Acinetobacter* spp. predominantly involves 3 mechanisms: production of β -lactamases, loss of outer

* Correspondence: gulsahmet@hotmail.com

membrane proteins, and upregulation of efflux pumps (2). The most common mechanism of resistance to β -lactam antibiotics is the production of β -lactamases (3). Acquired β -lactamases encoded by mobile genetic elements are an important resistance mechanism in *A. baumannii* (4).

One of the β -lactamases, *Pseudomonas* extended resistance (PER-1), was first identified in 1991 in a *Pseudomonas aeruginosa* strain isolated from a Turkish patient in France (5). The bla_{PER-1} gene is widespread in Turkey, particularly among *P. aeruginosa* and *Acinetobacter* spp. strains (6–8). Subsequently, PER-1-producing strains have been reported from various geographic regions (6–8). However, PER-1 type β -lactamases among clinical isolates from Turkish hospitals have not been investigated extensively. Furthermore, the current prevalence of PER-1 in the various regions of Turkey remains unknown. Therefore, the aim of this study was to determine the prevalence of PER-1 type β -lactamases in *A. baumannii* isolates from 11 hospitals in Turkey.

2. Materials and methods

A total of 763 nonrepetitive *A. baumannii* clinical isolates archived from 9 university hospitals and 2 state hospitals in Turkey from 2008 to 2011 were included in the study. The provinces that participated in the study are shown in the Figure. Antimicrobial susceptibility tests and molecular studies were performed in the microbiology laboratories of Afyon Kocatepe University. The isolates were identified by means of both conventional techniques (including oxidation–fermentation reactions in triple sugar iron agar, oxidase production, and colony appearance) and automated systems, including the VITEK system (bioMérieux VITEK System Inc.; bioMérieux, Marcy l’Etoile, France) and the Phoenix 100 system (Becton Dickinson and Company, Franklin Lakes, NJ, USA). The OXA-51 gene region is species-specific for *A. baumannii* (4). The identification of species was verified by the presence of the bla_{OXA-51} gene within the *A. baumannii* genome.

The susceptibilities of the isolates to imipenem, meropenem, cefepime, cefoperazone-sulbactam,

gentamicin, amikacin, and tobramycin were tested using the standard disk diffusion method on Mueller Hinton (MH) agar plates (Oxoid Ltd., Basingstoke, UK) and using the breakpoints defined by the Clinical and Laboratory Standards Institute (9). Antimicrobials were applied and stored according to the manufacturer’s instructions. *A. baumannii* ATCC 19606 was used as a reference strain for susceptibility tests.

For DNA extraction, a loopful of bacteria was taken from each fresh overnight culture on MH agar plates. DNA samples were extracted by incubating a pure bacterial suspension at 95 °C for 10 min in a lysis buffer, and debris was removed by centrifugation for 5 min at 12,000 \times g. The quantity and quality of the extracted DNA were assessed using a NanoDrop ND-1000 spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). The DNA concentration and the ratio of the optical density at 260/280 nm to evaluate the purity of the DNA samples were calculated simultaneously (10).

Based on sequences of bla_{PER-1} and bla_{OXA-51} of *A. baumannii* retrieved from the National Center for Biotechnology Information Entrez database, oligonucleotide primer sets PER-1F (5'- ATG AAT GTC ATT ATA AAA GC-3'), PER-1R (5'-AAT TTG GGC TTA GGG CAG AA -3'), OXA-51F (5'-TCA GCA AGA GGC ACA GTT TG-3'), and OXA-51R (5'-GCT GAA CAA CCC ATC CAG TT-3') were designed and used to amplify a single polymerase chain reaction (PCR) product of 925 bp and a 188-bp fragment, respectively. The annealing temperatures were 52 °C and 54 °C for the PER-1 and OXA-51 primer pairs, respectively. Conventional assays were used to optimize the PCR reaction using a TECHNE-TC-512 thermal cycler (Barloworld Scientific, Burlington NJ, USA) for detection of bla_{PER-1} and bla_{OXA-51} genes.

Conventional PCR reactions with genomic DNA were performed in a 25- μ L mixture according to the manufacturer’s protocol for the Maxima SYBR Green/ROX qPCR Master Mix (Thermo Fisher Scientific). The PCR reaction steps were as follows: preheating at 95 °C



Figure. The locations of study centers within Turkey.

for 10 min; 35 cycles consisting of 95 °C for 30 s, 52 °C for PER-1 and 54 °C for OXA-51 30 s, and 72 °C for 1 min; and a final extension incubation at 72 °C for 4 min. After optimization, PCR assays were completed using the real-time method (SLAN, Shanghai Odin Science & Technology Co, Shanghai, China). Real-time PCR reactions were performed in a 25-µL mixture according to the manufacturer's protocol for the PCR master mix. The real-time PCR reaction steps were as follows: initial denaturation at 95 °C for 5 min; 30 cycles consisting of 95 °C for 30 s, 52 °C for 30 s, and 72 °C for 1 min; and a final extension incubation at 72 °C for 2 min.

To confirm the PCR product, melting curve analysis and 1.0% agarose gel electrophoresis were performed. The chi-square test was used to compare proportions. $P < 0.05$ was deemed to indicate statistical significance. Statistical analyses were conducted using the SPSS 17 (SPSS Inc., Chicago, IL, USA).

3. Results

A total of 763 unique *A. baumannii* strains had been identified previously in clinical microbiology laboratories at study centers between 2008 and 2011. All of the 763 strains tested positive using primers specific for OXA-51. Among the isolates, the rates of resistance were as follows: 100% for the third-generation of cephalosporin (ceftazidime, ceftriaxone, and cefotaxime) 94.8% for ceftipime, 73.6% for cefoperazone/sulbactam, 76.3% for meropenem, 74.5% for imipenem, 77.1% for gentamicin, 75.3% for amikacin, and 36.5% for tobramycin. Although the frequency of resistance to all antibiotics was high, especially against carbapenems when grouped according to year, there were no significant differences among the rates of antimicrobial resistance in the various locations in Turkey (chi-square test; $P > 0.05$). The regions, years, and antimicrobial resistance distributions of *A. baumannii* isolates are summarized in the Table.

Table. PER-1 frequencies and the antimicrobial resistance profiles of *A. baumannii* isolates from various hospitals in Turkey.

City	Year	Location in Turkey	n	PER-1 (+) n/%	PER-1 (-) n/%	Antimicrobial resistance rate. %						
						IPM	MEM	FEP	SCF	GEN	AMK	TOB
*Afyonkarahisar	2008	West	42	10/23.8	32/76.2	47.6	54.8	100	52.3	83.3	71.4	64.2
*Kahramanmaraş	2008	Southeast	32	15/46.9	17/53.1	65.6	65.6	87.5	50.0	71.8	68.7	12.5
*Van	2008	East	62	46/74.2	16/25.8	45.2	50.1	95.1	73.5	90.3	70.9	9.7
*Afyonkarahisar	2009	West	79	21/26.6	58/73.4	60.8	60.8	97.5	67.1	48.1	48.1	32.9
*İstanbul	2009	Marmara	41	9/22.0	32/78.0	92.6	92.6	100	73.8	85.3	80.5	51.2
**Konya	2009	Middle	65	21/32.3	44/67.7	93.8	92.3	100	92.3	72.3	72.3	53.8
*Bolu	2009	North	43	31/72.1	12/27.9	81.4	81.4	100	88.3	100	88.3	32.5
*Afyonkarahisar	2010	West	105	1/0.9	104/99.1	83.8	82.9	97.1	61.9	80.0	66.6	39.0
**Ankara	2010	Middle	50	1/2.0	49/98.0	60.0	60.0	88.0	70.0	58.0	76.0	52.0
*Erzurum	2010	East	49	7/14.3	42/85.7	69.4	69.4	91.8	77.5	65.3	71.4	40.8
*Isparta	2010	South	94	11/11.7	83/88.3	72.3	76.6	82.9	60.6	76.6	88.3	20.2
*Konya	2010	Middle	74	11/14.9	63 / 85.1	98.7	98.7	98.7	97.3	93.2	79.7	52.7
*Van	2011	East	18	4/22.2	14 / 77.8	94.4	94.4	100	88.8	88.8	83.3	27.7
*Elazığ	2011	East	9	0/0	9 / 100	77.7	88.8	88.8	77.7	66.6	88.8	22.2
TOTAL			763	188/24.6	575/75.4	74.5	76.3	94.8	73.6	77.1	75.3	36.5

*: University hospital, **: state hospital, IPM: imipenem, MEM: meropenem, FEP: ceftipime, SCF: cefoperazone-sulbactam, GEN: gentamicin, AMK: amikacin, TOB: tobramycin.

The concentration and the purity of the extracted DNA from a representative study group of 25 randomly selected test samples were assessed using the NanoDrop ND-1000 spectrophotometer. The mean \pm standard deviation (SD) concentration of extracted DNA was 22.36 ± 7.11 ng/ μ L (minimum–maximum values, 8.97–49.37 ng/ μ L). The mean \pm SD purity of extracted DNA samples (A260/A280) was 1.69 ± 0.09 (minimum–maximum values, 1.60–1.85).

PER-1 was detected by PCR using specific primers in 188 (24.6%) of 763 isolates. The annual PER-1 detection frequencies were 52.2%, 35.9%, and 8.3% in 2008, 2009, and 2010, respectively. The rates of PER-1–producing isolates were 47.6%, 42.4%, 17.5%, 13.6%, and 11.7% in the northern, eastern, middle, western, and southern regions of Turkey, respectively. The rates of PER-1–producing isolates differed significantly among the regions of Turkey (chi-square test; $P < 0.001$). Furthermore, the rates of PER-1–producing isolates in Afyonkarahisar Province were 23.8%, 26.1%, and 1% in 2008, 2009, and 2010, respectively (chi-square test; $P < 0.001$); in Van Province they were 74.2% and 22.2% in 2008 and 2011, respectively (chi-square test; $P < 0.001$). There were statistically significant differences between the rates of isolation of PER-1–producing strains from different hospitals in the same city, and in different cities in the same region of Anatolia (chi-square test, $P = 0.017$). Other results are summarized in the Table.

4. Discussion

A. baumannii has become a major cause of hospital-acquired infections, because this pathogen is difficult to control due to its prolonged environmental survival. Moreover, treatment is complicated by its ability to develop resistance to multiple antibiotic agents. Thus, in the last 10 years, *A. baumannii* has emerged as a highly problematic pathogen because few antibiotics are effective against this organism (11).

A. baumannii has become resistant to almost all antimicrobial agents currently available, including broad-spectrum β -lactams and quinolones. Most strains are resistant to aminoglycosides and cephalosporins (11). Our results demonstrate that all isolates were resistant to third-generation cephalosporins. In addition, cefoperazone/sulbactam and aminoglycoside resistance rates (except those for tobramycin) were around 70% at the various centers in Turkey. Because of these very high resistance rates, these agent classes are unlikely to play an important role in the treatment of *A. baumannii* infections.

Among the 490 *A. baumannii* isolates from patients with serious infections in European hospitals participating in the 1997–2000 Meropenem Yearly Susceptibility Test Information Collection (MYSTIC) programs, the 2 carbapenems meropenem and imipenem showed the greatest clinically useful activity (12). Susceptibilities of *A. baumannii* to meropenem were very high (97%–100%)

in all countries except Italy (70%), Turkey (66%), and the United Kingdom (77%). Similar results were observed for imipenem (93%–100%), except for Italy (78%), Turkey (62%), and the United Kingdom (78%) (12). Baran et al. (13) reported that the imipenem resistance rate was lower than indicated in the MYSTIC report for Turkey (53.7%). These reports show that carbapenems (imipenem and meropenem) remain active against many strains; however, increasing numbers of clinical isolates of *A. baumannii* resistant to carbapenems are now being reported worldwide, reaching levels of $\geq 90\%$ at some centers (14,15). In the present study, we found that *A. baumannii* isolates have a high carbapenem resistance rate, which exceeded that in the MYSTIC report. Another important issue is the annual increase in rates of resistance; isolates from Afyonkarahisar, Konya, İstanbul, and Van provinces from the previous years exhibited particularly high rates of resistance. This could be associated with the increased isolation of multidrug-resistant (MDR) species and a change in antibiotic selection protocols. Thus, the use of carbapenem in the treatment of MDR *A. baumannii* infections has increased in recent years. As a result, a dramatic increase in carbapenem resistance has occurred (Table).

PER-1 β -lactamase has been considered to be significant only in Turkey for a number of years. However, the PER-1 β -lactamase has been detected in *P. aeruginosa* in many countries, including Turkey, France, Belgium, Spain, Italy, Poland, Romania, Japan, and South Korea (7,16–23). PER-1 production in *Acinetobacter* spp. has been reported most often in Turkey and Korea (24). PER-1 type β -lactamases in clinical isolates from Turkish hospitals have not been investigated extensively. Only 1 study by Hoşgör et al. (25) in İzmir was published and in that study the *bla*_{PER-1} gene was detected by PCR in 33 (19.5%) of a total of 169 gram-negative bacteria, including 17 (23.3%) of the 73 *P. aeruginosa* isolates and 16 (25%) of the 64 *A. baumannii* complex isolates. The rate of detection of PER-1 was available to date only for 274 *A. baumannii* isolates from 3 regions of Turkey. Nationwide surveys in Turkey revealed that 46% (1997), 37.8% (2001), 31% (2005), and 35.9% (2007) of *A. baumannii* isolates produced PER-1, as reported by Vahaboglu et al. (6,26), Kolayli et al. (7), and Erac and Gulay (8), respectively. The current prevalence and distribution of PER-1 in the various regions of Turkey are unknown. In the present study, the rate of detection of PER-1 in 763 *A. baumannii* clinical isolates collected from different regions, 10 cities, and 11 hospitals was 24.6% (minimum–maximum values, 0%–74.2%). The annual rate of PER-1 detection decreased gradually with time, with the most prominent decrease occurring in 2010. Furthermore, between 2008 and 2011, the resistance rates increased markedly. The data suggest that, in a hospital in which the prevalence of resistant *A. baumannii* is

increasing and clinical isolates demonstrate new antibiotic resistance patterns, it would be prudent to first consider the likelihood of transmission of these pathogens from exogenous sources. Another important reason for the decreased prevalence of PER-1 is transmission of mobile genes on plasmids among the *A. baumannii* population.

Differences in PER-1 prevalence have been observed among the regions of Turkey. PER-1 was prominent in isolates from Van Province in the eastern and Bolu Province in the northern region of Turkey. This difference in resistance profiles among regions has been expected by many researchers. We think that the regional differences identified in the present study can be attributed to changes in the patient population and treatment regimen preferences.

In conclusion, *A. baumannii* as a cause of nosocomial infections has shown a first-order upward trend in

recent years. The frequency of detection of PER-1 type extended-spectrum β -lactamases in *A. baumannii* species has decreased, and PER-1 is no longer a threat in terms of Turkey's resistance profile. However, the increased carbapenem resistance, together with MDR bacteria, represents a worrisome situation in this species. Thus, the mechanisms responsible for carbapenem resistance in *A. baumannii* isolates should be investigated. Moreover, strategies to prevent the spread of resistance should be instituted.

Acknowledgments

We thank staff member Şule Özkan of the Afyon Kocatepe University Microbiology Laboratory for contribution in the performing of the molecular tests. We also thank all the academics who sent *A. baumannii* strains to our laboratory for this study. There were no financial sources.

References

- Bergogne-Berezin E. Resistance of *Acinetobacter* spp. to antimicrobials: overview of clinical resistance patterns and therapeutic problems. In: Bergogne-Berezin E, Joly-Guillou ML, Towner KJ, editors. *Acinetobacter: Microbiology, Epidemiology, Infections, Management*. Boca Raton, FL, USA: CRC Press; 1996. pp. 133–183.
- Munoz-Price LS, Weinstein RA. *Acinetobacter* infection. *N Engl J Med* 2008; 358: 1271–1281.
- Towner KJ. Molecular basis of antibiotic resistance in *Acinetobacter* spp. In: Gerisher U, editor. *Acinetobacter Molecular Biology*. Norfolk, UK: Caister Academic Press; 2008. pp. 322–343.
- Brown S, Young HK, Amyes SGB. Characterisation of OXA-51, a novel class D cephalosporinase found in genetically unrelated clinical strains of *Acinetobacter baumannii* from Argentina. *Clin Microbiol Infect* 2005; 11: 15–23.
- Nordman P, Ronco E, Naas T, Dupont C, Michel Briand Y, Labia R. Characterization of a novel extended-spectrum β -lactamase from *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 1993; 37: 962–969.
- Vahaboglu H, Ozturk R, Aygun G, Coskuncan F, Yaman A, Kaygusuz A, Leblebicioglu H, Balik I, Aydin K, Oktun M. Widespread of PER-1-type extended-spectrum β -lactamases among nosocomial *Acinetobacter* and *Pseudomonas aeruginosa* isolates in Turkey: a nationwide multicenter study. *Antimicrob Agents Chemother* 1997; 41: 2265–2269.
- Kolayli F, Gacar G, Karadenizli A, Sanic A, Vahaboglu H. PER-1 is still widespread in Turkish hospitals among *Pseudomonas aeruginosa* and *Acinetobacter* spp. *FEMS Microbiol Lett* 2005; 249: 241–245.
- Erac B, Gulay Z. Molecular epidemiology of PER-1 extended spectrum β -lactamase among gram-negative bacteria isolated at a tertiary care hospital. *Folia Microbiol* 2007; 52: 535–541.
- Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing, 17th Informational Supplement. Wayne, PA, USA: Clinical and Laboratory Standards Institute; 2007.
- Bogner PN, Killeen AA. Extraction of nucleic acids. In: Coleman WB, Tsongalis GJ, editors. *Molecular Diagnostics: For the Clinical Laboratorian*. New York, NY, USA: Humana Press; 2007. pp. 25–30.
- Adams-Haduch JM, Paterson DL, Sidjabat HE, Pasculle AW, Potoski BA, Muto CA, Harrison LH, Doi Y. Genetic basis of multidrug resistance in *Acinetobacter baumannii* clinical isolates at a tertiary medical center in Pennsylvania. *Antimicrob Agents Chemother* 2008; 52: 3837–3843.
- Turner PJ, Greenhalgh JM; MYSTIC Study Group (Europe). The activity of meropenem and comparators against *Acinetobacter* strains isolated from European hospitals, 1997–2000. *Clin Microbiol Infect* 2003; 9: 563–567.
- Baran G, Erbay A, Bodur H, Onguru P, Akinci E, Balaban N, Cevik MA. Risk factors for nosocomial imipenem-resistant *Acinetobacter baumannii* infections. *Int J Infect Dis* 2008; 12: 16–21.
- Karageorgopoulos DE, Falagas ME. Current control and treatment of multidrug-resistant *Acinetobacter baumannii* infections. *Lancet Infect Dis* 2008; 8: 751–762.
- Kean SJ. Doripenem: a review of its use in the treatment of bacterial infections. *Drugs* 2008; 68: 2021–2057.
- Pagani L, Mantengoli E, Migliavacca R, Nucleo E, Pollini S, Spalla M, Daturi R, Romero E, Rossolini GM. Multifocal detection of multidrug-resistant *Pseudomonas aeruginosa* producing the PER-1 extended-spectrum β -lactamase in northern Italy. *J Clin Microbiol* 2004; 42: 2523–2529.

17. De Champs C, Poirel L, Bonnet R, Sirot D, Chanal C, Sirot J, Nordmann P. Prospective survey of beta-lactamases produced by ceftazidime-resistant *Pseudomonas aeruginosa* isolated in a French hospital in 2000. *Antimicrob Agents Chemother* 2002; 46: 3031–3034.
18. Naas T, Bogaerts P, Bauraing C, Degheldre Y, Glupczynski Y, Nordmann P. Emergence of PER and VEB extended-spectrum beta-lactamases in *Acinetobacter baumannii* in Belgium. *J Antimicrob Chemother* 2006; 58: 178–182.
19. Miro E, Mirelis B, Navarro F, Rivera A, Mesa RJ, Roig MC, Gomez L, Coll P. Surveillance of extended-spectrum beta-lactamases from clinical samples and faecal carriers in Barcelona, Spain. *J Antimicrob Chemother* 2005; 56: 1152–1155.
20. Empel J, Filczak K, Mrowka A, Hryniewicz W, Livermore DM, Gniadkowski M. Outbreak of *Pseudomonas aeruginosa* infections with PER-1 extended-spectrum beta-lactamase in Warsaw, Poland: further evidence for an international clonal complex. *J Clin Microbiol* 2007; 45: 2829–2834.
21. Naas T, Nordmann P, Heidt A. Intercountry transfer of PER-1 extended-spectrum beta-lactamase-producing *Acinetobacter baumannii* from Romania. *Int J Antimicrob Agents* 2007; 29: 226–228.
22. Yamano Y, Nishikawa T, Fujimura T, Yutsudou T, Tsuji M, Miwa H. Occurrence of PER-1 producing clinical isolates of *Pseudomonas aeruginosa* in Japan and their susceptibility to doripenem. *J Antibiot (Tokyo)* 2006; 59: 791–796.
23. Lee S, Park YJ, Kim M, Lee HK, Han K, Kang CS, Kang MW. Prevalence of Ambler class A and D beta-lactamases among clinical isolates of *Pseudomonas aeruginosa* in Korea. *J Antimicrob Chemother* 2005; 56: 122–127.
24. Jeong SH, Bae IK, Kwon SB, Lee K, Yong D, Woo GJ, Lee JH, Jung HI, Jang SJ, Sung KH et al. Investigation of a nosocomial outbreak of *Acinetobacter baumannii* producing PER-1 extended spectrum beta-lactamase in an intensive care unit. *J Hosp Infect* 2005; 59: 242–248.
25. Eraç B, Hoşgör-Limoncu M, Ermertcan Ş, Taşlı H, Aydemir Ş. Prevalence of *bla*_{PER-1} and integrons in ceftazidime-resistant Gram-negative bacteria at a university hospital in Turkey. *Jpn J Infect Dis* 2013; 66: 146–148.
26. Vahaboglu H, Coskuncan F, Tansel O, Ozturk R, Sahin N, Koksall I, Kocazeybek B, Tatman-Otkun M, Leblebicioglu H, Ozinel MA et al. Clinical importance of extended-spectrum β -lactamase (PER-1-type)-producing *Acinetobacter* spp. and *Pseudomonas aeruginosa* strains. *J Med Microb* 2001; 50: 642–645.