

1-1-2022

## Effects of some commercial products on root and crown rot caused by *Phytophthora cactorum* in apple cultivation

SUAT KAYMAK

Follow this and additional works at: <https://journals.tubitak.gov.tr/agriculture>



Part of the [Agriculture Commons](#), and the [Forest Sciences Commons](#)

---

### Recommended Citation

KAYMAK, SUAT (2022) "Effects of some commercial products on root and crown rot caused by *Phytophthora cactorum* in apple cultivation," *Turkish Journal of Agriculture and Forestry*. Vol. 46: No. 1, Article 2. <https://doi.org/10.3906/tar-2106-56>

Available at: <https://journals.tubitak.gov.tr/agriculture/vol46/iss1/2>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Agriculture and Forestry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Effects of some commercial products on root and crown rot caused by *Phytophthora cactorum* in apple cultivation

Suat KAYMAK 

General Directorate of Agricultural Research and Policies, Ankara, Turkey

Received: 17.06.2021 • Accepted/Published Online: 01.11.2021 • Final Version: 09.02.2022

**Abstract:** Root and crown rot caused by *Phytophthora cactorum* (Lebert & John) Schröeter is one of the most common diseases of apple (*Malus domestica*) in the world. Due to the negative influences of fungicides on the environment, using other applicable control measures is a plausible way for the management of the disease. The objective of the study was to evaluate the effects of eleven commercial products (bio-pesticides, fertilizers, and activators) on *P. cactorum* by comparing plant growth parameters (trunk diameter, plant height, number of branches, and root dry weight). Trials were conducted according to a randomized complete block design with three replicates in a greenhouse in 2 consecutive years. In the trials, saplings of Red Chief variety grafted onto MM106 rootstock were planted into pots containing soil mixture inoculated with *P. cactorum*. Two control groups (negative and positive) were also included in each treatment. Tested commercial products generally had significant ( $p < 0.01$ ) positive effects on plant growth parameters (trunk diameter, plant height, and root dry weight) by reducing the severity of the disease (50%–93%). Of the tested commercial products, Alexin 95 PS, Companion, Endo Roots Soluble and Subtilex had the highest efficacies. There were negative correlations between plant growth parameters (trunk diameter, plant height, and root dry weight) and severity of the disease, but root dry weight had significant ( $p < 0.01$ ) correlation ( $r = -0.56$ ) with the severity of the disease. Present findings revealed that some commercial products could reliably be used against *P. cactorum* as an alternative to pesticides in apple cultivation.

**Key words:** Disease control, *Malus domestica*, plant growth, root and crown rot

### 1. Introduction

Apple (*Malus domestica*) is one of the most economically important pome fruits that are widely grown in different ecological regions around the world. In recent years, apple production of the world has reached 86,142,197 tons, and Turkey is the fourth leading producer accounting for 3,625,960 tons of the world's production (FAO, 2020). However, *Phytophthora* species are among the most destructive soil-borne pathogens causing economic losses in orchards in fruit production (Erwin and Ribeiro, 1996; Sánchez et al., 2019). For example, in Bulgaria, it was reported that *Phytophthora* species caused drying the plants up to 10% in some orchards (Nakova, 2010), while they caused considerable losses, 2.5%–24.5% death in nurseries and 0.2%–77.5% in apple orchards in India (Sharma et al., 2014). Among *Phytophthora* species, *Phytophthora cactorum* (Lebert & John) Schröeter is the most frequently reported species in apple-growing regions (Jones and Aldwinckle, 1997; Ebrahimzadeh and Dolar, 2019).

Most of the apple orchards in our country were established with seedlings, and also M9, MM106, and M111 rootstocks were preferred as clone rootstocks (Bayav and Armağan, 2007). According to the findings obtained so far, all of these commercial rootstocks have been determined to be susceptible to *P. cactorum*. An integrated approach including chemical control and cultural practices has been recommended for root and crown rot management in apple orchards (Carisse and Khanizadeh, 2005). In this sense, Maden et al., (1995) reported that *P. cactorum* was the most widespread pathogen causing deaths in apple nurseries in Eğirdir county of Isparta Province, Turkey. Kurbetli and Demirci (2015) also found that *P. cactorum* caused significant yield losses in apple orchards in Turkey.

Root and crown rot symptoms caused by *P. cactorum* include reddish-brown in inner bark and wood tissue. Other symptoms of the pathogen are chlorosis of leaves and reduction of tree vitality and growth. As a result of the infections, infected trees slowly weaken and then collapse in a couple of years (Ellis, 2008; Nakova,

\* Correspondence: suatkaymak@tarimorman.gov.tr

2010). In management of the root and crown rot caused by *Phytophthora* species including *P. cactorum*, some pesticides such as fosetyl-al and phosphorous acid were found to be effective (Utkhede and Smith, 1991; Flett, 1996; Erkilic and Canihs, 1999; Sharma et al., 2014; Kurbetli and Demirci, 2015; Türkölmez and Derviş, 2017). However, widespread use of those and other pesticides cause negative effects on ecosystem and agriculture. Therefore, using resistant rootstocks and applying commercial products that do not have negative influences on ecosystem are the other applicable methods for control of *Phytophthora* species in apple cultivation. Some of the commercial products consist of various active ingredients including biological agents, plant nutrients, and activators. For example, plant growth promoting rhizobacteria (PGPR), plant growth-promoting fungi (PGPF), and arbuscular mycorrhizal fungi (AMF) have numerous benefits to plants such as nutrient intake, protection against soil-borne pathogens, and resistance to environmental stresses (Berendsen et al., 2012; Mendes et al., 2013; Pérez-Jaramillo et al., 2015; Paliwoda and Mikiciuk, 2020). PGPR, PGPF, and AMF can protect plants against pathogenic microorganisms such as bacteria, fungi, viruses, and nematodes and eliminate their harmful effects by activating induced systemic resistance (ISR) in plants (Pieterse et al., 2014; Hossain and Sultana, 2015; Etesami and Maheshwari, 2018).

However, existing commercial products should be tested specifically against pathogens causing serious problems in agriculture. Therefore, developing alternative efficient and environmentally friendly methods to control this disease is urgently needed. Due to its environmentally friendly and nontoxic characteristics, biological control is one of the most desired methods for managing root and crown rot disease.

In this study, a potential biocontrol agent was tested for antifungal activity and effective plant growth-promoting traits. The aims of this study were (i) to determine the effects of eleven commercial products (biopesticides, fertilizers, and activators) on severity of *P. cactorum* in vivo and (ii) to examine the effects of commercial products on plant growth parameters (trunk diameter, plant height, number of branches and root dry weight) of apple saplings.

**2. Materials and methods**

**2.1. Commercial products used in the trials**

Information about eleven commercial products used in the present trials is provided in Table 1.

**2.2. Inoculum preparation**

A virulent strain of *P. cactorum* [accession number: HM357616 in Genbank (<https://www.ncbi.nlm.nih.gov>)] was obtained from Plant Protection Central Research

**Table 1.** Trademark, ingredients, recommended application doses, and manufacturers of commercial products used in present trials.

Trademark	Active ingredient/microorganism	Recommended application doses	Company
Companion	<i>Bacillus subtilis</i> GB03	0.5 mL/m <sup>2</sup>	Growth products
Symbion Vam	<i>Glomus fasciculatum</i>	1 g/m <sup>2</sup>	Stanes
Green Miracle	Vegetable oil acid	0.2 mL/m <sup>2</sup>	Stanes
Cropset	<i>Lactobacillus acidophilus</i> , plant extract, MnSO <sub>4</sub> , FeSO <sub>4</sub> , CuSO <sub>4</sub>	0.06 mL/m <sup>2</sup>	Ant tarım (Improcrop EU)
Isr-2000	<i>Lactobacillus acidophilus</i> , plant extract, yeast extract, benzoic acid	0.1 mL/m <sup>2</sup>	Ant tarım (Improcrop EU)
Actinovate	<i>Streptomyces lydicus</i>	0.4 mL/m <sup>2</sup>	Mts agro
Subtilex	<i>Bacillus subtilis</i> MBI 600	0.5 mL/m <sup>2</sup>	Bioglobal
Tricho plus	<i>Trichoderma harzihanum</i>	0.04 g/m <sup>2</sup>	Bioglobal
Alexin 95 PS	Phosphorus pentoxide (P <sub>2</sub> O <sub>5</sub> ) 52%, Potassium oxide (K <sub>2</sub> O) 42%	0.7 g/m <sup>2</sup>	Sumitoma
Combat Plus	Plant activators	0.4 mL/m <sup>2</sup>	Bioglobal
Endo Roots Soluble	<i>Glomus intraradices</i> .....21 <i>Glomus aggregatum</i> .....20 <i>Glomus mosseage</i> .....20 <i>Glomus clarum</i> .....1 <i>Glomus monosporus</i> .....1 <i>Glomus deserticola</i> .....1 <i>Glomus brasilianum</i> .....1 <i>Glomus etunicatum</i> .....1 <i>Gigaspora margarita</i> .....1	0.5 mL/m <sup>2</sup>	Bioglobal

Institute, Ankara, Turkey. Five agar plugs (5 mm) from 6 day-old colonies of *P. cactorum* were added to jars (1 L) containing a mixture of volumes [carrot juice (400 mL) + vermiculite (560 mL) + oat seeds (40 ml)] and incubated at 23 °C for 21 days.

### 2.3. Greenhouse trials

The inoculum mixture in each jar was transferred to a pot (25-L) containing soil mixture [orchard soil and sand (1:1)] and homogeneously mixed into the pot. The inoculum constituted 4% of the total volume of each pot (Latorre et al., 2001). Two-month-old saplings of Red Chief variety grafted onto MM106 rootstock were used as test plants. Inoculations were carried out at the Fruit Research Institute (Eğirdir, Isparta Province, Turkey). One sapling of the variety was planted per pot and commercial products were applied at the recommended doses to the pots. In positive controls, the inoculum was applied to the pots without the commercial products. However, in negative controls, sterile water was only used. The trials were conducted according to a randomized complete block design with 3 replicates in two consecutive years (Figure 1).

The saplings were grown in the pots for 4 months in the greenhouse in June–September. In order to promote growth of *P. cactorum*, the pots were kept wet by drip irrigation.

### 2.4. Evaluation of the trials

#### 2.4.1. Plant growth parameters

Trunk diameter was measured from 10-cm above the grafting point with the use a caliper, while sapling height was measured from the soil level to the top with a meter stick. Number of branches per sapling was recorded by counting branches that are longer than 20 cm.

To determine root dry weights of the saplings, immediately after removing plants, the roots were washed under running tap water and consequently separated from the soil particles. They were kept at 25 °C for a few hours and then cut at the beginning of the root collars. The roots of each sapling were dried at 70 °C for 4 days and then weighed with the use of a digital scale.

#### 2.4.2. Disease severity and efficacy of the commercial products

Considering percentage of stems with lesion, disease severity was determined using 0–4 scale, where: 0 = no visible, 1, 2, 3 and 4 = 1%–25%, 26%–50%, 51%–75%, and 76%–100% with lesions, respectively (Tidball and Linderman, 1990) and the formula below.

$$\text{Disease severity (\%)} = \frac{\sum i (n_i \times v_i)}{N \times V}$$

where  $v_i$  is the scale class,  $n_i$  is the number in one class,  $N$  is the total number,  $V$  is the highest class,  $i$  is the number of classes.



**Figure 1.** Setting up the trials with saplings of Red Chief variety grafted onto MM106 rootstock.

To quantify disease severity values, percentage efficacy of each commercial product was determined according to the Townsend-Heuberger formula below (Karman, 1971).

$$\% \text{ efficacy} = [(I_c - I_t) / I_c] \times 100$$

where  $I_c$  is the disease severity of untreated control,  $I_t$  is the disease severity of the treatment.

#### 2.4.3. Data analysis

Data of the study were analyzed using SPSS 16 (IBM Corp., Armonk) software package. Means of each growth parameter, disease severity, and efficacy of each commercial product were categorized according to Duncan's multiple range test ( $p < 0.01$ ).

### 3. Results

According to average of 2 years, differences in growth parameters (trunk diameter, plant height, and root dry weight) of the trails were found to be significant ( $p < 0.01$ ). The highest mean trunk diameter (19.79 mm) was found in Subtilex trails, while the lowest one (16.78 mm) was determined in pathogen inoculated control. Mean trunk diameter of noninoculated control group was 17.31 mm. In addition to Subtilex, as compared to the pathogen inoculated control, mean trunk diameters, of Alexin 95 PS, Isr-2000, and Symbion Vam (18.92, 18.81, and 18.46 mm, respectively) were also found to be significant ( $p < 0.01$ ).

Mean sapling heights of the noninoculated and pathogen inoculated control groups were 151.78 and

145.22 cm, respectively. However, mean sapling heights in experimental trails ranged from 138.89 to 178.06 cm. As compared to the pathogen inoculated control group, mean sapling heights of Alexin 95 PS, Isr-2000, Actinovate, Green Miracle, Subtilex, and Cropset trails (181.11, 178.06, 175.15, 172.67, 172.17, and 171.39 cm, respectively) were found to be significant ( $p < 0.01$ ).

Mean number of branches of the pathogen inoculated control was 2.11, while it was 3.18 in the noninoculated control. The difference was significant ( $p < 0.01$ ), confirming destructiveness of the pathogen in the apple saplings. On the other hand, mean number of branches of the experimental trails were not significantly different from the pathogen inoculated control group. One example for evaluations of the results of the study is presented in Figure 2.

Mean root dry weights of the noninoculated and pathogen inoculated controls were 110.20 and 98.80 g, respectively. However, as compared to the pathogen inoculated controls, mean root dry weights of Alexin 95 PS, Subtilex, and Tricho plus trials (126.86, 125.24, and 121.65 g, respectively) were significantly different ( $p < 0.01$ ).

Mean disease severity of the pathogen inoculated control was 43.75%, while disease severity values of experimental trails ranged from 3.125 to 21.875%. The highest efficacy (93.75%) was detected in Alexin 95 PS trials (Table 2).



**Figure 2.** Evaluation of negative control (left), treatment (middle), and positive control (right) in companion (commercial product) application to the Red Chief variety grafted onto MM106 rootstock in the trials.

**Table 2.** Effects of the commercial products on plant growth parameters and disease severity in apple saplings of Red Chief variety grafted onto MM106 rootstock and efficacy rates of the commercial products.

Applications	Trunk diameter (mm)*	Sapling height (cm)*	Number of branches*	Root dry weight (g)	Disease severity (%)*	Efficacy (%)
Companion	17.00 ± 0.55 de (13.51–21.27)	153.61 ± 7.38 bcd (105.00–197.00)	1.69 ± 0.18 b (1.00–3.00)	112.34 ± 6.92 abc (83.98–152.05)	6.25 bc	87.5 ab
Symbion Vam	18.46 ± 0.42 abcd (16.64–23.21)	161.65 ± 5.26 abc (126.00–202.00)	1.72 ± 0.31 b (0.00–5.00)	120.58 ± 8.22 abc (77.00–159.70)	12.5 bc	62.5 abc
Green Miracle	17.87 ± 0.34 bcde (15.95–20.53)	172.67 ± 6.92 ab (110.00–215.00)	1.50 ± 0.20 b (1.00–4.00)	110.19 ± 5.49 abc (85.85–134.23)	15.62 bc	58.33 abc
Cropset	17.29 ± 0.40 cde (13.15–19.67)	171.39 ± 6.21 ab (129.00–221.00)	2.06 ± 0.29 b (1.00–5.00)	99.74 ± 7.68 bc (58.88–126.96)	21.875 b	50 c
Isr-2000	18.81 ± 0.55 abc (15.06–24.23)	178.06 ± 5.00 a (139.00–212.00)	1.89 ± 0.25 b (1.00–4.00)	113.76 ± 7.43 abc (69.05–141.09)	12.5 bc	62.5 abc
Actinovate	17.89 ± 0.46 bcde (14.96–20.15)	175.15 ± 5.03 a (151.00–215.00)	2.24 ± 0.22 b (1.00–4.00)	121.24 ± 8.82 abc (70.68–168.64)	12.5 bc	62.5 abc
Subtalex	19.79 ± 0.43 a (16.79–22.94)	172.17 ± 6.24 ab (113.00–211.00)	1.82 ± 0.20 b (1.00–3.00)	125.16 ± 6.16 a (93.00–151.66)	9.375 bc	75 abc
Tricho Plus	17.61 ± 0.53 bcde (12.79–20.78)	138.89 ± 6.70 d (90.00–193.00)	1.86 ± 0.23 b (1.00–4.00)	121.65 ± 6.82 ab (97.40–161.51)	18.75 bc	56.25 bc
Alexin 95 PS	18.92 ± 0.54 ab (14.21–24.74)	181.11 ± 7.57 a (122.00–238.00)	1.88 ± 0.21 b (1.00–4.00)	126.86 ± 5.68 a (95.50–147.65)	3.125 bc	93.75 a
Combat Plus	17.37 ± 0.33 cde (15.05–20.12)	163.67 ± 7.27 abc (85.00–204.00)	2.22 ± 0.33 b (1.00–6.00)	98.86 ± 7.04 c (72.25–133.68)	21.87 b	50 c
Endo Roots Soluble	17.85 ± 0.53 bcde (11.30–22.14)	164.83 ± 6.51 abc (112.00–210.00)	1.67 ± 0.14 b (1.00–3.00)	107.82 ± 6.41 abc (66.33–139.38)	6.25 bc	87.5 ab
C0	17.31 ± 0.34 cde (14.45–19.76)	151.78 ± 5.91 bcd (118.00–205.00)	3.18 ± 0.43 a (1.00–7.00)	110.20 ± 4.94 abc (93.55–140.85)	0 c	-
C1	16.78 ± 0.54 e (13.05–20.63)	145.22 ± 5.78 cd (113.00–193.00)	2.11 ± 0.24 b (1.00–4.00)	98.80 ± 5.52 c (68.35–119.66)	43.75 a	-
	F = 3.493, p = 0.000	F = 4.194, p = 0.000	F = 2.589, p = 0.003	F = 2.09, p = 0.020	F = 3.862, p = 0.002	

\* Mean values of 2 years, the same letters are not significantly different according to Duncan's multiple range test ( $p < 0.01$ ). Mean ± SE (min–max.), C0: Negative control, C1: Positive control

There was a negative correlation ( $r = -0.43$ ) between trunk diameter and disease severity. Similarly, plant height negatively correlated ( $r = -0.35$ ) with disease severity, but the correlation between root dry weight and disease severity ( $r = -0.56$ ) was significant ( $p < 0.01$ ).

#### 4. Discussion

Antifungal and plant growth-promoting effects of various strains of *Bacillus subtilis* were revealed in previous studies. For example, application of *B. subtilis* to tomato seedlings not only enhanced plant growth but also suppressed soil-borne pathogens including *Pythium aphanidermatum* (Kipngeno et al., 2015; Qiao et al., 2017). Plant growth bacteria excrete volatile compounds were found to be effective against soil-borne oomycetes including *Phytophthora cinnamomi* (Méndez-Bravo et al., 2018). In addition to volatile compounds, *B. subtilis* has functional

metabolites such as IturinA that causes damages to cell structure of *Phytophthora infestans* by inducing disruptions in cell membrane and organelle formation. For these reasons, *Bacillus subtilis* is capable of inhibiting mycelium growth of *P. infestans* (Wang et al., 2019). In a greenhouse study, *B. subtilis* decreased mortality of pistachio seedlings up to 80% against *Phytophthora pistaciae* (Moradi et al., 2018). Likewise, in the present study, as compared to positive control group, applications of Subtalex (*Bacillus subtilis* MBI 600) caused significant increases in trunk diameter, plant height, and root dry weight of the saplings by reducing severity of *Phytophthora cactorum*, indicating antifungal and plant growth-promoting effects of the commercial product.

As related to the other contents of the commercial products used in the present study, Symbion Vam (*Glomus fasciculatum*) and Endo Roots Soluble (*Glomus* spp.)

had other bio-pesticides with their *Glomus* contents. Applications of Symbion Vam (*Glomus fasciculatum*) caused significant increases in trunk diameter and root dry weight by decreasing severity of the disease, while effects of Endo Roots Soluble (*Glomus* spp.) on the plant growth parameters were not significant but its application caused significant reductions in severity of the disease. It is known that arbuscular mycorrhizal fungi (AMF) are potentially protective agents with their ability of forming symbiotic associations with root systems of various crops. For example, in a study, arbuscular mycorrhizal fungus *Glomus mosseae* decreased infections of *Phytophthora parasitica* in tomato by providing a systemic resistance for host plant through excreting hydrolytic enzymes (chitinase, chitosanase,  $\beta$ -1,3-glucanase and superoxide dismutase) (Poza et al., 2002). In another study, *Glomus* application increased plant height, pseudo-stem diameter and root weight of banana seedlings by suppressing *Fusarium oxysporum* f. sp. *cubense* (FocR4) (Castillo et al., 2019). Likewise, four commercial products (Bacto\_Prof, Endomyk\_Basic, Endomyk\_Conc and Endomyk\_Prof mycorrhizal) with mycorrhizal content decreased *Fusarium oxysporum* infections in tomato plants by significantly enhancing plant height (Al-Hmoud and Al-Momany, 2015). With regard to *Phytophthora*, applications of mycorrhizal fungi including *Glomus intraradices* increased plant height, stem diameter, number of leaves, and fresh aerial biomass and root growth of pepper and strawberry by suppressing *Phytophthora capsici* and *Phytophthora cactorum*, respectively (Hautsalo et al., 2016; Tena et al., 2016).

Apart from arbuscular mycorrhizal fungi, *Trichoderma* species in particular *Trichoderma harzianum* is closely associated with plant roots. In the present study, Tricho plus (*Trichoderma harzianum*) caused significant increases in root dry weight by reducing severity of *Phytophthora cactorum*. Similarly, in a study, *Phythisium ultimum*, *Phytophthora capsici* and *Phytophthora infestans* decreased fresh and dry shoot weight of tomato, while *Trichoderma harzianum* improved entire plant growth including root weight in treatment plots, indicating efficacy of application of *T. harzianum* against the soil-borne pathogens. The mechanism was related to inhibition of entrance of both pathogens to vascular bundle of tomato tissues, which provided reduction in size of infection areas of the pathogens (Uddin et al., 2018; Kariuki et al., 2020). In a 3-year-field experiment, *Trichoderma* spp. decreased soil populations of *Phytophthora cactorum* by reducing disease incidence up to 76.6% in strawberry cultivation (Porras et al., 2007). In another 2-year-field study, *Trichoderma harzianum* reduced severity of *Phytophthora capsici* up to 42% in pepper cultivation (Timila and Manandhar, 2020). All these supported the results of those commercial products tested in the present study.

In addition, of the tested commercial products, some of which had nutrient contents. In fact, plant nutrients may affect tolerance/resistance of plants to pathogens (Gupta et al., 2017). Fertilizers have significant physiological effects on plant growth (Li et al., 2012). For example, potato yields and infection caused by *Phytophthora infestans* are associated with supply of potassium. Application of potassium provided a protection for potato plants against *P. infestans* by enhancing phytoalexins and phenols in tubers, which stimulates production of biochemical compounds, pathogenesis-related enzymes and antioxidant enzyme activities in the host (Kowalska and Drożdżyński, 2018; Mohammadi et al., 2019). In a field study, applications of potassium phosphonate reduced incidence of root rot of papaya caused by *Phytophthora palmivora* by 47% (Vawdrey et al., 2004). Commercial product Phytogard that contains potassium phosphite led to preventive control of *Phytophthora plurivora* in beech plants (Rezende et al., 2020). Likewise, in the present study, of the tested commercial products, Alexin 95 PS (Phosphorus pentoxide + Potassium oxide), considered a fertilizer with its content, caused significant increases in trunk diameter, plant height, and root dry weight by reducing severity of the disease. Moreover, of the tested commercial products, the highest efficacy was detected in this commercial product. In addition to nutrients, plant extracts may play a role in plant resistance. Demirci and Dolar (2006) reported that extracts of dry cabbage, garlic, and alfalfa were effective in decreasing severity of *Phytophthora capsici* in vitro and in vivo. Similarly, in the present study, Isr-2000 (*Lactobacillus acidophilus* + plant extract + yeast extract + benzoic acid) led to significant increases in trunk diameter and plant height by reducing severity of *Phytophthora cactorum*, while other commercial product Cropset (*Lactobacillus acidophilus* + plant extract + MnSO<sub>4</sub> + FeSO<sub>4</sub> + CuSO<sub>4</sub>) increased plant height by reducing severity of *P. cactorum*.

Plant activators are capable of improving plant health through antifungal, antibacterial, oomycetocidal and act as elicitors of plant immunity through the routes of salicylic acid (SA) and jasmonic acid (JA) (Pye et al., 2013; Bitas et al., 2013; Schalchli et al., 2016). In this regard, some plant activators like HarpinEa and Salicylic acid might reduce infections of *Phytophthora infestans* and *Rhizoctonia solani* in tomato and chili by enhancing specific enzyme activities that provide systemic acquired resistance (SAR) in the host (Tosun et al., 2003; Baloch et al., 2018). For example, another plant activator, called pyrimidin-type plant activator (PPA), increased fresh weight of rice and Arabidopsis plants. In addition, PPA also promoted lateral root development and plant defense against pathogen invasion (Sun et al., 2015). In the present study, Combat Plus (plant activator) significantly reduced the severity of *P. cactorum*, which is in agreement with the results of the aforementioned previous studies.

As compared to positive and negative controls, sapling height significantly increased in Actinovate (*Streptomyces lydicus*) applied plots, indicating suppression of *Phytophthora cactorum*. In this regard, Ezziyyani et al. (2007) reported that *Streptomyces rochei* had a specific activity against *P. capsici*. Likewise, in another study, *Streptomyces* spp. was found to be effective in controlling root rot of alfalfa (*P. medicaginis*) and soybean (*P. sojae*) (Xiao et al., 2002). Besides, *Streptomyces* species were effective against *Rhizoctonia solani*, *Pythium aphanidermatum*, and *P. ultimum* in various crops (Hamdali et al., 2008; El-Tarabily et al., 2009; Goudjal et al., 2014). Suppression of plant pathogens by *Streptomyces* species is related to antibiotic production, volatile components, hydrolytic enzymes, and siderophore production (Palaniyandi et al., 2013; Wang et al., 2013; Sadeghi et al., 2017).

## 5. Conclusions

Pesticides used for disease control may generate serious consequences for environment and food safety, so there is a need for alternative management methods (Gupta et al., 2017). Therefore, alternative methods should be applicable and environment-friendly and they should be presented to farmers for management of plant diseases in agriculture. As a result of the present study, of the tested commercial

products, Alexin 95 PS (fertilizer), Companion (*Bacillus subtilis* GB03), Endo Roots Soluble (*Glomus* spp.), and Subtilix (*Bacillus subtilis* MBI 600) had the highest efficacies against *Phytophthora cactorum*. Applications of these products to newly planted saplings can protect them from soil-borne pathogens like *Phytophthora cactorum* in apple cultivation.

In this study, a potential biocontrol agent was tested for antifungal activity and effective plant growth-promoting traits. The conclusions achieved in this work help to understand the current state of the research field, and reveal new insights on the development of efficient biocontrol strategies for the control of *P. cactorum*. Furthermore, the present study also suggests that existing commercial products should be tested specifically against plant pathogens causing serious problems in agricultural production.

## Acknowledgments

This research was partially funded by the CORE Organic II Funding Bodies, being partners of the FP7 ERA-NET through BIO-INCROP Project (Project no. 249667), and partially by General Directorate of Agricultural Research and Policies, Ministry of Agriculture and Forestry, Republic of Turkey.

## References

- Al-Hmoud G, Al-Momany A (2015). Effect of four mycorrhizal products on fusarium root rot on different vegetable crops. *Plant Pathology & Microbiology* 6: 255. doi: 10.4172/2157-7471.1000255
- Baloch MS, Rajput NA, Atiq M, Rehman A, Khan SM et al. (2018). Effect of different plant activators against *Rhizoctonia solani* causing root rot of chilli. *Pakistan Journal of Phytopathology* 30 (01): 45-50. doi: 10.33866/phytopathol.030.01.0441
- Bayav A, Armağan G (2007). Adoption Levels of The Innovations and Research Findings, and Impact Assessments in The Apple Growing Enterprises in Isparta Province, Türkiye VIII. Tarım Ekonomisi Kongresi, Gıda Politikası Cildi p.168-181.
- Berendsen RL, Pieterse CMJ, Bakker P (2012). The rhizosphere microbiome and plant health. *Trends in Plant Science* 17 (8): 478-86. doi: 10.1016/j.tplants.2012.04.001
- Bitas V, Kim HS, Bennett JW, Kang S (2013). Sniffing on microbes: diverse roles of microbial volatile organic compounds in plant health. *Molecular Plant-Microbe Interactions*. 26: 835-843. doi: 10.1094/MPMI-10-12-0249-CR
- Carisse O, Khanizadeh S (2005). Relative resistance of newly released apple rootstocks to *Phytophthora cactorum*. *Canadian Journal of Plant sciences* 86: 199-204. doi: 10.4141/P04-176
- Castillo AG, Puig CG, Cumagun C (2019). Non-synergistic effect of *Trichoderma harzianum* and *Glomus* spp. in reducing infection of fusarium wilt in banana. *Pathogens* (Basel, Switzerland) 8 (2): 43. doi: 10.3390/pathogens8020043
- Demirci F, Dolar FS (2006). Effects of some plant materials on phytophthora blight (*Phytophthora capsici* Leon.) of pepper. *Turkish Journal of Agriculture Forestry* 30 (4): 247-252.
- Ebrahimzadeh R, Dolar FS (2019). Diseases of apples caused by *Phytophthora* spp., symptoms and descriptions. *Black Sea Journal of Agriculture* 2 (2): 109-118.
- Ellis MA (2008). *Phytophthora* root and crown rot of fruit trees. Ohio State University Extension. [Available: <https://ohioline.osu.edu/factsheet/plpath-fru-06>] [Accessed: 15 April 2021].
- El-Tarabily KA, Nassar AH, Hardy GE, Sivasithamparam K (2009). Plant growth promotion and biological control of *Pythium aphanidermatum*, a pathogen of cucumber, by endophytic actinomycetes. *Journal of Applied Microbiology* 106: 13-26. doi: 10.1111/j.1365-2672.2008.03926.x
- Erkılıç A, Canhoş Y (1999). Determination of the effect of fosetyl-al against citrus gummosis disease caused by *Phytophthora citrophthora* (Smith and Smith) leonien. *Turkish Journal of Agriculture Forestry* 23: 419-424.



- Erwin DC, Ribeiro OK (2005). *Phytophthora* diseases worldwide. APS Press, 2nd Ed.
- Etesami H, Maheshwari DK (2018). Use of plant growth promoting rhizobacteria (PGPRs) with multiple plant growth promoting traits in stress agriculture: Action mechanisms and future prospects. *Ecotoxicology and Environmental Safety* 156: 225-246.
- Ezziyiani M, Requena ME, Egea-Gilbert C, Candela ME (2007). Biological Control of *Phytophthora* Root Rot of Pepper Using *Trichoderma harzianum* and *Streptomyces rochei* in Combination. *Journal of Phytopathology* 155: 342-349. doi: 10.1111/j.1439-0434.2007.01237.x
- Food and Agriculture Organization of the United Nations (2020). FAOSTAT [online]. Website <http://www.fao.org/faostat/en/> [accessed 02 April 2021].
- Flett S (1996). Developing safe foliar spraying of phosphonic acid to control peach and apple *Phytophthora*. Project AP253, Final Report. Horticultural Research and Development Corporation, New South Wales.
- Goudjal Y, Toumatia O, Yekkour A, Sabaou N, Mathieu F et al. (2014). Biocontrol of *Rhizoctonia solani* damping-off and promotion of tomato plant growth by endophytic actinomycetes isolated from native plants of Algerian Sahara. *Microbiological Research* 169: 59–65. doi: 10.1016/j.micres.2013.06.014
- Gupta N, Debnath S, Sharma S, Sharma P, Purohit J (2017). Role of nutrients in controlling plant diseases in sustainable agriculture. A review. In book: *Agriculturally Important Microbes for Sustainable Agriculture* (pp.217-262). doi: 10.1007/978-981-10-5343-6-8
- Hamdali H, Hafidi M, Virolle MJ, Ouhdouch Y (2008). Growth promotion and protection against damping-off of wheat by two rock phosphate solubilizing actinomycetes in a P-deficient soil under greenhouse conditions. *Applied Soil Ecology* 40: 510–517. doi: 10.1016/j.apsoil.2008.08.001
- Hautsalo J, Vestberg M, Parikka P, Kukkonen S, Karhu S et al. (2016). Biological control of strawberry crown rot is substrate dependent phenomenon. *Journal of Berry Research* 6: 65–79. doi: 10.3233/JBR-150110
- Hossain MM, Sultana F (2015). Genetic variation for induced and basal resistance against leaf pathogen *Pseudomonas syringae* pv. *tomato* DC3000 among *Arabidopsis thaliana* accessions. *Springerplus* 4: 296. doi: 10.1186/s40064-015-1070-z
- Jones AL, Aldwinckle HS (1997). *Compendium of Apple and Pear diseases*. APS perss. Third Ed.
- Karman M (1971). *General Information on Plant Protection Research. Establishment of Trials and Evaluation Principles*. Bölge Zirai Mücadele Araştırma Enstitüsü Mesleki Kitaplar Serisi. İzmir-Bornova.
- Kariuki WG, Mungai NW, Otaye DO, Thuita M, Muema E et al. (2020). Antagonistic Effects of Biocontrol Agents Against *Phytophthora infestans* and growth Stimulation in tomatoes. *African Crop Science Journal* 28: 55–70. doi: 10.4314/acsj.v28i1.5S
- Kipngeno P, Losenge T, Maina N, Kahangi E, Juma P (2015). Efficacy of *Bacillus subtilis* and *Trichoderma asperellum* against *Pythium aphanidermatum* in tomatoes. *Biological Control* 90: 92-95. doi: 10.1016/j.biocontrol.2015.05.017
- Kowalska J, Drożdżyński D (2018). Effects of potassium fertilisation on potato late blight and yield. Short Communication. *Plant Protection Science* 54: 87–91. doi: 10.17221/79/2017-PPS
- Kurbetli İ, Demirci F (2015). Effects of some fungicides against *Phytophthora cactorum* (Lebert & John) Schröeter causing root and crown rot of apple trees. *Plant Protection Bulletin* 55 (1): 31-40.
- Latorre BA, Rioja ME, Wilcox WF (2001). *Phytophthora* species associated with crown and root rot of apple in Chile. *Plant Disease* 85: 603-606.
- Li X, Wei J, Li Y (2012). Physiological Effects of Nitrogen, Phosphorus and Potassium on Blueberry 'Bluecrop'. *Acta Horticulture* 926: 347-351 doi: 10.17660/ActaHortic.2012.926.48
- Maden S, Erzurum K, Yürüt A, Güreş M, Erkal Ü (1995). Studies on the determination of the causes of drying seen in apple rootstocks M.M.106, VII. Turkey Phytopathology Congress Proceedings, 199–203.
- Mendes R, Garbeva P, Raaijmakers JM (2013). The rhizosphere microbiome: significance of plant beneficial, plant pathogenic and human pathogenic microorganisms. *FEMS Microbiology Reviews* 37: 634-663. doi: 10.1111/1574-6976.12028
- Méndez-Bravo A, Cortazar-Murillo EM, Guevara-Avendaño E, Ceballos-Luna O, Rodríguez-Haas B et al. (2018). Plant growth-promoting rhizobacteria associated with avocado display antagonistic activity against *Phytophthora cinnamomi* through volatile emissions. *PLoS ONE* 13 (3): e0194665. doi: 10.1371/journal.pone.0194665
- Mohammadi MA, Zhang Z, Xi Y, Han H, Lan F et al. (2019). Effects of potassium phosphite on biochemical contents and enzymatic activities of Chinese potatoes inoculated by *Phytophthora infestans*. *Applied Ecology and Environmental Research* 17 (2): 4499-4514. doi: 10.15666/aeer/1702\_44994514
- Moradi M, Nejad FJ, Bonjar GHS (2018). Efficacy of *Bacillus subtilis* native strains for biocontrol of *Phytophthora* crown and root rot of pistachio in Iran. *Tropical Plant Pathology* 43: 306–313. doi: 10.1007/s40858-018-0226-0
- Nakova M (2010). Monitoring of *Phytophthora* species on fruit trees in Bulgaria. *European Journal of Plant Pathology* 128: 517-525.
- Palaniyandi SA, Yang SH, Suh JW (2013). Extracellular proteases from *Streptomyces phaenopurpureus* ExPro138 inhibit spore adhesion, germination and appressorium formation in *Colletotrichum coccodes*. *Journal Applied Microbiology* 115: 207–217. doi: 10.1111/jam.12212
- Paliwoda D, Mikiciuk G (2020). Use of rhizosphere microorganisms in plant production—A Review Study. *Ecological Engineering* 21 (8): 292-310. doi: 10.12911/22998993/126597
- Pérez-Jaramillo JE, Mendes R, Raaijmakers JM (2015). Impact of plant domestication on rhizosphere microbiome assembly and functions. *Plant Molecular Biology* 90: 635-644. doi: 10.1007/s11103-015-0337-7

- Pieterse CMJ, Zamioudis C, Berendsen RL, Weller DM, Van Wees SCM et al. (2014). Induced systemic resistance by beneficial microbes. *Annual Review Phytopathology* 52: 347-375. doi: 10.1146/annurev-phyto-082712-102340
- Porrás M, Barrau C, Arroyo FT, Santos B, Blanco C et al. (2007). Reduction of *Phytophthora cactorum* in strawberry fields by *Trichoderma* spp. and soil solarization. *Plant Disease* 91: 142-146.
- Pozo MJ, Cordier C, Dumas-Gaudot E, Gianinazzi S, Barea JM (2002). Concepción Azcón-Aguilar, Localized versus systemic effect of arbuscular mycorrhizal fungi on defence responses to *Phytophthora* infection in tomato plants. *Journal Experimental Botany* 53 (368): 525-534. doi: 10.1093/jexbot/53.368.525
- Pye MF, Hakuno F, MacDonald JD, Bostock RM (2013). Induced resistance in tomato by SAR activators during predisposing salinity stress. *Frontiers in Plant Science* 4: 116. doi: 10.3389/fpls.2013.00116
- Qiao J, Yu X, Liang X (2017). Addition of plant-growth-promoting *Bacillus subtilis* PTS-394 on tomato rhizosphere has no durable impact on composition of root microbiome. *BMC Microbiology* 17: 131. doi: 10.1186/s12866-017-1039-x
- Rezende DC, Brandão DFR, Brand SC, Blumer S, Pascholati SF et al. (2020). Potassium phosphite and its implications for the control of *Phytophthora plurivora* in beech. *Research, Society and Development [S.l.]* 9 (10): e5629108824. doi: 10.33448/rsd-v9i10.8824
- Sadeghi A, Koobaz P, Azimi H, Karimi E, Akbari AR (2017). Plant growth promotion and suppression of *Phytophthora drechsleri* damping-off in cucumber by cellulase-producing *Streptomyces*. *Biological Control* 62 (6): 805-819. doi: 10.1007/s10526-017-9838-4
- Sánchez AD, Ousset MJ, Sosa MC (2019). Biological control of *Phytophthora* collar rot of pear using regional *Trichoderma* strains with multiple mechanisms. *Biological Control* 135: 124-134. doi: 10.1016/j.biocontrol.2019.05.012
- Schalchli H, Tortella GR, Rubilar O, Parra L, Hormazabal E et al. (2016). Fungal volatiles: an environmentally friendly tool to control pathogenic microorganisms in plants. *Crit. Rev. Nature Biotechnology* 36: 144-152. doi: 10.3109/07388551.2014.946466
- Sharma IM, Negi HS, Sharma S (2014). Integrated management of collar rot in apple caused by *Phytophthora cactorum*. *Indian Phytopathology* 67 (2): 168-173.
- Sun T-J, Lu Y, Narusaka M, Shi C, Yang YB et al. (2015). A Novel Pyrimidin-Like Plant Activator Stimulates Plant Disease Resistance and Promotes Growth. *PLoS ONE* 10 (4): e0123227. doi: 10.1371/journal.pone.0123227
- Tidball CJ, Linderman RG (1990). *Phytophthora* root and stem rot of apple rootstocks from stool beds. *Plant Disease* 74 (2): 141-146.
- Tena AR, Aguilar EEQ, Enríquez GR, Pérez LL (2016). Mycorrhizae in *Capsicum annuum* L. to promote growth and biosecurity against *Phytophthora capsici* L. *Revista Mexicana de Ciencias Agrícolas* 7 (4): 857-870.
- Timila RD, Manandhar S (2020). Biocontrol efficacy of *Trichoderma* spp. against phytophthora blight of pepper. *Nepalese Horticulture* 14: 15-20. doi: 10.3126/nh.v14i1.30600
- Tosun N, Karabay NU, Turkusay H, Aki C, Turkan I et al. (2003). The effect of harpinea as plant activator in control of bacterial and fungal diseases of tomato. *Acta Hort* 613: 251-254. doi: 10.17660/ActaHortic.2003.613.39
- Türkölmez S, Derviş S (2017). Activity of metalaxyl-m+mancozeb, fosetyl-al, and phosphorous acid against *Phytophthora* crown and root rot of apricot and cherry caused by *Phytophthora palmivora*. *Plant Protection Science* 53: 216-225. doi: 10.17221/38/2016-PPS
- Uddin MN, Ur Rahman U, Khan W, Uddin N, Muhammad M (2018). Effect of *Trichoderma harzianum* on tomato plant growth and its antagonistic activity against *Phytophthora ultimum* and *Phytophthora capsici*. *Egyptian Journal of Biological Pest Control* 28: 32. doi: 10.1186/s41938-018-0032-5
- Utkhede RS, Smith EM (1991). Effects of fosetyl-al, and *Enterobacter aerogenes* on crown and root rot of apple trees caused by *Phytophthora cactorum* in British Columbia. *Plant Diseases* 75: 406-409.
- Vawdrey LL, Grice KE, Peterson RA, De Faveri J (2004). The use of metalaxyl and potassium phosphonate, mounds, and organic and plastic mulches, for the management of *Phytophthora* root rot of papaya in far northern Queensland. *Australas. Plant Pathology* 33: 103-107.
- Wang C, Wang Z, Qiao X, Li Z, Li F et al. (2013). Antifungal activity of volatile organic compounds from *Streptomyces alboflavus* TD-1. *FEMS Microbiology Letters* 341: 45-51. doi: 10.1111/1574-6968.12088
- Wang Y, Zhang C, Wu L, Wang Le, Gao W et al. (2019). Inhibitory effect of *Bacillus subtilis* WL-2 and its IturinA lipopeptides against *Phytophthora infestans*. *BioRxiv* 751131. doi: 10.1101/751131
- Xiao K, Kinkel LL, Samac DA (2002). Biological control of *Phytophthora* root rots on Alfalfa and Soybean with *Streptomyces*. *Biological Control* 23: 285-295. doi: 10.1006/bcon.2001.1015
- Zheng HZ, Kim YW, Lee HJ, Park RD, Jung WJ et al. (2004). Quantitative changes of PR proteins and antioxidative enzymes in response to *Glomus intraradices* and *Phytophthora capsici* in pepper (*Capsicum annuum* L.) plants. *Journal of Microbiology and Biotechnology* 14 (3): 553-562.