

1-1-1998

Direct Sums and the Schur Property

BETÜL TANBAY

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

TANBAY, BETÜL (1998) "Direct Sums and the Schur Property," *Turkish Journal of Mathematics*: Vol. 22: No. 3, Article 9. Available at: <https://journals.tubitak.gov.tr/math/vol22/iss3/9>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

DIRECT SUMS AND THE SCHUR PROPERTY

B. Tanbay

Abstract

It is a known fact that ℓ^1 , the dual space of the null sequences c_0 , has the Schur property, that is, weakly convergent sequences in ℓ^1 are norm convergent. In this paper, we prove that if $(X_\alpha)_{\alpha \in I}$ are Banach spaces and $X = (\oplus_{\alpha \in I} X_\alpha)_1$ their l_1 -sum, then the space X has the Schur property iff each factor X_α has it.

Key words and phrases: Schur property, Banach spaces, AMS Subject Classification: 46 B 20.

1. Introduction

A Banach space X is said to have the Schur property if every weakly convergent sequence in X is norm convergent; equivalently if weakly compact subsets of X are norm compact. It is a known fact that ℓ^1 , the dual space of the null sequences c_0 , has the Schur property. Many other results about the Schur property can be found in the literature. In [3], the authors show that the dual space of the group C^* -algebra $C^*(G)$ of a compact group G has the Schur property. In [1], W. S. Brown shows under a very mild condition that the dual space of every commutative subalgebra of the operator algebra $\mathcal{K}(\mathcal{H})$ of the compact operators on a Hilbert space \mathcal{H} has the Schur property. Continuing this work, in [8], A. Ülger characterizes the closed subspaces and subalgebras of $\mathcal{K}(\mathcal{H})$ whose duals have the Schur property. The characterizations of C^* -algebras whose duals have the Schur property has been given by A. Lau and A. Ülger [3], and of commutative C^* -algebras whose duals have the Schur property by A. Pelczynski and Z. Semadeni [5]. J. Diestel gave the connection of the Schur property with the Dunford-Pettis property [2].

If two Banach spaces have the Schur property, then so do their injective and projective tensor products [4], [6].

In this paper, we prove that if $(X_\alpha)_{\alpha \in I}$ are Banach spaces and $X = (\oplus_{\alpha \in I} X_\alpha)_1$ their l_1 -sum, then the space X has the Schur property if each factor X_α has it. The technique used to prove that ℓ^1 has the Schur property is the inspiration for most of the results [7].

Our notation is quite standard. For any Banach space $\langle X, \| \cdot \| \rangle$, we denote its dual by X^* and its closed unit ball by X_1 . The natural duality between X and X^* is denoted as $\langle u, f \rangle$ or as $\langle f, u \rangle$.

2. Direct sums of spaces with the Schur property

Proposition *Let $(X_\alpha)_{\alpha \in I}$ be Banach spaces and $X = (\oplus_{\alpha \in I} X_\alpha)_1$ be their l_1 -sum. The space X has the Schur property iff each factor X_α has it.*

Proof. If a space has the Schur property, then any closed subspace of the space clearly has the Schur property. Hence the implication is true. For the reverse implication, first recall that a Banach space has the Schur property iff all of its closed separable subspaces have the Schur property. So we can assume that each X_α is separable and take $I = \mathbf{N}$. Since $X = (\oplus_{k \in \mathbf{N}} X_k)_1$ is separable, the closed unit ball of $X^* = (\oplus_{k \in \mathbf{N}} X_k^*)_\infty$ under its w^* -topology is metrizable ([9], II.A.15). It follows that a weakly null sequence $\{a_n\}$ in X is norm-null if any w^* -null sequence $\{f_n\}$ in the unit ball of X^* satisfies $\lim_{n \rightarrow \infty} \langle f_n, a_n \rangle = 0$.

Let then $\{a_n\}_{n \in \mathbf{N}}$ be a weakly null sequence in X . Then $a_n = \{b_{n,k}\}_{k \in \mathbf{N}}$ and for all $k \in \mathbf{N}$, the sequence $\{b_{n,k}\}_{n \in \mathbf{N}}$ is weakly null in X_k . As X_k has the Schur property, $\lim_{n \rightarrow \infty} \|b_{n,k}\| = 0$, for all $k \in \mathbf{N}$. On the other hand, let $\{f_n\}_{n \in \mathbf{N}}$ be a w^* -null sequence in the unit ball of X^* . Each f_n is of the form $f_n = \{g_{n,k}\}_{k \in \mathbf{N}}$, and for all $k \in \mathbf{N}$, the sequence $\{g_{n,k}\}$ is w^* -converging in X_k^* to zero. To prove that $\lim_{n \rightarrow \infty} \langle f_n, a_n \rangle = 0$, it is enough to show that the sum $\sum_{k \in \mathbf{N}} |\langle g_{n,k}, b_{n,k} \rangle|$ converges to zero uniformly in $n \in \mathbf{N}$, i.e. $\sup_{n \in \mathbf{N}} \sum_{k > N} |\langle g_{n,k}, b_{n,k} \rangle| \rightarrow 0$, as $N \rightarrow \infty$. In other words, we have to show that,

$$\forall \varepsilon > 0 \exists N \in \mathbf{N} \forall \mathbf{n} \in \mathbf{N} \sum_{\mathbf{k} > \mathbf{N}} |\langle \mathbf{g}_{\mathbf{n}, \mathbf{k}}, \mathbf{b}_{\mathbf{n}, \mathbf{k}} \rangle| < \varepsilon \quad (*)$$

Assume (*) is false. So, there is an $\varepsilon > 0$ such that,

$$\forall N \in \mathbf{N} \exists \mathbf{n} \in \mathbf{N} \sum_{\mathbf{k} > \mathbf{N}} |\langle \mathbf{g}_{\mathbf{n}, \mathbf{k}}, \mathbf{b}_{\mathbf{n}, \mathbf{k}} \rangle| \geq \varepsilon \quad (**)$$

Consider a sequence of positive numbers $\{\delta_k\}$ such that $\sum_{k=1}^{\infty} \delta_k < \frac{\varepsilon}{4}$. We are going to construct two strictly increasing sequences $\{n_k\}_{k \geq 1}$ and $\{N_k\}_{k \geq 0}$ such that

- 1) $\sum_{p > N_k} \|b_{n_k, p}\| \leq \delta_k$ for each $k \geq 1$,
- 2) $\sum_{p=1}^{N_{k-1}} |\langle g_{n_k, p}, b_{n_{k-1}, p} \rangle| \leq \delta_k$ for each $n \geq n_k$,
- 3) $\sum_{p > N_{k-1}} |\langle g_{n_k, p}, b_{n_k, p} \rangle| \geq \varepsilon$.

Start with $n_1 = 1$ and $N_0 = 0$. Since $a_{n_1} \in X$, i.e. $\sum_{p \in \mathbf{N}} \|b_{n_1, p}\| < \infty$, we can choose N_1 such that $\sum_{p > N_1} \|b_{n_1, p}\| \leq \delta_1$. Since $\{g_{n, 1}\}_{n \in \mathbf{N}}, \dots, \{g_{n, N_1}\}_{n \in \mathbf{N}}$ converge to zero in $(X_1^*, w^*), \dots, (X_{N_1}^*, w^*)$ respectively, there exists $\bar{n}_2 > n_1$ such that

$$\forall n \geq \bar{n}_2, \sum_{p=1}^{N_1} |\langle g_{n, p}, b_{n_1, p} \rangle| \leq \delta_2.$$

By (**), there exists $n_2 \geq \bar{n}_2$ such that

$$\sum_{p > N_1} |\langle g_{n_2, p}, b_{n_2, p} \rangle| \geq \varepsilon.$$

Now let $N_2 > N_1$ such that $\sum_{p > N_2} \|b_{n_2, p}\| \leq \delta_2$.

Since $\{g_{n, 1}\}_{n \in \mathbf{N}}, \dots, \{g_{n, N_2}\}_{n \in \mathbf{N}}$ converge to zero in $(X_1^*, w^*), \dots, (X_{N_2}^*, w^*)$ respectively, there exists $\bar{n}_3 > n_2$ such that

$$\forall n \geq \bar{n}_3, \sum_{p=1}^{N_2} |\langle g_{n, p}, b_{n_2, p} \rangle| \leq \delta_3.$$

Let $n_3 \geq \bar{n}_3$ and $N_3 > N_2$ be such that, $\sum_{p=1}^{N_2} |\langle g_{n_3, p}, b_{n_2, p} \rangle| \leq \delta_3$ and $\sum_{p > N_3} \|b_{n_3, p}\| \leq \delta_3$ and so on...

Now let us choose a sequence $\{\gamma_p\}$ such that

for $N_0 + 1 \leq p \leq N_1$, $\gamma_p \langle g_{n_1,p}, b_{n_1,p} \rangle = |\langle g_{n_1,p}, b_{n_1,p} \rangle|$
 for $N_1 + 1 \leq p \leq N_2$, $\gamma_p \langle g_{n_2,p}, b_{n_2,p} \rangle = |\langle g_{n_2,p}, b_{n_2,p} \rangle|$
 for $N_{k-1} + 1 \leq p \leq N_k$, $\gamma_p \langle g_{n_k,p}, b_{n_k,p} \rangle = |\langle g_{n_k,p}, b_{n_k,p} \rangle|$

We shall define an element h in X^* as

$$h = (\gamma_1 g_{n_1,1}, \gamma_2 g_{n_1,2}, \dots, \gamma_{N_1} g_{n_1,N_1}, \gamma_{N_1+1} g_{n_2,N_1+1}, \gamma_{N_1+2} g_{n_2,N_1+2}, \dots, \gamma_{N_2} g_{n_2,N_2}, \gamma_{N_2+1} g_{n_3,N_2+1}, \gamma_{N_2+2} g_{n_3,N_2+2}, \dots)$$

If we denote h as $h = (h_p)_{p \geq 1}$, then we have $\|h\| = \sup_{p \geq 1} \|h_p\| \leq 1$ and

$$\langle h, a_{n_k} \rangle = \sum_{p=1}^{\infty} \langle h_p, b_{n_k,p} \rangle =$$

$$\sum_{j=1}^{k-1} \sum_{p=N_{j-1}+1}^{N_j} \gamma_p \langle g_{n_j,p}, b_{n_j,p} \rangle + \sum_{p=N_{k-1}+1}^{N_k} |\langle g_{n_k,p}, b_{n_k,p} \rangle| + \sum_{p=N_k}^{\infty} \gamma_p \langle g_{n_k,p}, b_{n_k,p} \rangle.$$

Using the inequalities 1), 2), 3), and $\|g_{n_k,p}\| \leq 1$ we get:

$$\begin{aligned} |\langle h, a_{n_k} \rangle| &\geq - \sum_{j=1}^{k-1} \delta_j + \sum_{p=N_{k-1}+1}^{N_k} |\langle g_{n_k,p}, b_{n_k,p} \rangle| - \delta_k \\ &\geq - \sum_{j=1}^{k-1} \delta_j + \sum_{p > N_{k-1}} |\langle g_{n_k,p}, b_{n_k,p} \rangle| - \sum_{p \leq N_{k+1}} |\langle g_{n_k,p}, b_{n_k,p} \rangle| - \delta_k \\ &\geq - \sum_{j=1}^{k-1} \delta_j + \sum_{p > N_{k-1}} |\langle g_{n_k,p}, b_{n_k,p} \rangle| - 2\delta_k \\ &\geq \epsilon - \sum_{j=1}^{k-1} \delta_j - 2\delta_k \\ &\geq \epsilon - 2 \sum_{j=1}^{\infty} \delta_j > 2\frac{\epsilon}{2}. \end{aligned}$$

This contradicts the fact that $\{a_n\}$ converges weakly to zero. Consequently, (*) holds, that is

$$\lim_{K \rightarrow \infty} \sup_{n \in \mathbb{N}} \sum_{k=1}^K |\langle g_{n,k}, b_{n,k} \rangle| = 0.$$

It follows that

$$\lim_{n \rightarrow \infty} \sum_{k=1}^{\infty} |\langle g_{n,k}, b_{n,k} \rangle| = \sum_{k=1}^{\infty} \lim_{n \rightarrow \infty} |\langle g_{n,k}, b_{n,k} \rangle|.$$

As $\|b_{n,k}\| \rightarrow 0$, $|\langle g_{n,k}, b_{n,k} \rangle| \rightarrow 0$, so that,

$$\lim_{n \rightarrow \infty} \sum_{k=1}^{\infty} |\langle g_{n,k}, b_{n,k} \rangle| = 0.$$

As $|\langle f_n, a_n \rangle| \leq \sum_{k=1}^{\infty} |\langle g_{n,k}, b_{n,k} \rangle|$, we conclude that $\lim_{n \rightarrow \infty} \langle f_n, a_n \rangle = 0$, and so $\|a_n\|$ converges to zero. \square

References

- [1] S. W. Brown, *Weak Sequential Convergence in the Dual of an Algebra of Compact Operators*, J. Operator Theory **33**, (1995), 33-42.
- [2] J. Diestel, *A Survey of Results Related to the Dunford-Pettis Property*, Contemporary Mathematics **2**, AMS (1980), 15-60.
- [3] A. Lau and A. Ülger, *Some Geometric Properties on the Fourier and Fourier Stieltjes Algebras of Locally Compact Groups, Arens Regularity and Related Problems* Trans. Amer. Math. Soc. **337** (1993), 321-359.
- [4] F. Lust, *Produits Tensoriels Injectifs d'Espaces de Sidon*, Colloq. Math. **32**, (1975), 286-289.
- [5] A. Pelczynski and Z. Semadeni, *Spaces of Continuous Functions (III)*, Studia Mathematica T. XVIII, (1959).

TANBAY

- [6] R. Ryan, *The Dunford-Pettis Property and Projective Tensor Products*, Bull. Polish Acad. Sc. **35** No. 11-12, (1987), 785-792.
- [7] B. Tanbay, *A condition for the Schur property*, preprint.
- [8] A. Ülger, *Subspaces and subalgebras of $\mathcal{K}(\mathcal{H})$ whose Duals have the Schur Property*, J. Operator Theory **37** (1997), 371-378.
- [9] P. Wojtaszczyk, *Banach Spaces for Analysts*, Cambr. Studies in Adv. Math. **25**, (1996).

Betül TANBAY
Boğaziçi University

Received 21.10.1998

CONTENTS

<i>A Combinatorial Definition of n-Types of Simplicial Commutative Algebras</i>	243
Z. ARAVASI, M. KOÇAK, M. ALP	
<i>Pullbacks of Crossed Modules and cat^1-Groups</i>	273
M. ALP	
<i>Difference Method for a Singularly Perturbed Initial Value Problem</i>	283
G. M. AMIRALIYEV	
<i>The Norm in Taxicab Geometry</i>	295
C. EKİNCİ, I. KOCAYUSUFOĞLU, Z. AKÇA	
<i>Derivation of Separable Amplitude Equations by Multiple Scales Method</i>	309
M. N. ÖZER, D. ESER	
<i>A New Integrable Reduction of the Coupled Nls Equation</i>	319
M. N. ÖZER	
<i>On The Discrete Squeezing Property for Semilinear Wave Equations</i>	335
A. EDEN	
<i>The Dual of the Bochner Space $L^p(\mu, E)$ For Arbitrary μ</i>	343
B. CENGİZ	
<i>Direct Sums and the Schur Property</i>	349
B. TANBAY	