

1-1-1998

On the Parabolic Class Number of Some Subgroups of Hecke Groups

R. KESKİN

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

KESKİN, R. (1998) "On the Parabolic Class Number of Some Subgroups of Hecke Groups," *Turkish Journal of Mathematics*: Vol. 22: No. 2, Article 8. Available at: <https://journals.tubitak.gov.tr/math/vol22/iss2/8>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

ON THE PARABOLIC CLASS NUMBER OF SOME SUBGROUPS OF HECKE GROUPS

Refik Keskin

Abstract

In this paper we calculate the parabolic class number of subgroups of Hecke groups $H(\sqrt{2}), H(\sqrt{3})$.

Subject Classification: 20 G-H

Keywords: Parabolic class number, orbit Fuchsian group.

1. Introduction

By a Fuchsian group Λ we will mean a finitely generated discrete subgroup of $PSL(2, \mathbb{R})$, the group of conformal homeomorphisms of the upper-half plane. The most general presentation for Λ is

Generators;

$$\begin{array}{ll} a_1, b_1, \dots, a_g, b_g & \text{(Hyperbolic)} \\ x_1, x_2, \dots, x_r & \text{(Elliptic)} \\ p_1, p_2, \dots, p_s & \text{(Parabolic)} \end{array}$$

Relations;

$$x_1^{m_1} = x_2^{m_2} = \dots = x_r^{m_r} = \prod_{i=1}^g [a_i, b_i] \prod_{j=1}^r x_j \prod_{k=1}^s p_k = 1.$$

We then say Λ has signature (see [1])

$$(g, m_1, m_2, \dots, m_r; s).$$

Hecke introduced an infinite class of discrete groups $H(\lambda_q)$ of linear fractional transformation preserving the upper-half plane. $H(\lambda_q)$ is the group generated by

$$S(z) = z + \lambda_q, \quad T(z) = \frac{-1}{z}$$

where $\lambda_q = 2 \cos(\pi/q)$, q is an integer, $q \geq 3$. When $q = 3$ we have the modular group Γ . When $q = 4$ or 6 the resulting group are $H(\sqrt{2}), H(\sqrt{3})$. These two groups are of particular interest since they are the only Hecke groups, aside from the modular group, whose elements are completely known.

It is well known ([2], [3]) that $H(\sqrt{m}), m = 2, 3$, consists of the mappings of all the following two types.

- i) $T(z) = \frac{az + b\sqrt{m}}{c\sqrt{m}z + d}, a, b, c, d \in \mathbb{Z}, ad - bcm = 1,$
- ii) $T(z) = \frac{a\sqrt{m}z + b}{cz + d\sqrt{m}}, a, b, c, d \in \mathbb{Z}, adm - bc = 1.$

2. Parabolic Class Number

From now on, m will stand for 2 or 3

Lemma 1. $H(\sqrt{m})$ act transitively on $\sqrt{m}\hat{\mathbb{Q}} = \{\frac{x}{s}\sqrt{m} : \frac{x}{s} \in \mathbb{Q}\} \cup \{\infty\}$.

$\sqrt{m}\mathbb{Q}$ is the largest subset of \mathbb{R} on which $H(\sqrt{m})$ acts transitively.

Proof. Let $\frac{x}{y}\sqrt{m} \in \sqrt{m}\hat{\mathbb{Q}} \setminus \{\infty\}$ with $(x, y) = 1$. Then $m|y$ or $m \nmid y$. Since $(x, y) = 1$, we can find $a, b \in \mathbb{Z}$ such that $xa - yb = 1$. If $m|y$, then we take

$$T(z) = \frac{xz + b\sqrt{m}}{\frac{x}{m}\sqrt{m}z + a},$$

so, we have $T(\infty) = \frac{x}{y}\sqrt{m}$.

Let $m \nmid y$. In this case $(mx, y) = 1$, and thus there exist some, $a, b \in \mathbb{Z}$ such that $mxa - yb = 1$. Similarly, if we take

$$S(z) = \frac{x\sqrt{m}z + b}{yz + \sqrt{ma}},$$

then $S(\infty) = \frac{x}{y}\sqrt{m}$.

Let $n \in \mathbb{N}$. Define

$$H_0^m(n) = \{T \in H(\sqrt{m}) : c \equiv 0 \pmod{n}\}.$$

Then $H_0^m(n)$ is a subgroup of $H(\sqrt{m})$. □

Let Γ be the modular group, and $\Gamma_0(n)$ be the subgroup of Γ such that $c \equiv 0 \pmod{n}$. Then,

Lemma 2. [5] $|\Gamma : \Gamma_0(n)| = n \prod_{p|n} \left(1 + \frac{1}{p}\right)$.

Lemma 3. If $(m, n) = 1$, then $|H(\sqrt{m}) : H_0^m(n)| = n \prod_{p|n} \left(1 + \frac{1}{p}\right)$, if $(m, n) = m$,

then $|H(\sqrt{m}) : H_0^m(n)| = 2n \prod_{p|n} \left(1 + \frac{1}{p}\right)$ where $p \neq m$.

Proof. We will give the proof in case where $(m, n) = m$.

Let

$$H = \left\{ T \in H(\sqrt{m}) : T(z) = \frac{az + b\sqrt{m}}{c\sqrt{m}z + d} \right\}.$$

As $m|n$, $H_0^m(n) \subset H \subset H(\sqrt{m})$. It is obvious that $|H(\sqrt{m}) : H| = 2$. Let θ be the mapping from H to $\Gamma_0(m)$ defined as follows. If

$$T(z) = \frac{az + b\sqrt{m}}{c\sqrt{m}z + d},$$

then

$$\theta(T)(z) = \frac{az + b}{cmz + d}.$$

It can be shown that θ is an isomorphism, and $\theta(H_0^m(n)) = \Gamma_0(mn)$.

On the other hand, $\Gamma_0(mn) \subset \Gamma_0(m) \subset \Gamma$. Therefore,

$$\begin{aligned} |H(\sqrt{m}) : H_0^m(n)| &= |H(\sqrt{m}) : H| |H : H_0^m(n)| = 2|\Gamma_0(m) : \Gamma(mn)| \\ &= 2 \frac{|\Gamma : \Gamma_0(mn)|}{|\Gamma : \Gamma_0(m)|} = 2n \prod_{p|n} \left(1 + \frac{1}{p}\right) \text{ where } p \neq m. \end{aligned}$$

If $(m, n) = 1$, the proof can be done in a similar way. □

We now give the following lemmas without proof. Proofs are similar to those for the modular group Γ in [4].

Lemma 4. Let Λ be a subgroup of finite index in $H(\sqrt{m})$. Then, the parabolic class number of Λ is the number of orbits of Λ on $\sqrt{m}\hat{\mathbb{Q}}$.

Lemma 5. *Let Λ be a subgroup of finite index in $H(\sqrt{m})$. Then, the parabolic class number s of Λ satisfies $1 \leq s \leq N$, where N is the index $|H(\sqrt{m}) : \Lambda|$; in particular s is finite.*

We now give our main theorems.

Theorem 1. *If $(m, n) = 1$, then the parabolic class number of $H_0^m(n)$ is*

$$\sum_{d|n} \varphi\left(\left(d, \frac{n}{d}\right)\right)$$

Before giving the proof we will give some lemmas.

Lemma 6. *Let $(m, n) = 1$, and $\frac{r}{s}\sqrt{m} \in \sqrt{m}\hat{\mathbb{Q}}$ with $m|s$, then we can find some $T \in H_0^m(n)$ such that $T(\frac{r}{s}\sqrt{m}) = \frac{r_1}{s_1}\sqrt{m}$ with $(m, s_1) = 1$ (we represent ∞ as $\frac{1}{0}\sqrt{m}$).*

Proof. Since $(m, n) = 1$, there exist some $a, b \in \mathbb{Z}$ such that $1 = ma - nb$. Let

$$T(z) = \frac{a\sqrt{m}z + b}{nz + \sqrt{m}}.$$

Then $T \in H_0^m(0)$, and

$$T\left(\frac{r}{s}\sqrt{m}\right) = \frac{arm + bs}{(rn + s)\sqrt{m}} = \frac{ar + bs/m}{rn + s}\sqrt{m}.$$

It can be easily shown that $(m, rn + s) = 1$. If we take $r_1 = ar + bs/m$, and $s_1 = rn + s$, then $T(\frac{r}{s}\sqrt{m}) = \frac{r_1}{s_1}\sqrt{m}$ with $(m, s_1) = 1$. \square

Lemma 7. *Let $(m, n) = 1$, and $\frac{k}{s}\sqrt{m} \in \sqrt{m}\hat{\mathbb{Q}}$ with $(k, s) = 1$. If $(m, s) = 1$, then there exist some $T \in H_0^m(n)$ such that*

$$T\left(\frac{k}{s}\sqrt{m}\right) = \frac{k_1}{s_1}\sqrt{m} \text{ with } s_1|n.$$

Proof. $(km, s) = 1$ since $(k, s) = (m, s) = 1$. Let $s_1 = (s, n)$. Then $s_1 = (s, n) = (s, kmn)$. Therefore there exist some integers c_1, d_1 such that

$$\frac{kmn}{s_1}c_1 + \frac{s}{s_1}d_1 = 1.$$

Since $(d_1, \frac{kmn}{s_1}) = 1$, there exists an integer t such that $(d_1 - \frac{kmn}{s_1}t, mn) = 1$. Let $d = d_1 - \frac{kmn}{s_1}t$ and $c = c_1 + \frac{s}{s_1}t$. Then

$$\frac{kmn}{s_1}c + \frac{s}{s_1}d = 1.$$

□

On the other hand, $(d, cmn) = 1$, since $(d, mn) = (d, c) = 1$. Hence, we can find some integers x, y such that $xd - ycmn = 1$. If we take

$$T(z) = \frac{xz + y\sqrt{m}}{cn\sqrt{m}z + d},$$

then, we have $T(\frac{k}{s}\sqrt{m}) = \frac{k_1}{s_1}\sqrt{m}$ where $k_1 = xk + ys$ and $s_1 = cnmk + ds$.

It is obvious that $T \in H_0^m(n)$. On the other hand, it can be seen that $(k_1, s_1) = 1$.

Lemma 8. *Let $(m, n) = 1$. If $d_1|n$ and $(a_1d_1) = (a_2, d_1) = 1$, then $\frac{a_1}{d_1}\sqrt{m}$ is conjugate to $\frac{a_2}{d_1}\sqrt{m}$ under $H_0^m(n)$ if and only if $a_1 \equiv a_2 \pmod{t}$ where $t = (d_1, \frac{n}{d_1})$.*

Proof. Let $a_1 \equiv a_2 \pmod{t}$ and $n_1 = n/d_1$. Then, $t = (d_1, n_1)$, and (a_1a_2, d_1) , and $(a_1a_2, d_1) = 1$. Furthermore, $(m, d_1) = 1$ since $(m, n) = 1$. Therefore $(a_1a_2m, d_1) = 1$, and thus $(n_1a_1a_2m, d_1) = t$. Since $t|a_1 - a_2$, $mn_1a_1a_2x + d_1y = a_2 - a_1$ has a solution. That is, there exist some integers k, s such that $mn_1a_1a_2k + a_1 + d_1s = a_2$. Hence, we obtain $aa_1 + bd_1 = a_2$. On the other hand, if we take $c = n_1d_1k$ and $d = 1 - mn_1a_1k$, we obtain $mca_1 + dd_1 = d_1$.

Furthermore,

$$ad - bcm = a(1 - mn_1a_1k) - bmn_1d_1k = a - (aa_1 + bd_1)mn_1k = 1.$$

Let

$$T(z) = \frac{az + b\sqrt{m}}{c\sqrt{m}z + d}.$$

Then it is clear that $T \in H_0^m(n)$ and $T(\frac{a_1}{d_1}\sqrt{m}) = \frac{a_2}{d_1}\sqrt{m}$.

Now let $\frac{a_1}{d_1}\sqrt{m}$ be equivalent to $\frac{a_2}{d_1}\sqrt{m}$ by some $T \in H_0^m(n)$. Then it is easily seen that

$$T(z) = \frac{az + b\sqrt{m}}{cn\sqrt{m}z + d} \text{ where } ad - bcmn = 1.$$

Hence, we obtain

$$\frac{aa_1 + bd_1}{cna_1m + dd_1}\sqrt{m} = \frac{a_2}{d_1}\sqrt{m},$$

that is,

$$\frac{aa_1 + bd_1}{cna_1m + dd_1} = \frac{a_2}{d_1}.$$

Since

$$d(aa_1 + bd_1) - b(cna_1m + dd_1) = a_1,$$

and

$$a(cna_1m + dd_1) - cnm(aa_1 + bd_1) = d_1$$

we have $(aa_1 + bd_1, cna_1m + dd_1) = 1$. Therefore, there exists some $u = \pm 1$ such that

$$aa_1 + bd_1 = ua_2,$$

and

$$cna_1m + dd_1 = ud_1.$$

It can be easily shown that $a_1 \equiv a_2 \pmod{t}$. □

Proof of Theorem 1. It is sufficient to calculate the number of orbits of $H_0^m(n)$ on $\sqrt{m}\hat{\mathbb{Q}}$. Then from Lemma 6, Lemma 7, and Lemma 8, the number of orbits of $H_0^m(n)$ on $\sqrt{m}\hat{\mathbb{Q}}$ is $\sum_{d|n} \varphi\left(d, \frac{n}{d}\right)$ where φ is Euler's function.

We can deduce the following.

Lemma 9. *If $\frac{k}{s} \in \hat{\mathbb{Q}}$ with $(k, s) = 1$, then there exists some $T \in \Gamma_0(n)$ such that $T\left(\frac{k}{s}\right) = \frac{k_1}{s_1}$ with $s_1|n$ where we represent ∞ as $\frac{1}{0}$.*

Lemma 10. *If $d_1|n$ and $(a_1, d_1) = (a_2, d_1) = 1$, then $\frac{a_1}{d_1}$ is conjugate to $\frac{a_2}{d_1}$ under $\Gamma_0(n)$ if and only if $a_1 \equiv a_2 \pmod{t}$ where $t = (d_1, \frac{n}{d_1})$.*

Then it is easily seen that the number of orbits of $\Gamma_0(n)$ on $\hat{\mathbb{Q}}$ is

$$\sum_{d|n} \varphi\left(d, \frac{n}{d}\right),$$

which is the parabolic class number of $\Gamma_0(n)$.

Theorem 2. *If $m|n$, then the parabolic class number of $H_0^m(n)$ is*

$$\sum_{d|n} \varphi\left(\left(d, \frac{nm}{d}\right)\right).$$

Proof. If $m|n$, then

$$H_0^m(n) = \{T \in H : c \equiv 0 \pmod{n}\}.$$

We define ϕ from the set of orbits of $\Gamma_0(mn)$ to the set of orbits of $H_0^m(n)$ as follows.

$$\phi\left(\Gamma_0(mn)\frac{r}{s}\right) = H_0^m(n)\frac{r}{s}\sqrt{m}.$$

Then ϕ is well defined one to one and onto function. The proof then follows. \square

References

- [1] Singerman, D.: Subgroups of Fuchsian groups and finite permutation groups, *Bulletin London Math. Soc.*, **2**, 319-323 (1970).
- [2] Hutchinson, J. I.: On a class of automorphic functions, *Trans. Amer. Math. Soc.*, **5**, 1-11 (1902).
- [3] Young, J.: On the group belonging to the sign $(0, 3; 2, 4, \infty)$ and the functions belonging to it. *Trans. Amer. Math. Soc.*, **5**, 81-104 (1904).
- [4] Jones, G. A., and Singerman, D.: *Complex Functions: an algebraic and geometric viewpoint*. Cambridge, Cambridge University Press, 1987.
- [5] Schoeneberg, B.: *Elliptic Modular Functions*. Berlin, Heidelberg, New York, Springer Verlag 1974.
- [6] Miyake, T.: *Modular Forms*. Springer Verlag 1989.
- [7] Akbaş, M.: *The Normalizer of Modular Subgroups*. Ph. D. Thesis, Southampton University.

Refik KESKİN
Karadeniz Technical University
Department of Mathematics
61080 Trabzon-TURKEY

Received 16.04.1997