

1-1-1997

## Augmented Graded Rings

Mashhoor REFAI

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

REFAI, Mashhoor (1997) "Augmented Graded Rings," *Turkish Journal of Mathematics*: Vol. 21: No. 3, Article 10. Available at: <https://journals.tubitak.gov.tr/math/vol21/iss3/10>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## AUGMENTED GRADED RINGS

*Mashoor Refai*

### Abstract

In this paper we study the augmented graded rings and give the relationship between these rings and stronger properties of graded rings.

### 0.Introduction.

Let  $G$  be a multiplicative group with identity  $e$  and  $R$  be an associative  $G$ -graded ring with unity 1. Then  $R_e$  (the identity component of  $R$ ) is a subring of  $R$  and  $1 \in R_e$  (See [5]). One of the most important problems in graded ring theory is to study the link between a certain property for  $R$  or  $(R, G)$  and  $R_e$ .

Now, one can think about grading  $R_e$  by a group  $G$ . This types of rings appear naturally when we deal with the group ring of  $G$  over a  $G$ -graded ring  $R$ .

In this paper, we discuss the  $G$ -graded rings in which the identity component is itself a  $G$ -graded ring satisfying some related conditions with the graduation of  $G$ . We call these rings augmented  $G$ -graded rings. Also, we give the relationship between these rings and the stronger properties of  $G$ -graded rings given in [8]. This simple effort will help for studying those rings in details and that will be a good tool to solve many problems in Graded Ring Theory.

### 1.Preliminaries.

In this section we give some basic definitions and facts of graded rings. For more details one can look in [8].

**Definition 1.1.** Let  $G$  be a group with identity  $e$ . Then a ring  $R$  is  $G$ -graded if there exist additive subgroups  $R_g$  of  $R$  such that  $R = \bigoplus_{g \in G} R_g$  and  $R_g R_h \subseteq R_{gh}$  for all  $g, h \in G$ . We denote for this graduation by  $(R, G)$  and  $R_e$  will be the identity component of  $R$ . The elements of  $R_g$  are called homogeneous of dimension  $g$ . For  $x \in R$ , we write  $x_g$  for the component of  $x$  in  $R_g$ , so that  $x$  can be written uniquely as  $\sum_{g \in G} x_g$ .

We denote  $\text{supp} (R, G) = \{g \in G : R_g \neq 0\}$ . Note that  $\text{supp} (R, G)$  need not be a subgroup of  $G$ .

**Definition 1.2.** For a  $G$ -graded ring  $R$  we say :

- 1)  $(R, G)$  is strong if  $R_g R_h = R_{gh}$  for all  $g, h \in G$ . But this definition is equivalent to  $1 \in R_g R_{g^{-1}}$  for all  $g \in G$  (See proposition 1.6 of [3]).
- 2)  $(R, G)$  is first strong if  $1 \in R_g R_{g^{-1}}$  for all  $g \in \text{supp} (R, G)$ .
- 3)  $(R, G)$  is second strong if  $R_g R_h = R_{gh}$  for all  $g, h \in \text{supp} (R, G)$  and  $\text{supp} (R, G)$  is a monoid in  $G$ .
- 4)  $(R, G)$  is faithful if for any  $a_g \in R_{g^{-1}}$ ,  $a_g R_h \neq 0$  and  $R_h a_g \neq 0$  for all  $h \in G$ .
- 5)  $(R, G)$  is nondegenerate if for any  $a_g \in R_g$ ,  $a_g R_{g^{-1}} \neq 0$  and  $R_{g^{-1}} a_g \neq 0$ .

The nondegenerate and faithful properties are motivated by the work of Cohen and Rowen in [1]. The strongly graded rings have been studied by E. C.Dade in [2], where they were called clifford systems. In [8] we studied the other stronger properties and gave relationship between all of them.

**Lemma 1.3** ([5]).  $R_e$  is a subring of  $R$  and  $1 \in R_e$ .

## 2. Augmented Graded Rings.

In this section we study the augmented graded rings and then give the relationship between these rings and some stronger properties  $G$ -graded rings.

**Definition 2.1** A ring  $R$  is said to be an augmented  $G$ -graded ring if it satisfies the following conditions:

- 1)  $R = \bigoplus_{g \in G} R_g$  where  $R_g$  is an additive subgroup of  $R$  and  $R_g R_h \subseteq R_{gh}$  for all  $g, h \in G$ . ( $R$  is a  $G$ -graded ring).
- 2) If  $R_e$  is the identity component of the graduation given in 1 then  $R_e = \bigoplus_{g \in G} R_{e-g}$  where  $R_{e-g}$  is an additive subgroup of  $R_e$  and  $R_{e-g} R_{e-h} \subseteq R_{e-gh}$  for all  $g, h \in G$ . ( $R_e$  is a  $G$ -graded ring).
- 3) For each  $g \in G$ , there exists  $r_g \in R_g$  such that  $R_g = \bigoplus_{h \in G} R_{e-h} r_g$  we assume  $r_e = 1$ .

- 4) If  $g, h \in G$  and  $r_g, r_h$  are both non-zero, then  $r_g r_h = r_{gh}$  and  $(x r_g)(y r_h) = xy r_{gh}$  for all  $x, y \in R_e$ .

We need condition 3 of the definition to give a connection between the graduation of  $R$  and the graduation of  $R_e$ . Also, condition 4 is necessary for proving some of the following useful remarks.

**Remarks 2.2.**

- 1) Condition 3 of the definition implies  $R_h = R_e r_h$  for all  $h \in G$ .

To see this, let  $x \in R_h$ . Then  $x = s_{e-g_1} r_h + \dots + s_{e-g_n} r_h$  where  $s_{e-g_i} \in R_{e-g_i}$ , and hence  $x = (s_{e-g_1} + \dots + s_{e-g_n}) r_h \in R_e r_h$ . Conversely, let  $s r_h \in R_e r_h$ . Then  $s r_h = (s_{e-g_1} r_h + \dots + s_{e-g_n} r_h) \in R_h$  where  $s_{e-g_i} \in R_{e-g_i}$ .

- 2) Condition 4 of the definition is equivalent to :

$(x_{e-h_1} r_g)(y_{e-h_2} r_h) = x_{e-h_1} y_{e-h_2} r_{gh}$  for all  $x_{e-h_1} \in R_{e-h_1}, y_{e-h_2} \in R_{e-h_2}$  and non-zero  $r_g, r_h$ . Clearly condition 4 implies this condition. Let  $x, y \in R_e$ . For simplicity we assume  $x = x_{e-h_1} + x_{e-h_2}$  and  $y = y_{e-g_1} + y_{e-g_2}$ . Then  $(x r_g)(y r_h) = x_{e-h_1} y_{e-g_1} r_{gh} + x_{e-h_1} y_{e-g_2} r_{gh} + x_{e-h_2} y_{e-g_1} r_{gh} + x_{e-h_2} y_{e-g_2} r_{gh} = (xy) r_{gh}$ .

- 3)  $R_g$  is a  $G$ -graded  $R_e$ -module with the usual multiplication on  $R$  and with the graduation  $R_{g-h} = R_{e-h} r_g$  for all  $h \in G$ .

- (i)  $R_{g-h}$  is an additive subgroup of  $R_g$ .
- (ii)  $R_{e-e} R_{g-h} = R_{e-e} R_{e-h} r_g \subseteq R_{e-h} r_g, i.e., R_{g-h} \in R_{e-e}$  -mod.
- (iii)  $R_{e-h_1} R_{g-h_2} = (R_{e-h_1} R_{e-h_2}) r_g \subseteq R_{e-h_1 h_2} \quad r_g = R_{g-h_1 h_2}$ .

Clearly  $R_g = \bigoplus_{h \in G} R_{g-h}$ .

- 4)  $R_{g-h} R_{g'-h'} \subseteq R_{gg'-hh'}$  for all  $g, g', h, h' \in G$ .  $R_{g-h} R_{g'-h'} = R_{e-h} r_g R_{e-h'} r_{g'}$ . If  $r_g = 0$  or  $r_{g'} = 0$  then done. Suppose  $r_g, r_{g'}$  are both non zero. By condition 4 of the definition we have  $R_{g-h} R_{g'-h'} = R_{e-h} R_{e-h'} r_{gg'} \subseteq R_{e-hh'} \quad r_{gg'} = R_{gg'-hh'}$ .
- 5) If  $r_g, r_{g^{-1}}$  are both non-zero then  $r_g R_e = R_e r_g$ . Let  $x \in R_e$ . Then  $r_g x = r_g x.1 = (r_g x) r_e = (r_g x)(r_{g^{-1}} r_g) = (1.r_g)(x r_{g^{-1}}) (1.r_g) = x r_g r_{g^{-1}} r_g = x r_g$ .

In the following example , we see that if  $R$  is an associative ring with unity 1 and  $G$  is a group with identity  $e$  then  $R$  can be regarded as an augmented  $G$ -graded ring.

**Example 2.3.** Let  $R$  be any ring and  $G$  be any group. Then  $R$  is an augmented  $G$ -graded ring with :

$$R_g = \begin{cases} R & \text{if } g = e \\ O & \text{if } g \neq e \end{cases}, R_{e-g} = \begin{cases} R & \text{if } g = e \\ O & \text{if } g \neq e \end{cases},$$

$$\text{and } r_g = \begin{cases} 1 & \text{if } g = e \\ O & \text{if } g \neq e \end{cases}.$$

1. Clearly  $R_g$  is an additive subgroup of  $R$ ,  $R_g R_h \subseteq R_{gh}$  for all  $g, h \in G$  and  $R = \bigoplus_{g \in G} R_g$ .
2.  $R_{e-g}$  is an additive subgroup of  $R_e$ ,  
 $R_{e-g} R_{e-h} \subseteq R_{e-gh}$  and  $R_e = \bigoplus_{g \in G} R_{e-g}$ .
3.  $R_g = \bigoplus_{h \in G} R_{e-h} r_g$ .
4. If  $r_g, r_h$  are both non-zero then  $g = h = e$ , and hence  $r_g r_h = 1 = r_e = r_{gh}$ . Also,  $(x r_g)(y r_h) = x y r_{gh}$  for all  $x, y \in R_e$ . We call this graduation the trivial augmented graduation of  $R$  by  $G$ .

Now, we will try to connect the augmented graded rings with the following facts which we proved in [8].

**Proposition 2.4 ([8]).**

1.  $(R, G)$  strong  $\implies (R, G)$  first strong  $\implies (R, G)$  second strong.
2.  $(R, G)$  strong  $\implies (R, G)$  first strong  $\implies (R, G)$  nondegenerate.
3.  $(R, G)$  strong  $\implies (R, G)$  faithful  $\implies (R, G)$  nondegenerate.
4.  $(R, G)$  first strong  $\iff (R, G)$  is second strong and nondegenerate.
5.  $(R, G)$  strong  $\iff (R, G)$  is second strong and faithful.
6.  $(R, G)$  second strong and  $\text{supp } (R, G) \leq G \implies (R, G)$  first strong.
7.  $(R, G)$  second strong and  $G$  finite group  $\implies (R, G)$  first strong.

**Remark 2.5.** If  $|G| \geq 2$  then the trivial augmented  $G$ -graded rings are not faithful. In fact,  $\text{supp}(R, G) = \text{supp}(R_e, G) = \{e\}$ . Therefore, if  $R$  is an augmented  $G$ -graded ring then  $(R, G)$  is not necessarily faithful.

**Proposition 2.6.** Let  $(R, G)$  be augmented  $G$ -graded ring and  $\text{supp}(R, G) = G$ . Then  $R$  is a strongly  $G$ -graded ring.

**Proof.** : Suppose  $(R, G)$  is augmented  $G$ -graded ring. Let  $g \in G$ . Since  $\text{supp}(R, G) = G$ ,  $R_g$  and  $R_{g^{-1}}$  are non-zero and hence there exist non-zero elements  $r_g \in R_g$  and  $r_{g^{-1}} \in R_{g^{-1}}$ . By condition 4,  $1 = r_e = r_g r_{g^{-1}} \in R_g R_{g^{-1}}$ , i.e.,  $1 \in R_g R_{g^{-1}}$  and hence  $R$  is a strongly  $G$ -graded ring.

From this proposition we notice that the augmented graded rings are very useful in case  $\text{supp}(R, G) = G$ . In the following example,  $R = K[x]$  is an augmented  $\mathbb{Z}$ -graded ring with  $\text{supp}(R, \mathbb{Z}) = \mathbb{N} \cup \{o\}$  and  $\text{supp}(R_o, \mathbb{Z}) = \{o\}$ . But in Example 2.9, we use the same ring with different graduations of  $R$  and  $R_o$  such that  $\text{supp}(R, \mathbb{Z}) = \{o\}$  and  $\text{supp}(R_o, \mathbb{Z}) = \mathbb{N} \cup \{o\}$ .  $\square$

**Example 2.7.** Let  $R = K[x]$ . Then  $R$  is an augmented  $\mathbb{Z}$ -graded ring with :

$$R_j = \begin{cases} K & \text{if } j = O \\ Kx^j & \text{if } j > O \\ O & \text{if } j < O \end{cases}, R_{o-i} = \begin{cases} K & \text{if } i = O \\ O & \text{if } i \neq O \end{cases},$$

$$\text{and } r_j = \begin{cases} 1 & \text{if } j = O \\ x^j & \text{if } j > O \\ O & \text{if } j < O \end{cases}.$$

1.  $R = \bigoplus_{j \in \mathbb{Z}} R_j$  ( This is the usual graduation of  $K[x]$  by  $\mathbb{Z}$ ), and  $\text{supp}(R, \mathbb{Z}) = \mathbb{N} \cup \{o\}$ .
2.  $R_o = \bigoplus_{i \in \mathbb{Z}} R_{o-i}$  (This is the trivial graduation of  $K$  by  $\mathbb{Z}$ ), and  $\text{supp}(R_o, \mathbb{Z}) = \{o\}$ .
3. Clearly  $R_j = \bigoplus_{i \in \mathbb{Z}} R_{o-i} r_j$ .
4. If  $r_i, r_j$  are non-zero then  $i, j \geq o$  and hence  $r_i r_j = r_{i+j}$ . Since  $R$  is commutative ring we have  $(x r_i)(y r_j) = (xy) r_{i+j}$ .

**Remark 2.8.** If  $R$  is an augmented  $G$ -graded ring then  $(R, G)$  is not necessarily nondegenerate. In the previous example choose  $g = 1$  and  $a_g = x$ . Then  $a_g \in R_{g-o}$ ,  $R_{g^{-1}} = R_{-1} = o$  and hence  $a_g R_{g^{-1}} = o$ , i.e.,  $(R, G)$  is degenerate.

**Example 2.9.** Let  $R = K[x]$  and  $G = \mathbb{Z}$ . Then  $R$  is augmented  $G$ -graded ring with :

$$R_j = \begin{cases} R & \text{if } j = 0 \\ O & \text{if } j \neq 0 \end{cases}, R_{o-j} = \begin{cases} K & \text{if } j = 0 \\ Kx^j & \text{if } j > 0 \\ O & \text{if } j < 0 \end{cases},$$

$$\text{and } r_j = \begin{cases} 1 & \text{if } j = 0 \\ O & \text{if } j \neq 0 \end{cases}.$$

1.  $R = \bigoplus_{j \in \mathbb{Z}} R_j$ , and  $\text{supp}(R, \mathbb{Z}) = \{0\}$ .
2.  $R_o = \bigoplus_{i \in \mathbb{Z}} R_{o-i}$ , and  $\text{supp}(R_o, \mathbb{Z}) = \mathbb{N} \cup \{0\}$ . Conditions 3 and 4 are obvious.

Now, we give two essential examples of augmented  $G$ -graded rings. In fact, we have built the concept augmented  $G$ -graded ring by taking the common properties between those examples.

**Example 2.10.** Let  $R$  be a  $G$ -graded ring and let  $R[G]$  be the group ring of  $R$  over  $G$ . Then  $R[G]$  is an augmented  $G$ -graded ring with :  $R[G]_g = R.g, R[G]_{e-g} = R.g.e$  and  $r_g = 1.g$  for all  $g \in G$ .

1.  $R[G]_g$  is an additive subgroup of  $R[G]$ ,  $R[G]_g R[G]_h = (R.g)(R.h) = R.g.h = R[G]_{gh}$  and  $R[G] = \bigoplus_{g \in G} R[G]_g$ . Also,  $\text{supp}(R[G], G) = G$ .
2.  $R[G]_{e-g} = R.g.e$  is an additive subgroup of  $R[G]_e, R[G]_{e-g} R[G]_{e-h} = R.g.e.R.h.e \subseteq R.g.h.e = R[G]_{e-gh}$  and  $R[G]_e = \bigoplus_{g \in G} R[G]_{e-g}$ . Also,  $\text{supp}(R[G]_e, G)$  is not necessarily  $G$ .
3.  $R[G]_g = R.g = \bigoplus_{h \in G} R.h.g = \bigoplus_{h \in G} (R.h.e)(1.g) = \bigoplus_{h \in G} R[G]_{e-h} r_g$ .
4.  $r_g r_h = (1.g)(1.h) = 1.gh = r_{gh}$ . Let  $x, y \in R$  and  $x.e, y.e \in R[G]_e$ . Then  $(x.e r_g) = (y.e r_h) = (x.e)(1.g)(y.e)(1.h) = (x.g)(y.h) = xy.gh = (x.e)(y.e)r_{gh}$ .

**Example 2.11.** Let  $R$  be a  $G$ -graded ring. Let  $\overline{R[G]}$  be the left free  $R$ -module with basis  $G$ .

Let  $\lambda_\sigma \in R_\sigma$  and  $\lambda_{\sigma'} \in R_{\sigma'}$ , then for the elements  $\lambda_\sigma \tau, \lambda_{\sigma'} \tau'$  define  $(\lambda_\sigma \tau)(\lambda_{\sigma'} \tau') = (\lambda_\sigma \lambda_{\sigma'}) \sigma'^{-1} \tau \sigma' \tau'$ . With this product  $\overline{R[G]}$  is an associative ring with unity  $1.e$  (See [4]).

Now  $\overline{R[G]}$  is an augmented  $G$ -graded ring with :

$$\overline{R[G]}_g = \bigoplus_{\sigma \in G} R_{g\sigma^{-1}\sigma}, \overline{R[G]}_{e-g} = R_g g^{-1} \text{ and } r_g = 1.g \text{ for all } g \in G.$$

1.  $\overline{R[G]}_g$  is an additive subgroup of  $\overline{R[G]}$ , and  $\overline{R[G]}_g \overline{R[G]}_h \subseteq \overline{R[G]}_{gh}$  because  $R_{g\sigma^{-1}\sigma} R_{h\tau^{-1}\tau} \subseteq R_{g\sigma^{-1}h\tau^{-1}\tau} h^{-1}\sigma h \subseteq \overline{R[G]}_{gh}$  for all  $\sigma, \tau \in G$ . Also,  $\overline{R[G]} = \bigoplus_{g \in G} \overline{R[G]}_g$ .
2.  $\overline{R[G]}_{e-g} = R_g g^{-1}$  is an additive subgroup of  $R[G]_e$ ,  $\overline{R[G]}_{e-g} \overline{R[G]}_{e-h} = (R_g g^{-1})(R_h h^{-1}) \subseteq R_{gh} h^{-1} g^{-1} = \overline{R[G]}_{e-gh}$  and  $\overline{R[G]}_e = \bigoplus_{g \in G} \overline{R[G]}_{e-g}$ .
3.  $\overline{R[G]}_h = \bigoplus_{\sigma \in G} R_{h\sigma^{-1}\sigma}$ . But  $R_{h\sigma^{-1}\sigma} = (R_{h\sigma^{-1}\sigma} h^{-1})(1.h) = (R_{h\sigma^{-1}\sigma} h^{-1})r_h$ , i.e.,  $\overline{R[G]}_h = \bigoplus_{g \in G} \overline{R[G]}_{e-g} r_h$  for all  $h \in G$ .
4. Let  $\lambda_\sigma \sigma^{-1}, \lambda_\tau \tau^{-1} \in R[G]_e$  and  $g, h \in G$ . Then  $(\lambda_\sigma \sigma^{-1})r_h (\lambda_\tau \tau^{-1})r_g = (\lambda_\sigma \sigma^{-1})(1.h)(\lambda_\tau \tau^{-1})(1.g) = (\lambda_\sigma \sigma^{-1} h)(\lambda_\tau \tau^{-1} g) = \lambda_\sigma \lambda_\tau (\tau^{-1} \sigma^{-1} hg)$ . On the other hand,  $(\lambda_\sigma \sigma^{-1})(\lambda_\tau \tau^{-1})r_{hg} = (\lambda_\sigma \lambda_\tau \tau^{-1} \sigma^{-1})(1.hg) = \lambda_\sigma \lambda_\tau (\tau^{-1} \sigma^{-1} hg)$ . Clearly,  $r_h r_g = (1.h)(1.g) = 1.hg = r_{hg}$ .

**Remark 2.12.** If  $R$  is an augmented  $G$ -graded ring, then  $R \in R_{e-gr}$  with the usual product in  $R$  and with  $R_{(g)} = \bigoplus_{h \in G} R_{h-g}$ .

1.  $R_{(g)}$  is an additive subgroup of  $R$  and  $R_{(g)} \in R_{e-e}$ -mod.
2.  $R_{e-g} R_{(h)} \subseteq R_{(gh)}$  for all  $g, h \in G$  because  $R_{e-g} R_{\sigma^{-1}h} \subseteq R_{\sigma^{-1}gh} \subseteq R_{(gh)}$  for all  $\sigma \in G$ . Clearly  $R = \bigoplus_{g \in G} R_{(g)}$ .

$$\text{In Example 2.10, } R[G]_{(g)} = \bigoplus_{h \in G} R[G]_{h-g} = \bigoplus_{h \in G} R_g.h.$$

$$\text{In Example 2.11, } \overline{R[G]}_{(g)} = \bigoplus_{h \in G} \overline{R[G]}_{h-g} = \bigoplus_{h \in G} R_g g^{-1}h.$$

**Definition 2.13 ([10]).** Let  $R$  be a ring with two gradations  $(R, G)$  and  $(R, H)$ . Then  $(R, G)$  is almost equivalent to  $(R, H)$  if there exists a ring isomorphism  $f : R \rightarrow R$  such that for each  $h \in H$  there exists  $g \in G$  with  $f(R_g) = R_h$ .

In [8, 10] we proved that first strong, second strong and nondegenerate are preserved between almost equivalent gradations, but the faithfulness and strong properties are not. We finish this paper by giving similar result on the augmented graded rings.



**Proposition 2.14 .** *Let  $R$  be an augmented  $G$ -graded ring such that  $\text{supp}(R, G) = \text{supp}(R_{e_G}, G)$  and  $R_{g_1} R_{g_2} \neq 0$  for all  $g_1, g_2 \in \text{supp}(R, G)$ . Suppose  $(R, G)$  is almost equivalent to  $(R, H)$ . Then  $R$  is an augmented  $H$ -graded ring with the same graduation of  $R$  by  $H$  and with a graduation on  $R_{e_H}$  defined as follows :*

$$R_{e_H} - h = \begin{cases} f(R_{e_G} - g) & \text{if } h \in \text{supp}(R, H) \text{ where } f(R_g) = R_h \\ O & \text{if } h \notin \text{supp}(R, H) \end{cases}$$

$$\text{Let } r_h = \begin{cases} f(r_g) & \text{if } h \in \text{supp}(R, G) \text{ and } f(R_g) = R_h \\ O & \text{otherwise} \end{cases}$$

**Proof.** First , we show that  $R_{e_H}$  is an  $H$ -graded ring. Since  $f$  is an isomorphism ,  $R_{e_H-h}$  is an additive subgroup of  $R_{e_H}$ . Let  $h_1, h_2 \in \text{supp}(R, H)$ . Then  $R_{e_H-h_1} R_{e_H-h_2} = f(R_{e_G-g_1})f(R_{e_G-g_2})$  where  $R_{h_1} = f(R_{g_1})$  and  $R_{h_2} = f(R_{g_2})$ .

Since  $g_1, g_2 \in \text{supp}(R, G)$ ,  $R_{g_1} R_{g_2} \neq 0$  and hence  $O \neq f(R_{g_1} R_{g_2}) \subseteq R_{h_1 h_2}$ . So ,  $f(R_{g_1 g_2}) = R_{h_1 h_2}$  (See Lemma 2.2 of [10]). Therefore ,  $R_{e_H-h_1} R_{e_H-h_2} \subseteq f(R_{e_G-g_1 g_2}) = R_{e_H-h_1 h_2}$ . Clearly ,  $R_{e_H} = f(R_{e_G}) = \bigoplus_{g \in G} f(R_{e_G-g}) = \bigoplus_{h \in H} R_{e_H-h}$  ,

*i.e.*,  $R_{e_H}$  is an  $H$ -graded ring.

Let  $h \in \text{supp}(R, H)$ . Then  $R_h = f(R_g)$  for some  $g \in \text{supp}(R, G)$ . But  $f(R_g) = \bigoplus_{\sigma \in G} f(R_{e_G-\sigma} r_g) = \bigoplus_{\sigma \in G} f(R_{e_G-\sigma})f(r_g) = \bigoplus_{\tau \in H} R_{e_H-\tau} r_h$ .

Suppose  $h_1, h_2 \in \text{supp}(R, H)$ . Then  $r_{h_1} = f(r_{g_1})$  and  $r_{h_2} = f(r_{g_2})$  where  $f(R_{g_1}) = R_{h_1}$  and  $f(R_{g_2}) = R_{h_2}$ . Now ,  $r_{h_1} r_{h_2} = f(r_{g_1} r_{g_2}) = f(r_{g_1 g_2}) \neq 0$ . But  $f(R_{g_1 g_2}) = R_{h_1 h_2}$  implies  $f(r_{g_1 g_2}) = r_{h_1 h_2}$ , *i.e.*,  $r_{h_1} r_{h_2} = r_{h_1 h_2}$ .

Finally, let  $x, y \in R_{e_H}$ . Then  $(x r_{h_1})(y r_{h_2}) = f(x') f(r_{g_1}) f(y') f(r_{g_2})$  where  $f(x') = x$  and  $f(y') = y$ . Hence  $f(x') f(r_{g_1}) f(y') f(r_{g_2}) = f(x' r_{g_1} y' r_{g_2}) = f(x' y' r_{g_1 g_2}) = f(x') f(y') f(r_{g_1 g_2}) = x y r_{h_1 h_2}$ .  $\square$

### References

- [1] M.Cohen and L.Rowen, Group graded rings, Comm. in Algebra 11 (11) , 1253-1270 (1983).
- [2] E.C.Dade , Compounding Clifford's Theory, Ann. of Math. 19 , 236-290 (1970).
- [3] E.C.Dade , Group-graded rings and modules, Math. Z. 174, 241 - 262 (1980).
- [4] C.Năstăsescu , Group rings of graded rings. Applications, J. of pure and applied algebra 33 , 313-335 (1984).

## REFAI

- [5] C.Năstăsescu and F. Van Oystaeyn , Graded Ring Theory , Mathematical Library 28 , North Holland , Amsterdam (1982).
- [6] M.Refai , Modules over  $K[x]$  , J. Institute of Mathematics & Computer Sciences (Math.Series) 5 (2) , 223-229 (1992) .
- [7] M.Refai, Totally finite DG-A -modules , M.E.T.U. Journal of Pure and Applied Sciences 24 (1-2) , (1991).
- [8] M.Refai , Various types of strongly G-graded rings , Alexandria Engineering Journal 32 (2) , 91-95 (1993) .
- [9] M.Refai and M.Obiedat, On a strongly-support graded rings , Math. Japonica 39 (3) , 1-5 (1994).
- [10] M.Refai and M.Obiedat, On graduations of  $K[x_1, x_2, \dots, x_m]$  J.Institute of Mathematics & Computer Sciences (Math. Series) , 6 (3) , 241 - 252 (1993) .
- [11] M. Refai, On Noetherian modules graded by G.sets, Acta Mathematica Hungarica Journal, 69 (3), 211-219 (1995).

### Genişletilmiş Katmanlı Halkalar

#### Özet

Bu çalışmada genişletilmiş katmanlı halkalar incelenmiş ve kuvvetli katmanlı halkalarla olan ilişkileri ortaya konmuştur.

Mashhoor REFAI  
Department of Mathematics  
Yarmouk University  
Irbid-JORDAN

Received 3.4.1996