

1-1-1997

Exotic structures and adjunction inequality

Selman AKBULUT

Rostislav MATVEYEV

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

AKBULUT, Selman and MATVEYEV, Rostislav (1997) "Exotic structures and adjunction inequality," *Turkish Journal of Mathematics*: Vol. 21: No. 1, Article 4. Available at: <https://journals.tubitak.gov.tr/math/vol21/iss1/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Exotic structures and adjunction inequality

Selman Akbulut and Rostislav Matveyev

1. Introduction

Here we want to reprove and strengthen some old difficult theorems of 4 manifolds by the aid of recently proven modern tools. One of the important recent results of smooth 4-manifolds is Eliashberg's topological description of compact Stein manifolds, that is complex manifolds which admit strictly plurisubharmonic Morse function (PC-manifolds):

Theorem 1.1. ([E]) *Let $X = B^4 \cup (1\text{-handles}) \cup (2\text{-handles})$ be four-dimensional handlebody with one 0-handle and no 3- or 4-handles. Then*

- *The standard PC-structure on B^4 can be extended over 1-handles so that manifold $X_1 = B^4 \cup (1\text{-handles})$ has pseudo-convex boundary.*
- *If each 2-handle is attached to ∂X_1 along a Legendrian knot with framing one less than Thurston-Bennequin framing of this knot then the symplectic form and complex structure on X_1 can be extended over 2-handles to a symplectic form on X , which makes X a PC manifold.*

Lisca and Matić showed that PC manifolds imbed naturally into Kähler surfaces:

Theorem 1.2. ([LM]) *Every PC manifold X can be holomorphically embedded as a domain into a minimal Kähler surface S with ample canonical bundle and $b_2^+(S) > 1$, such that the induced Kähler form on X agrees with the symplectic form of the PC structure.*

Theorem 1.3. *A minimal Kähler surface X , with $b_2^+(X) > 1$ and an ample canonical bundle, can not contain a smoothly embedded 2-sphere $\Sigma \subset X$ with $\Sigma.\Sigma \geq -1$.*

This theorem roughly follows from the fact that Kähler surfaces have nonzero Seiberg-Witten invariants (see [B] and [MF] in case $\Sigma.\Sigma = -1$, and [FS] in case $\Sigma.\Sigma \geq 0$). Also Kähler surfaces satisfy the adjunction inequality of Kronheimer and Mrowka:

Theorem 1.4. ([KM1], [MST]) *Let X be a closed smooth 4-manifold with $b_2^+(X) > 1$, with a nonzero Seiberg Witten invariant (e.g. X Kähler surface) corresponding to the line bundle $L \rightarrow X$. Let $\Sigma \subset X$ be a compact oriented embedded surface with $\Sigma.\Sigma \geq 0$ when Σ is not a sphere, then*

$$2g(\Sigma) - 2 \geq \Sigma.\Sigma + |c_1(L).\Sigma|$$

The first author was partially supported by an NSF grant DMS-9626204

After discussing the basics, we will show some 4-manifold theorems can be obtained as easy corollaries of these three basic theorems.

2. Definitions

Let us recall some basic facts (e.g. [G], [E]). Any PC-manifold X induces a contact structure on the boundary 3-manifold $Y = \partial X$, which is isomorphic to the restriction of the dual K^* of the canonical line bundle $K \rightarrow X$. Furthermore if α is an oriented Legendrian knot in Y (a knot whose tangents lie in the contact planes) bounding an oriented surface F in X , then the “rotational number” is defined to be the relative Chern class $rot(\alpha, F) = c_1(K^*, v)$ of the induced 2-plane bundle $K^* \rightarrow F$ with respect to the tangent vector field v of α (i.e. the obstruction to extending v to a section of K^* over F). Also this contact structure gives the so called Thurston-Bennequin framing $tb(\alpha)$.

The simplest example of a PC-manifold is $B^4 \subset \mathbf{C}^2$ with the induced symplectic structure. We choose coordinates in $\mathbf{R}^3 \subset S^3 = \partial B^4$, so that the induced contact structure ξ_0 on \mathbf{R}^3 is the kernel of the form $\lambda_0 = dz + xdy$. By Theorem 1.1 this PC-structure extends across 1-handles attached to B^4 ; we draw the attaching balls of each 1-handle on the plane $\{x = 0, z = \text{constant}\}$. Any link L in S^3 can be isotoped to a Legendrian link, we can achieve this by first isotoping L so that all crossings are left handed as in the first picture of Figure 1. For example, we can turn a right handed crossing to a left handed crossing by the local isotopy as in the second picture of Figure 1.




FIGURE 1

Projection of a Legendrian link to the yz -plane has intersections only coming from left handed crossing, no vertical tangencies and all minima and maxima in y -direction are cusps (as the projection of the knot in Figure 3 to the yz -plane). Moreover, every projection with these properties is a projection of some Legendrian link. Rotational number of an oriented Legendrian knot α in S^3 does not depend the surface $F \subset S^3$ it bounds. Invariants $rot(K)$ and $tb(K)$ can be calculated by:

$$rot(\alpha) = 1/2(\text{Number of “downward” cusps} - \text{Number of “upward” cusps})$$

$$tb(\alpha) = bb(\alpha) - c(\alpha)$$

where $bb(\alpha)$ is the blackboard (yz -plane) framing of the projection of α , $c(\alpha)$ is the number of right cusps, and “downward” and “upward” cusps are calculated in the obvious way. Because the local isotopy in Figure 1 introduces one right cusp (hence a -1 contribution to calculation of $tb(K)$) we can incorporate this to “self crossing number” calculation by reading the crossings numbers in a modified way as in in Figure 2. For example in the

knot of Figure 3 we can calculate $tb(K) = 5 - 4 = 1$ and $rot(K) = 0$. A useful corollary to these theorems is the following generalization of the Bennequin inequality:




FIGURE 2




FIGURE 3

Corollary 2.1. *Let X be a PC manifold, $F \subset X$ be a two-dimensional submanifold of X , such that $\alpha = \partial F \subset \partial X$ is Legendrian with respect to induced contact structure and f is framing on α induced by a trivialization of the normal bundle of F in X , then*

$$-\chi(F) \geq [tb(\alpha) - f] + |rot(\alpha)|$$

where $tb(\alpha)$ and $rot(\alpha)$ are the Thurston-Bennequin framing and rotational number of α

To prove this we attach a 2-handle to X along α with the framing $tb(\alpha) - 1$, and apply Theorems 1.1, 1.2, and 1.3 to the resulting manifold and closed surface $F' = F \cup_{\partial} D$, where D is the core 2-disc of the 2-handle.

3. Applications

Theorem 3.1. ([A1]) *Let W be the contractible manifold of Figure 4. Let $f : \partial W \rightarrow \partial W$, be the involution induced by an involution of S^3 (as described in [A1]) with $f(\gamma) = \gamma'$, where γ and γ' are circles in ∂W shown on Figure 4. Then $f : \partial W \rightarrow \partial W$ does not extend to a diffeomorphism $F : W \rightarrow W$ (but it extends to a homeomorphism)*

Proof. By applying Theorem 1.1 to W (the second picture of W in Figure 4) we see that W has a PC structure. Also since γ is slice in W if f extended to a diffeomorphism $F : W \rightarrow W$, then γ' would be slice also. But this contradicts the inequality of Theorem 2.1 (here $F = D^2$, $f = 0$, and $tb(\gamma') = 0$). \square




FIGURE 4

Theorem 3.2. ([A2]) *Let Q_1 and Q_2 be the manifolds obtained by attaching 2-handles to B^4 along the knots K_1 and K_2 with -1 framings as in Figure 5, then Q_1 and Q_2 are homeomorphic but not diffeomorphic to each other, even interiors are not diffeomorphic to each other.*




FIGURE 5

Proof. By Theorem 1.1 Q_1 has a PC structure, by Theorem 1.2 Q_1 is a domain in a minimal Kähler surface S with $b_2^+(S) > 1$. If Q_1 were diffeomorphic to Q_2 , the generator of $H_2(Q_1; \mathbf{Z})$ would be represented by a smooth embedded sphere (since K_2 is a slice knot the generator of $H_2(Q_1; \mathbf{Z})$ represented by a smooth sphere) with -1 self intersection, violating Theorem 1.3. \square

Let $K \subset S^3$ be a Legendrian knot, and K'_0 be a 0-push off of K (the zero framing is the framing induced from the normal vector field of K in the oriented surface in S^3 bounding K).

Proposition 3.3. *We can move K'_0 to a Legendrian knot K_0 by an isotopy which fixes K , such that $tb(K_0) = -|tb(K)|$.*

Proof. $tb(K) = bb(K) - c(K)$. If $bb(K) \leq 0$ then K'_0 is just the blackboard push-off of K with $-2bb(K)$ right half twist, but this is a projection of a Legendrian link and $bb(K_0) = bb(K)$ and $c(K_0) = c(K)$ hence $tb(K_0) = tb(K)$ If $tb(K) \leq 0$ and $bb(K) > 0$,




FIGURE 6

then K'_0 is the blackboard push-off with $2bb(K)$ left handed half twist. Knot K has $2c(K)$ cusps, and $2c(K) \geq 2bb(K)$. We can produce a Legendrian picture by placing all half twists near cusps as in Figure 7. Thus $tb(K_0) = tb(K)$.




FIGURE 7

If $tb(K) > 0$, then $bb(K) > 0$ and again K'_0 is the blackboard push-off with $2bb(K)$ left half twist. By above trick of placing half left twist near cusps, we can get rid of $2c(K)$ left twist, and we are left with $2bb(K) - 2c(K)$ left twist each of which contributes -1 to $tb(K_0)$ as in Figure 8. Hence $tb(K_0) = tb(K) - [2bb(K) - 2c(K)] = tb(K) - 2tb(K) = -tb(K)$




FIGURE 8

□

By Proposition 3.3 we can produce many 0-push offs K_0^i , $i = 1, 2, \dots, k$ such that $tb(K_0^i) = -|tb(K)|$. So if $tb(K) \leq 0$ then all K_0^i have same tb , if $tb(K) > 0$ then all K_0^i but the original knot K has the same tb .

Theorem 3.4. *If $K \subset S^3$ a Legendrian knot with $tb(K) \geq 0$, then all iterated positive Whitehead doubles of $Wh_n(K)$ are not slice. In fact if $Q_n^r(K)$ is the manifold obtained by attaching a 2-handle to B^4 along $Wh_n(K)$ with a framing $-1 \leq r \leq 0$, then there is no smoothly embedded 2-sphere in $Q_n^r(K)$ representing the generator of $H_2(Q_n^r(K); \mathbf{Z})$.*

Proof. The first positive Whitehead double is obtained by connecting K and K_0 by a left handed cusp as in Figure 9 which contributes $+1$ to tb , hence

$$tb(Wh(K)) = tb(K) + tb(K_0) + 1 = tb(K) - tb(K) + 1 = 1.$$

So by iteration we get $tb(Wh_n(K)) = 1$. But Corollary 2.1 says that any slice knot L must have $tb(L) \leq -1$.




FIGURE 9

For the second part, observe that since $r \leq 0 = tb(Wh_n(K)) - 1$, by Theorem 1.1 $Q_n^r(K)$ is PC. By Theorem 1.2 we can imbed $Q_n^r(K)$ into Kähler surface S . Then by Theorem 1.3 we can not have a smoothly embedded 2-sphere $\Sigma \subset Q_n^r(K) \subset S$ representing $H_2(Q_n^r(K); \mathbf{Z})$, since $\Sigma \cdot \Sigma = r \geq -1$. \square

Remark 3.1. L.Rudolf has previously shown that $Wh_n(K)$ are not slice if $tb(K) \geq 0$ ([R]).

Remark 3.2. Proof of Proposition 3.2 gives: If K'_r is the r -framing push off of a knot K , then K'_r can be isotoped to a Legendrian knot K_r fixing K with $(tb(K_r) - r) = -|tb(K) - r|$

Remark 3.3. If ξ is the contact structure on $\Sigma = \partial W$ induced by the PC manifold W of Theorem 3.1, and $f^*(\xi)$ is the “pull-back” contact structure on Σ , then it follows that the contact structures ξ and $f^*(\xi)$ are homotopic through 2-plane fields but not isotopic through contact structures ([AM]).

References

- [A1] S. Akbulut, A Fake compact contractible 4-manifold. *J. Diff. Geom* Vol. 33 (1991) pp. 335-356.
- [A2] S. Akbulut, An exotic 4-manifold *J. Diff. Geom* Vol. 33 (1991) pp. 357-361.
- [AM] S. Akbulut and R. Matveyev, A note on Contact Structures (to appear Pac. Jour. of Math).

AKBULUT & MATVEYEV

- [B] R. Brussee, The Kähler class and the C^∞ properties of Kähler surfaces, New York Journal of Mathematics volume 2 1996 (also in alg-geom 9503004).
- [G] R.Gompf, Handlebody construction of Stein surfaces (preprint).
- [E] Y. Eliashberg, Topological characterization of Stein manifolds of dimension > 2 . *International J. of Math.* Vol. 1, No 1 (1990) pp.29-46.
- [FS] R. Fintushel and R. Stern, Immersed spheres in 4-manifolds and the immersed Thom Conjecture, Proceedings of the 3rd Gokova Geometry-Topology Conference, 27-39 (1994).
- [KM1] P. Kronheimer and T. Mrowka, On genus of embedded surfaces in the projective plane, Math. Res. Letters 1 (1994), 797-808.
- [KM2] P. Kronheimer and T. Mrowka, Monopoles and contact structures (to appear).
- [LM] P. Lisca and G. Matic, Tight contact structures and Seiberg Witten invariants.
- [MF] J. Morgan and R. Friedman, Algebraic surfaces and Seiberg-Witten invariants (preprint).
- [MST] J. Morgan, Z.Szabo, C.H. Taubes, The generalized Thom conjecture (to appear).
- [R] L. Rudolf, An obstruction to sliceness via contact geometry and “classical” gauge theory, Invent.math. 119, 155-163 (1995).

MICHIGAN STATE UNIVERSITY, DEPT OF MATHEMATICS, E. LANSING MI 48824, USA
E-mail address: akbulut@math.msu.edu, matveyev@math.msu.edu