

1-1-1996

ON THE $\ell_{\{P\}}$ NORMS OF ALMOST CAUCHY-TOEPLITZ MATRICES

D. BOZKURT

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

BOZKURT, D. (1996) "ON THE $\ell_{\{P\}}$ NORMS OF ALMOST CAUCHY-TOEPLITZ MATRICES," *Turkish Journal of Mathematics*: Vol. 20: No. 4, Article 10. Available at: <https://journals.tubitak.gov.tr/math/vol20/iss4/10>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

ON THE ℓ_p NORMS OF ALMOST CAUCHY-TOEPLITZ MATRICES

D. Bozkurt

Abstract

In this study, we have given the definition of almost Cauchy-Toeplitz matrix. i.e. its elements are $t_{ij} = a(i = j)$ and $t_{ij} = 1/(i - j)$ ($i \neq j$) such that a is a real number. We have found a lower and upper bounds for the ℓ_p norm of this matrix. Furthermore, we have done the proof of the conjecture that were given by myself for the spectral norm of this matrix.

Key Words: Cauchy, Toeplitz, norm, Cauchy-Toeplitz matrix, matrix norm.

1. Introduction

In this study, we have given the definition of almost Cauchy-Toeplitz matrix (i.e. its elements are $t_{ij} = a(i = j)$ and $t_{ij} = 1/(i - j)$ such that a is a real number).

It is called Cauchy and Toeplitz matrices as in the following forms:

$$C_n = [1/(x_i - y_j)]_{i,j=1}^n$$

where $x_i \neq y_j$ for all i, j and

$$T_n = [t_{j-i}]_{i,j=1}^n$$

respectively. It is easy to check that an arbitrary Cauchy- Toeplitz matrix of order n is of the form

$$T_n = \left[\frac{1}{g + (i - j) \cdot h} \right]_{i,j=1}^n \quad (1.1)$$

where g and h are some numbers. We assume that $h \neq 0$ and the quotient $g/h \notin Z$ (Z is the set of integers).

In [5], E. E. Tyrtyshnikov found a lower bound for the spectral norm of Cauchy-Toeplitz matrix such that $h = 1$ and $g = 1/2$. He had computed that with $\varepsilon = 10^{-4}$ for $n = 40, 60, 100$ then numbers of singular values $\rho_{1n} \geq \rho_{2n} \geq \dots \geq \rho_{nn}$ of the matrix T_n satisfying the condition $\rho_{jn} > \pi - \varepsilon$ are equal to 6,7,8 respectively.

A few years ago these matrices attracted the attention of C. Moler, who had experimentally discovered that most of their singular values are clustered near π . Recently S. Parter gave an explanation of this phenomenon [3].

In [1], we had generalized the bounds that E. E. Tyrtyshnikov had found. i. e. we had established that

$$\frac{\sqrt{n}\pi}{|h|} - s_n \leq \| T_n \| \leq \frac{\sqrt{n}\pi}{|h|} \tag{1.2}$$

where $h \neq 0, g/h$ is not integer, $s_n > 0$ and $s_n = O(1/n)$.

In Section 2 we have given the definition of almost Cauchy-Toeplitz matrix. We have found a lower and an upper bounds for the ℓ_p norm of this matrix. Furthermore, we have done the proof of the conjecture that we had given in [2] for the spectral norm of this matrix.

2. The ℓ_p Norms of the almost Cauchy-Toeplitz Matrices

Let A be any nxn matrix. The ℓ_p norm of the matrix A is

$$\| A \|_p = \left[\sum_{i,j=1}^n |a_{ij}|^p \right]^{1/p} \tag{2.1}$$

where $p \geq 1$. We can write the equation (2.1) in the following form:

$$\| A \|_p = \left[\sum_{i,j=1}^n |a_{ij}|^p \right]^{1/p} = \left[\sum_{i=1}^n (\|_{1i} e_i \|^p + \| a_{2i} e_i \|^p + \dots + \| a_{ni} e_i \|^p) \right]^{1/p} \tag{2.2}$$

where e_i ($1 \leq i \leq n$) are the standard basis of R^n .

Definition *The matrix A_n is called almost Cauchy-Toeplitz matrix if*

$$A_n = \begin{cases} a, i = j \\ \frac{1}{i-j}, i \neq j \end{cases} \tag{2.3}$$

where $a \in R$ (R is the set of the real numbers) and $1 \leq i, j \leq n$.

In [2] we had shown that the following theorem is valid $p > 2$.

Theorem 2.1.[2] *Let the matrix A_n be as in (2.3). Then*

$$\sqrt[p]{|a|^p - \alpha_n} \leq n^{-1/p} \| A_n \|_p \leq \sqrt[p]{|a|^p + 2\zeta(p)}$$

is valid where $\zeta(p)$ is Riemann's zeta function, $a_n > 0, \alpha_n = O(1/n)$ and $p > 2$.

Proof. From (2.2), we have

$$\| A \|_p = \left[\sum_{i,j=1}^n |a_{ij}|^p \right]^{1/p} = \left[\sum_{i=1}^n (|a_{1i}|^p \| e_i \|^p + |a_{2i}|^p \| e_i \|^p + \dots + |a_{ni}|^p \| e_i \|^p) \right]^{1/p}.$$

Since e_i are the standard basis of R^n for all $i (1 \leq i \leq n)$, its norms are equal to 1. Then, we obtain

$$\begin{aligned} \| A_n \|_p &= \left\{ n|a|^p + 2 \left[(n-1) + (n-2) \left(\frac{1}{2}\right)^p + (n-3) \left(\frac{1}{3}\right)^p + \dots + \left(\frac{1}{n-1}\right)^p \right] \right\}^{1/p} \\ &= \left(n|a|^p + 2 \sum_{k=1}^{n-1} \frac{n-k}{k^p} \right)^{1/p} \end{aligned}$$

Hence

$$\| A_n \|_p = \left(n|a|^p + 2n \sum_{k=1}^{n-1} \frac{1}{k^p} - 2 \sum_{k=1}^{n-1} \frac{1}{k^{p-1}} \right)^{1/p}. \tag{2.4}$$

Multiplying the equation (2.4) with $n^{-1/p}$, then

$$n^{-1/p} \| A_n \|_p = \left(|a|^p + 2 \sum_{k=1}^{n-1} \frac{1}{k^p} - \frac{2}{n} \sum_{k=1}^{n-1} \frac{1}{k^{p-1}} \right)^{1/p}. \tag{2.5}$$

From (2.5)

$$\lim_{n \rightarrow \infty} n^{-1/p} \| A_n \|_p = \lim_{n \rightarrow \infty} \left(|a|^p + 2 \sum_{k=1}^{n-1} \frac{1}{k^p} - \frac{2}{n} \sum_{k=1}^{n-1} \frac{1}{k^{p-1}} \right)^{1/p}.$$

Since

$$\sum_{k=1}^{\infty} \frac{1}{k^p} = \zeta(p) \text{ ve } \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^{n-1} \frac{1}{k^{p-1}} = 0,$$

then we obtain

$$n^{-1/p} \| A_n \|_p \leq \sqrt[p]{|a|^p + 2\zeta(p)}. \tag{2.6}$$

from (2.5). Thus , we have found upper bound for A_n .

From (2.5) we have

$$n^{-1/p} \| A_n \|_p \geq \left(|a|^p - \frac{2}{n} \sum_{k=1}^{n-1} \frac{1}{k^{p-1}} \right)^{1/p} . \tag{2.7}$$

If we get

$$\alpha_n = \frac{2}{n} \sum_{k=1}^{n-1} \frac{1}{k^{p-1}}$$

in (2.7), $\alpha_n > 0$ and

$$\lim_{n \rightarrow \infty} \frac{\alpha_n}{1/n} \leq 2\zeta(p-1).$$

Then $\alpha_n = O(1/n)$. Hence

$$n^{-1/p} \| A_n \|_p \geq \sqrt[p]{|a|^p - \alpha_n} \tag{2.8}$$

Thus, the proof of the theorem is completed from (2.6) and (2.8). □

Theorem 2.2. *Let the matrix A_n be as in (2.3). Then*

$$\begin{aligned} |a| + o(n) \leq n^{-1} \| A_n \|_p &\leq |a| + \gamma + \log n + O(1/n), \quad (p = 1) \\ |a| \leq n^{-1/p} \| A_n \|_p &\leq \sqrt[p]{|a|^p + \frac{2^p}{2^{p-1} - 1}}, \quad (1 < p \leq 2) \\ \sqrt[p]{|a|^p + \zeta(p)} \leq n^{-1/p} \| A_n \|_p &\leq \sqrt[p]{|a|^p + 2\zeta(p)} \end{aligned} \tag{2.9}$$

are valid where $\zeta(p)$ is Riemann's zeta-function and γ is Euler's constant.

The lower bound in (2.9) is better than the lower bound in Theorem 2.1.

Proof. Let us do three steps the proof of theorem.

Step 1. Let $p = 1$. From (2.5) we obtained

$$\left. \begin{aligned} n^{-1} \| A_n \|_1 &= |a| + 2 \sum_{k=1}^{n-1} \frac{1}{k} - \frac{2(n-1)}{n} \\ &= |a| + 2 \sum_{k=1}^{n-1} \frac{1}{k} - 2 + \frac{2}{n} \\ &\leq |a| + 2 \sum_{k=1}^{n-1} \frac{1}{k} + \frac{2}{n} \end{aligned} \right\} . \tag{2.10}$$

Since $\frac{2}{n} \rightarrow 0$ and $\sum_{k=1}^n \frac{1}{k} = \gamma + \log n + O(1/n)$ for the sufficient large n , then

$$n^{-1} \| A_n \|_1 \leq |a| + \gamma + \log n + O(1/n). \tag{2.11}$$

We have

$$n^{-1} \| A_n \|_1 \geq |a| + \sum_{k=2}^n \frac{1}{k}$$

from the first equation of (2.10). Taking

$$\alpha_n = \sum_{k=2}^n \frac{1}{k},$$

then $\alpha_n = o(n)$ (i.e. $\lim_{n \rightarrow \infty} \alpha_n/n = 0$). We have

$$n^{-1} \| A_n \|_1 \geq |a| + o(n) \tag{2.12}$$

Step 2. Let $1 < p \leq 2$. We obtain

$$n^{-1/p} \| A_n \|_p \leq \left(|a|^p + 2 \sum_{k=1}^{n-1} \frac{1}{k^p} \right)^{1/p}$$

from (2.5). Since

$$\sum_{k=1}^{\infty} \frac{1}{k^p} \leq \frac{2^{p-1}}{2^{p-1} - 1}$$

according to p-test for the series, we find

$$n^{-1/p} \| A_n \|_p \leq \sqrt[p]{|a|^p + \frac{2^p}{2^{p-1} - 1}}. \tag{2.13}$$

From (2.5) we have

$$n^{-1/p} \| A_n \|_p \geq \left(|a|^p - \frac{2}{n} \sum_{k=1}^{n-1} \frac{1}{k^{p-1}} \right)^{1/p}.$$

Then since

$$\lim_{n \rightarrow \infty} \frac{2}{n} \sum_{k=1}^{n-1} \frac{1}{k^{p-1}} = 0,$$

we obtain

$$n^{-1/p} \| A_n \|_p \geq |a|. \tag{2.14}$$

Step 3. Let $p > 2$. Since the proof of the right side of the inequality (2.9) was done Theorem 2.1, here we will do the proof of the left side of the inequality (2.9). We have

$$2 \sum_{k=1}^{n-1} \frac{1}{k^p} - \frac{2}{n} \sum_{k=1}^{n-1} \frac{1}{k^{p-1}} \geq \sum_{k=1}^{n-1} \frac{1}{k^p}$$

from the equation (2.5) as $n \rightarrow \infty$. Then

$$n^{-1/p} \| A_n \|_p \geq \left(|a|^p + \sum_{k=1}^{n-1} \frac{1}{k^p} \right)^{1/p}.$$

Since

$$\sum_{k=1}^{\infty} \frac{1}{k^p} = \zeta(p),$$

we have

$$n^{-1/p} \| A_n \|_p \geq \sqrt[p]{|a|^p + \zeta(p)}. \tag{2.15}$$

The proof of the Theorem 2.2 is completed from (2.11), (2.12), (2.13), (2.14) and (2.15). \square

Conjecture 2.1[2] *Let the matrix A_n be as in (2.3). Then*

$$\| A_n \|_2 \leq \sqrt{a^2 + \pi^2}$$

where $\| \cdot \|_2$ is the spectral norm of the matrix A_n .

Proof. Let $H_n = H_n^* = i(A_n - aI)$ with spectral radius $r(H_n)$ where I is the unit matrix of order n . Then A_n is normal matrix with spectrum

$$\sigma(A_n) = \{a + i\lambda : \lambda \in \sigma(H_n)\},$$

and hence

$$\| A_n \|_2 = r(A_n) = \sqrt{a^2 + r(H_n)^2}.$$

Let us consider the symmetric matrix

BOZKURT

$$\tilde{H}_n = \frac{1}{i} \begin{bmatrix} 0 & H_n^T \\ H_n & 0 \end{bmatrix}$$

The matrix \tilde{H}_n is permutation-similar to the block Toeplitz matrix \hat{H}_n as in the following form:

$$\tilde{H}_n = \begin{bmatrix} 0 & 0 & -1 & 0 & -\frac{1}{2} & 0 & \dots \\ 0 & 0 & 0 & 1 & 0 & \frac{1}{2} & \dots \\ 1 & 0 & 0 & 0 & -1 & 0 & \dots \\ 0 & -1 & 0 & 0 & 0 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

After all the odd rows have been multiplied by the imaginary unit i and the even ones by $-i$, the result is Hermitian Toeplitz matrix $T_n = [t_{i-j}]_{i,j=0}^{2n-1}$. Let $f(x)$ be a real-valued function and $u = [u_0, u_1, \dots, u_{2n-1}]^T$. Then we have

$$(T_n u, u) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \left| u_0 + u_1 e^{ix} + \dots + u_{2n-1} e^{i(2n-1)x} \right|^2 f(x) dx,$$

$$(u, u) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \left| u_0 + u_1 e^{ix} + \dots + u_{2n-1} e^{i(2n-1)x} \right|^2 dx$$

[4, pp. 63] and

$$\| H_n \|_2 = \| T_n \|_2 = \max_{u \neq 0} \left| \frac{(T_n u, u)}{(u, u)} \right| \leq \pi \tag{2.16}$$

[5, pp. 9 and 3, pp. 120]. Let $T_n(f)$ be a Hermitian Toeplitz form. If a and b are arbitrary real numbers, the eigenvalue associated with the function $a + bf(x)$ will be $a + b\lambda$ [4, pp. 64]. So, we have

$$\| A_n \|_2 = r(A_n) \leq \sqrt{a^2 + \pi^2}.$$

Thus, the proof of the conjecture is completed. □

BOZKURT

References

- [1] Bozkurt, D.: On the Bounds of Cauchy-Toeplitz Matrices. Siberian Journal of Differential Equations, Volume: 3, No: 2, pp. 135-142 (1996).
- [2] Bozkurt, D.: Almost Cauchy-Toeplitz Matrislerinin ℓ_p Normları Üzerine. VIII. Ulusal Matematik Sempozyumu, (1995).
- [3] Parter, S. V.: On the Distribution of the Singular Values of Toeplitz Matrices. Linear Algebra and Its Appl. 80:115-130 (1986).
- [4] Szegő, G. and Grenander, U.: Toeplitz Forms and Their applications. University of California Press (1958).
- [5] Tyrthyhnikov, E. E.: Cauchy-Toeplitz Matrices and Some Applications. Linear Algebra and Its Appl. 149:1-18 (1991).

ALMOST CAUCHY-TOEPLITZ MATRİSLERİNİN ℓ_p NORMLARI

Özet

Bu çalışmada, Almost Cauchy-Toeplitz matris tanımını yaptık (yani a herhangi bir reel sayı olmak üzere elemanları $t_{ij} = a(i = j)$ ve $t_{ij} = 1/(i - j)(i \neq j)$ şeklinde olan matris). Bu matrisin ℓ_p normu için bir alt ve üst sınır bulduk. Ayrıca bu matrisin spektral normuyla ilgili bir konjektürün ispatını yaptık.

Durmuş BOZKURT
Department of Mathematics Faculty of Art and
Science of Selçuk University,
42031, Kampus, Konya-TURKEY
e-mail: dbozkurt@karatay.cc.selcuk.edu.tr

Received 26.1.1996