

1-1-1996

EXTENSION OF CARISTI-KIRK'S THEOREM

M. O. DIALLO

M. OUDADESS

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

DIALLO, M. O. and OUDADESS, M. (1996) "EXTENSION OF CARISTI-KIRK'S THEOREM," *Turkish Journal of Mathematics*: Vol. 20: No. 2, Article 2. Available at: <https://journals.tubitak.gov.tr/math/vol20/iss2/2>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

EXTENSIONS OF CARISTI- KIRK'S THEOREM

M.O. Diallo, M. Oudadess

Abstract

We give some extensions and/or improvements, to uniform spaces and to multi-valued mappings, of Caristi-Kirk's theorem.

Key words and phrases: Uniform spaces, multi-valued mappings, fixed point theorem, maximal element, weak p -contraction mappings.

1. Introduction

It was observed that certain fixed point theorems can be deduced from the following result:

Let (E, \leq) be an ordered set which admits a maximal element.

Let $f : E \rightarrow E$ be a mapping such that $x \leq f(x)$ for every x in E .

Then f has a fixed point.

This result served as a basis for certain theorems about the existence of maximal elements ([1], [2], [3], [6]), and hence fixed point theorems. Considered spaces are often metric spaces endowed with an order defined via the distance ([2], [3], [4], [5]).

Ekland's variational principle, which concerns the existence of maximal elements ([2], [6]) and its generalizations allowed simple proof of Caristi-Kirk's theorem ([2],[3],[8]).

Recently, V. Conserva ([5]) gave a slight improvement of this theorem in metric spaces.

In this paper we give some extensions and/or improvements to uniform spaces and to multi-valued mappings, of Caristi- Kirk's theorem.

Let us notice that proofs we give here go along the lines of those given in the case of metric or topological vector spaces ([2], [3], [4], [5]).

In the following, for a uniform space E , we consider a family $(d_i)_{i \in I}$ of semi-metrics which defines its uniform structure and such that $\sup_{i \in I} d_i(x, y) < +\infty$, for all x, y in E .

Let E be a uniform space and $p : E \rightarrow \mathbb{R}_+$ a positive real functional on E . Define a partial order on E as follows:

$x \leq y$ if and only if $d_i(x, y) \leq p(x) - p(y)$, for all $i \in I$

For an x in E we put $S(x) = \{y \in E/x \leq y\}$.

Let A be a subset of E , $\text{diam}(A) = \sup_{i \in I} (\sup_{\substack{x \in A \\ y \in A}} d_i(x, y))$ will be called diameter of A .

We denote by 2^E , the set of all nonempty subsets of E .

I-Single Valued Mappings

We will begin with the following result:

Theorem I-1. *Let E be a uniform space and $p : E \rightarrow \mathbb{R}_+$ a real functional which is lower semi-continuous (l.s.c.). Let $f : E \rightarrow E$ be an arbitrary self-mapping of E .*

(I-1): *there exists an x in E such that, for all $i \in I$,*
 $d_i(y, f(y)) \leq p(y) - p(f(y))$, for all $y \in S(x)$.

(I-2): *any Cauchy sequence in $S(x)$ converges in E .*

Then f has a fixed point which is maximal in (E, \leq) .

Proof. Construct a sequence $(x_n)_n$ in E inductively as follows: $x_1 = x$; when x_1, x_2, \dots, x_n have been chosen, let $a_n := \inf p(S(x_n))$ and take x_{n+1} in $S(x_n)$ such that $p(x_{n+1}) \leq a_n + 1/n$.

Then $x_n \leq x_{n+1}$ and for any y in $S(x_n)$ we have $a_{n-1} \leq a_n \leq p(y) \leq p(x_n) \leq a_{n-1} + 1/n - 1$.

In particular, for $n \leq m$ we have $0 \leq p(x_n) - p(x_m) \leq 1/n - 1$.

This shows that $(x_n)_n$ is a Cauchy sequence and that, for all $i \in I$, $d_i(x_n, x_m)$ converges to zero when n tends to infinity. Hence $\text{diam}(S(x_n))$ converges to zero for all n .

By hypothesis $(x_n)_n$ converges to an x_0 in E . On the other hand, by the construction of $(x_n)_n$, we have, for all $i \in I$, $d_i(x_n, x_{n+k}) \leq p(x_n) - p(x_{n+k})$, for all $k \geq 0$.

Hence, allowing k tend to infinity we have $d_i(x_n, x_0) \leq p(x_n) - p(x_0)$, for all n and for all $i \in I$. This means that $x_n \leq x_0$ for all n . Therefore $x_0 \in S(x)$ and

$$d_i(x_0, f(x_0)) \leq p(x_0) - p(f(x_0)), \text{ for all } i \in I \text{ i.e. } x_0 \leq f(x_0)$$

Let now $(y_n)_n$ be a sequence such that $x_n \leq y_n$ for all n . Then $\lim_n y_n = x_0$, for $\text{diam}(S(x_n))$ converges to zero for all n .

Finally, suppose that y in E is such that $x_0 \leq y$. Then we also have $x_n \leq y$ for all n and it follows that $y = x_0$ (take $y_n := y$ for all n in the preceding sequence), i.e. x_0 is maximal and then $f(x_0) = x_0$. \square

We have the following corollary:

Corollary I-2. *Let E be a sequentially complete uniform space and $p : E \rightarrow \mathbb{R}_+$ a l.s.c. real functional. Let f be an arbitrary self-mapping of E . Suppose that there exists an x in E such that $d_i(y, f(y)) \leq p(y) - p(f(y))$, for all y in $S(x)$ and for all $i \in I$.*

Then f has a fixed point which is a maximal element in (E, \leq) .

As a consequence of this result we have

Corollary I-3. (*Caristi-Kirk's theorem*). Let (E, d) be a complete metric space and $p : E \rightarrow \mathbb{R}_+$ a l.s.c. real functional. Let f be a self-mapping of E such that $p(x, f(x)) \leq p(x) - p(f(x))$, for all x in E . Then f has a fixed point.

Analyzing the proof of Theorem I-1, we can state the following

Theorem I-4. Let E be a uniform space and $p : E \rightarrow \mathbb{R}_+$ a l.s.c. real functional. Let f be an arbitrary self-mapping of E . Suppose that:

(I-3): there exists an x in E such that, for all $i \in I$,
 $d_i(y, f(y)) \leq p(y) - p(f(y))$, for every y in $S(x)$

(I-4): any nondecreasing sequence in $S(x)$ is relatively compact.

Then f has at least one fixed point which is maximal in (E, \leq) .

Proof. Let $(x_n)_n$ be a sequence defined as follows: $x_1 = x$; when x_1, x_2, \dots, x_n have been chosen let $a_n := \inf p(S(x_n))$ and take x_{n+1} in $S(x_n)$ such that $p(x_{n+1}) \leq a_n + 1/n$. The sequence $(x_n)_n$ is increasing. One shows that, for all $i \in I, d_i(x_n, x_m) \leq p(x_n) - p(x_m) \leq 1/n - 1/m$, for $n \leq m$. Hence $(x_n)_n$ is a Cauchy sequence; moreover $\text{diam}(S(x_n))$ converges to zero for all n .

By hypothesis, $(x_n)_n$ is relatively compact. Therefore there exists a subsequence $(x_{n_k})_k$ of $(x_n)_n$ which converges to an x_0 in E . Since $(x_n)_n$ is a Cauchy sequence, it also converges to x_0 . By the same argument as in the proof of Theorem I-1 we get that $x_0 \in S(x)$ and that x_0 is maximal. Thus by hypothesis, for all $i \in I, d_i(x_0, f(x_0)) \leq p(x_0) - p(f(x_0))$, i.e. $x_0 \leq f(x_0)$. Thus $f(x_0) = x_0$.

□

Remark I-5. Instead of condition (I-4) in Theorem I-4, if we suppose that $S(x)$ is complete for each x in E , the conclusion of the theorem still holds.

If we suppose likewise that E is sequentially complete, the condition (I-4) is no more needed. Corollary I-2 can again be obtained as a consequence.

II-Multi-Valued Mappings

Now we give an extension of Theorem I-1 to the case of certain multi-valued mapping, namely those which in some way are p -contractive. Thus we improve some of the results of M-H. Shih ([8]) which are of Caristi- Kirk type.

We slightly soften a definition of M-H. Shih ([8]).

Definition II-1. Let A be a subset of E . A multi-valued mapping $f : E \rightarrow 2^E$ is said to be a weak p -contraction on A , if there exists a real functional $p : E \rightarrow \mathbb{R}_+$ such that for each x in A and $y \in f(x)$, $d_i(x, y) \leq p(x) - p(y)$, for all $i \in I$.

f is said to be a p -contraction on A , if for each x in A and all y in $f(x)$, $d_i(x, y) \leq p(x) - p(y)$, for all $i \in I$.

f is said to be a weak p -contraction (respectively a p -contraction) in the sense of Shih, if $A = E$, E being a metric space.

We have the following result:

Theorem II-2. Let E be a uniform space and $f : E \rightarrow 2^E$ a closed multi-valued mapping. Suppose that:

(II-1): there exists an x in E such that f is a weak p -contraction on $S(x)$;

(II-2): any Cauchy sequence in $S(x)$ converges in E .

Then f has a fixed point.

Proof. Endow E with the partial order corresponding to p and construct a sequence $(x_n)_n$ as follows: $x_1 = x$ and for $n > 1$, take x_{n+1} in $f(x_n)$ such that $x_n \leq x_{n+1}$ (this is possible for f is a weak p -contraction on $S(x)$). One shows that $(x_n)_n$ is a Cauchy sequence. Therefore by hypothesis, there exists $x^* \in f(x^*)$, i.e. x^* is a fixed point of f .

Moreover, we have, for all $i \in I$,

$$d_i(x_n, x_{n+k}) \leq p(x_n) - p(x_{n+k}), \text{ for } k \geq 0.$$

Tending k to infinity we obtain

$$d_i(x_n, x^*) \leq p(x_n), \text{ for all } i \in I, n = 0, 1, 2 \dots$$

As a consequence we get the following: □

Corollary II-3. Let E be a sequentially complete uniform space and $f : E \rightarrow 2^*$ a closed multi-valued mapping. Suppose that f is a weak p -contraction. Then f has a fixed point.

Corollary II-4 (M-H. Shih ([8])). Let (E, d) be a complete metric space and $f : E \rightarrow 2^E$ a closed multi-valued mapping. Suppose that f is a weak p -contraction. Then f has a fixed point.

We will need the following statement, in uniform spaces, of Ekeland's variational principle. A. Brøndsted ([3]) stated it differently (in uniform spaces). Here we give a statement directly applicable to our case.

Theorem II-5. *Let E be a sequentially complete uniform space and $p : E \rightarrow \mathbb{R}$ a l.s.c. real functional which is bounded below. Then there exists an x in E such that:*

$$(II - 3) : \forall y \neq x, \exists i_o \in I : p(y) > p(x) - d_{i_o}(x, y).$$

Now we can state the following

Theorem II-6 *Let E be a uniform space and $f : E \rightarrow 2^E$ a multi-valued mapping. Suppose that:*

(II-4): there exists an x in E such that f is a weak p -contraction on $S(x)$ with p being I.s.c. and $S(x)$ complete. Then f has a fixed point.

Proof. By Theorem II-5, there exists $v \in S(x)$ such that for every $w \neq v$, there exists $i_o \in I$ such that $p(w) - p(v) > -d_{i_o}(w, v)$. We assert that $v \in f(v)$. Indeed, if not, then $p(w) - p(v) > d_{i_o}(w, v)$, for each w in $f(v)$. Whence a contradiction to the p -contractness of f on $S(x)$. \square

Corollary II-7. *Let E be a sequentially complete uniform space and $f : E \rightarrow 2^E$ a multi-valued mapping. Suppose that f is a weak p -contraction with p being I.s.c. Then f has a fixed point.*

From the previous corollary we deduce the following result:

Corollary II-8 (M-H. Shih ([8])). *Let (E, d) be a complete metric space and $f : E \rightarrow 2^E$ a multi-valued mapping. Suppose that f is a weak p -contraction with p being I.s.c. Then f has a fixed point.*

Remark II-9. Replacing weak p -contractness by p -contractness we get special cases of the results above and in particular some results of M-H. Shih ([8]).

Acknowledgments

The first author is grateful to Islamic Educational, Scientific and Cultural Organization (I.S.E.S.C.O.) for the scholarship extended during the academic years of 1991-92 and 1992-93. Thanks are due to the staff of the scientific Department of ISESCO in Rabat (Morocco).

References

- [1] Bishop, E. and Phelps, R.R., The support functionals of a convex set. Proc. Symp. in Pure Mathematics, Vol 7: 27-35 (1962).
- [2] Bressis, H. and Browder, F.E., A general principle on ordered sets in nonlinear functional analysis, Adv. in Math., 21, 355-364 (1976).
- [3] Brøndsted, A., On a lemma of Bishop and Phelps, Pac.J. Math., Vol.55, n° 2, 335-341 (1974).
- [4] — — — — — — — — — —, Fixed points and partial orders, Proc. of the Amer. math. Soc, Vol. 60, 365-366 (1976).
- [5] Conserva, V., Fixed points and partial orders, Pitmann Research Notes in Math., Serie 252: 97-99, (1991).
- [6] Ekeland, I., On the variational principle, J. Math. Anal. Appl., 47, 324-355 (1974)
- [7] Kirk, W.A., Caristi's fixed point theorem and the theory of normal solvability, fixed point theorem and its applications, Acad. Press N.Y.,S.F., and London, 109-120 (1976).
- [8] Shih, M-H., Fixed point for mappings majorized by real functionals, Hokkaido Math - Journal, vol. 9, 18-35 (1980).
- [9] Turinici, M, Maximal elements in a class of order complete metric spaces., Math. Japonica, n° 5, 511-517 (1980).

CARISTI- KIRK THEOREMİNİN BİR GENELLEŞTİRİLMESİ

Özet

Bu makalede Caristi- Kirk teoreminin düzgün uzaylara ve çok-değerli temsillere genişletilmeleri verilmiştir.

M.O. DIALLO, M. OUDADESS
Ecole Normale Supérieure-TAKADDOUM
B.P. 5118, Rabat, MAROC.

Received 23.3.1994