

1-1-1996

The minimal genus of an embedded surface of non-negative square in a rational surface

Daniel RUBERMAN

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

RUBERMAN, Daniel (1996) "The minimal genus of an embedded surface of non-negative square in a rational surface," *Turkish Journal of Mathematics*: Vol. 20: No. 1, Article 10. Available at: <https://journals.tubitak.gov.tr/math/vol20/iss1/10>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

The minimal genus of an embedded surface of non-negative square in a rational surface

Daniel Ruberman

1. Introduction

The long-standing conjecture of Thom on the minimal genus of an embedded surface in \mathbf{CP}^2 carrying a given homology class was resolved in the fall of 1994 by Kronheimer-Mrowka [KM94] and Morgan–Szabo–Taubes (to appear). Upon hearing the argument used in [KM94], I saw how to extend that proof to all rational surfaces, provided that the self-intersection of the homology class in question is non-negative. This note contains that extension. I subsequently learned that the paper of Morgan–Szabo–Taubes will include a more general result applying to any Kähler surface with $b_+^2 = 1$. Their method is less computational than that presented here.

For the purposes of the paper, a rational surface will be a 4-manifold diffeomorphic to $S^2 \times S^2$ or to $\mathbf{CP}^2 \# n \overline{\mathbf{CP}}^2$ and will be denoted by X . We will make no notational distinction between an embedded surface in X and the homology class which it carries. Choose a basis $\{S_0, S_1, \dots, S_n\}$ for the homology of X_n . Here S_0 is the complex line (with its usual orientation) in \mathbf{CP}^2 and the other S_i are the exceptional curves in the $\overline{\mathbf{CP}}^2$'s, oriented so that $-S_i$ is a complex curve. Denote by H the Poincaré dual of S_0 , and by E_i the Poincaré dual of $-S_i$, so that $E_i \cdot S_i = 1$. These classes form a basis of $H_2(X)$, so we may write (in homology) $\Sigma = \sum a_i S_i$, and will denote by $|\Sigma|$ the class $\sum |a_i| S_i$.

Theorem 1.1. *Let X be a rational surface with canonical class K_X , and let Σ be an embedded surface with self-intersection $\Sigma \cdot \Sigma > 0$. Then the genus $g(\Sigma)$ satisfies*

$$2g - 2 \geq K_X \cdot |\Sigma| + \Sigma \cdot \Sigma \tag{1}$$

If $\Sigma \cdot \Sigma = 0$, then inequality (1) holds if, in addition $K_X \cdot |\Sigma| \geq 0$.

Corollary 1.2. *A complex curve in X minimizes the genus in its homology class.*

This work was done during a visit to the Mathematical Institute in Oxford, which was supported by grant #24833 from the EPSRC. I would like to thank Peter Kronheimer for explaining the Seiberg-Witten invariants to me and for keeping me informed about his work, and Benoit Gérard for a careful reading of this paper.

Proof of Corollary: A complex curve in X_n , other than one of the exceptional curves $-S_i$, must have positive intersection with S_0 and all of the $-S_i$. So the coefficients all are non-negative, and $|\Sigma| = \Sigma$. For classes of positive square, the corollary thus follows directly from the adjunction formula, which states that equality holds in (1) for Σ a complex curve. For Σ of square 0, the adjunction formula again implies that (apart from a few cases where Σ is a rational curve), the inequality $K_X \cdot |\Sigma| \geq 0$ holds, so that again Σ is genus-minimizing. \square

Because the manifolds $S^2 \times S^2$ or $\mathbf{CP}^2 \# \overline{\mathbf{CP}^2}$ admit orientation reversing diffeomorphisms, the theorem applies as well to classes of negative self intersection. As a corollary, we thus get the minimal genus for any homology class in those manifolds:

Corollary 1.3. *The minimal genus of a surface in $S^2 \times S^2$ carrying the homology class (a, b) in the obvious basis, with $ab \neq 0$, is $(|a| - 1)(|b| - 1)$. The minimal genus of a surface in $\mathbf{CP}^2 \# \overline{\mathbf{CP}^2}$ homologous to $a_0 S_0 + a_1 S_1$, assuming $|a_0| > |a_1|$, is given by*

$$\frac{(|a_0| - 1)(|a_0| - 2)}{2} - \frac{|a_1|(|a_1| + 1)}{2}$$

If $|a_0| < |a_1|$, then the genus is given by the same formula with the roles of the a_i reversed.

The statements about $S^2 \times S^2$ are proved via the diffeomorphism of $S^2 \times S^2 \# \overline{\mathbf{CP}^2}$ with $\mathbf{CP}^2 \# 2\overline{\mathbf{CP}^2}$. In $S^2 \times S^2$, the classes $(a, 0)$ and $(0, b)$ are represented by embedded spheres, so the corollary determines the minimal genus in any homology class. A similar remark applies to the missing case in the corollary, i.e. the classes $n(S_0 \pm S_1)$ are all represented by embedded spheres. We do not know in general what the minimal genus is for classes of square 0 in X_n .

2. Basic results

Our proof is a straightforward extension of the method of Kronheimer-Mrowka, which is in turn based on the new 4-manifold invariants derived from the Seiberg-Witten equation [Wit94]. We extract the basics of those invariants from the paper [KM94], to which we refer the reader for additional details.

Suppose now that X is a 4-manifold diffeomorphic to $\mathbf{CP}^2 \# n\overline{\mathbf{CP}^2}$, and that X has been equipped with a Riemannian metric g . Let L be a complex line bundle on X which is the dual of K_X , so that $c_1(L) = 3H - E = 3H - \sum E_i$ in the notation above. Let ω_g be a self-dual harmonic 2-form, normalized so that the cohomology class $[\omega_g]$ which it carries lies in the same component of the positive cone in $H^2(X, \mathbf{R})$ as the hyperplane class H . Let $W \subset H^2(X)$ be the ‘wall’ defined by the condition $x \cup c_1(L) = 0$. If the metric g satisfies the genericity condition that $[\omega_g] \notin W$, then Kronheimer-Mrowka define $n(g)$ as the number of points (counted modulo 2) in the 0-dimensional solution space to the perturbed Seiberg-Witten equations. If g_t is a path of metrics with g_0, g_1 generic, so that the corresponding path $[\omega_{g_t}]$ is transverse to W , then $n(g)$ changes by the intersection

number of W and $[\omega_{g_t}]$. Finally, for X a rational surface, they calculate that $n(g) = 1$ when $c_1(L) \cup [\omega_g]$ is negative.

3. Proof of the theorem

Suppose first that Σ is a surface in X_n with $\Sigma \cdot \Sigma \geq 0$, for which the inequality 1 fails to hold. In brief, Σ is a counterexample to theorem 1.1. Express $[\Sigma]$ as $\sum_{i=0}^n a_i S_i$, and notice that if any of the a_i is negative, then there is another counterexample Σ' with $a'_i = -a_i$. For, there is an orientation preserving diffeomorphism $\varphi : X \rightarrow X$ with $\varphi_*(S_i) = -S_i$, and $\varphi_*(S_j) = S_j$ for $i \neq j$. Let $\Sigma' = \varphi(\Sigma)$; then it is readily seen to be a counterexample as well. So we may as well assume that all the $a_i \geq 0$. We also make the remark, following [KM, Lemma 7.7] that if Σ is a counterexample to theorem 1.1, then the homology class $r\Sigma$ (for any positive r) also contains a counterexample.

With these preliminary observations in hand, suppose that Σ is a counterexample, with $\Sigma \cdot \Sigma = m \geq 0$, and form the class $\tilde{\Sigma} = \Sigma + \sum_{i=n+1}^{n+m} S_i$ in $X_n \# m \overline{\mathbf{C}\mathbf{P}^2}$. Evidently $\tilde{\Sigma}$ has the same genus as Σ , and self-intersection 0. Choose a sequence of metrics g_R with increasingly long cylinders $Y \times [-R, R]$, where $Y \cong S^1 \times \tilde{\Sigma}$ is the boundary of the tubular neighborhood of $\tilde{\Sigma}$. Normalize the corresponding harmonic forms ω_R so that the $[\omega_R] \cup H = 1$.

Lemma 3.1. *Suppose that $\Sigma \cdot \Sigma > 0$, or that $\Sigma \cdot \Sigma = 0$ and $K_X \cdot \Sigma > 0$. If Σ is a counterexample, then there is a counterexample Σ' in the class $r[\Sigma]$, ($r \geq 1$) so that applying the above construction to Σ' , then R sufficiently large implies that $c_1(L) \cup [\omega_R]$ is negative. (Here $L = -K_{X_n \# m \overline{\mathbf{C}\mathbf{P}^2}}$).*

Proof. In homology, Σ may be written as $\sum_{i=0}^n a_i S_i$; recall that we have assumed that all the $a_i \geq 0$. Because Σ has non-negative square, we must have that $a_0 \geq 1$. Also, $[\omega_R] = H + \sum_{i=1}^{n+m} x_i E_i$, where the coefficient 1 of H is due to our normalization. Since $\omega_R \cup \omega_R > 0$, we must have that $\sum x_i^2 < 1$. Now

$$\begin{aligned} [\omega_R] \cup c_1(L) &= [\omega_R] \cdot \tilde{\Sigma} + [\omega_R] \cup (3 - a_0)H + [\omega_R] \cup \sum_{i=1}^n (a_i - 1)E_i \\ &= [\omega_R] \cdot \tilde{\Sigma} + (3 - a_0) - \sum_{i=1}^n x_i(a_i - 1) \end{aligned}$$

The argument in Lemma 10 of [KM94] shows that $[\omega_R] \cdot \tilde{\Sigma} \rightarrow 0$ as $R \rightarrow \infty$. So it suffices to show that the conditions $a_0^2 - \sum_{i=1}^n a_i^2 > 0$ and $\sum_{i=1}^n x_i^2 < 1$ imply that

$$(3 - a_0) - \sum_{i=1}^n x_i(a_i - 1)$$

is negative. To approach this, maximize the above expression (as a function of x_1, \dots, x_n) with the constraint $\sum_{i=1}^n x_i^2 = 1$, in the hopes that it will be negative. The maximum value is readily found to be

$$3 - a_0 + \sqrt{\sum_{i=1}^n (a_i - 1)^2}$$

Assuming, as we may by taking a multiple of Σ , that $a_0 > 3$, this maximum is negative if

$$(a_0 - 3)^2 > n + \sum_{i=1}^n a_i^2 - 2 \sum_{i=1}^n a_i$$

i.e., if

$$a_0^2 - \sum_{i=1}^n a_i^2 - 6a_0 + 2 \sum_{i=1}^n a_i > n - 9 \quad (*)$$

Unfortunately it is not always true that $(*)$ holds, even if a_0 is very large. (Take, for example $n = a_0 - 1$, $a_1 = a_0 - 2$, and the other $a_i = 2$). However, if the a_i are all replaced by ra_i , then the left-hand side of $(*)$ becomes

$$r^2(a_0^2 - \sum_{i=1}^n a_i^2) + r(-6a_0 + 2 \sum_{i=1}^n a_i)$$

So if either $\Sigma \cdot \Sigma = a_0^2 - \sum_{i=1}^n a_i^2 > 0$, or $\Sigma \cdot \Sigma = 0$ and $K_X \cdot \Sigma = -3a_0 + \sum_{i=1}^n a_i > 0$, then $(*)$ will hold for some $r > 1$. So if Σ were a counterexample with positive square, or with 0 square and for which $\sum_{i=1}^n a_i > 3a_0$, let Σ' be a counterexample in the homology class $r[\Sigma]$, where r is chosen so that

$$(3 - ra_0) + \sum x_i(ra_i - 1) < 0$$

for all $\{x_i\}$ satisfying $\sum x_i^2 < 1$. □

Proof of Theorem 1.1: Suppose that $\Sigma \cdot \Sigma > 0$, or that $\Sigma \cdot \Sigma = 0$ and $K_X \cdot |\Sigma| > 0$. Assume, perhaps replacing Σ by some large positive multiple, that Σ is a counterexample in the homology class $a_0 S_0 + \sum a_i S_i$, chosen so that all the $a_i \geq 0$ and that lemma 3.1 applies to Σ . Starting from the fact that the invariant $n(g) = 1$ for any metric g such that $c_1(L) \cup [\omega_g] < 0$, Kronheimer-Mrowka show that

$$c_1(L) \cdot \tilde{\Sigma} \geq -(2g - 2)$$

But

$$\begin{aligned} c_1(L) \cdot \tilde{\Sigma} &= c_1(L) \cdot \Sigma + c_1(L) \cdot \sum_{i=n+1}^{n+m} S_i \\ &= -K_X \cdot \Sigma - \Sigma \cdot \Sigma \end{aligned}$$

So

$$2g - 2 \geq K_X \cdot \Sigma + \Sigma \cdot \Sigma$$

This leaves only the possible exception that $\Sigma \cdot \Sigma = 0$, and that $K_X \cdot \Sigma = 0$. In this case, we must also suppose that Σ is non-trivial in homology. The content of inequality 1 in this instance is merely that Σ is not represented by an embedded sphere. But if it were, we proceed by doing surgery on Σ , resulting in a definite manifold X' . The condition that $K_X \cdot \Sigma = 0$ means that Σ is orthogonal to the characteristic class $3S_0 + \sum_{i \geq 1} S_i$. Thus $H_2(X')$ has a characteristic element of square less (in absolute value) than its rank, which readily implies that the intersection form is non-standard, contradicting Donaldson's theorem [Don87, DK90]. \square

The condition that there exist a 'short' characteristic element, as in the last paragraph of the proof, is precisely the ingredient necessary to use the monopole equations in a simplified proof of Donaldson's theorem.

References

- [DK90] S.K. Donaldson and P.B. Kronheimer, *The Geometry of Four-Manifolds*, Clarendon Press, Oxford, 1990.
- [Don87] S.K. Donaldson, *The orientation of Yang-Mills moduli spaces and 4-manifold topology*, J. Diff. Geo. **26** (1987), 397–428.
- [KM] P.B. Kronheimer and T.S. Mrowka, *Embedded surfaces and the structure of Donaldson's polynomial invariants*, J. Diff. Geo., **41** (1995), 573–734.
- [KM94] P.B. Kronheimer and T.S. Mrowka, *The genus of embedded surfaces in the projective plane*, Math. Res. Lett. **1** (6) (1994), 797–808.
- [Wit94] E. Witten, *Monopoles and four-manifolds*, Math. Res. Lett. **1** (6) (1994), 769–796.

DEPARTMENT OF MATHEMATICS, BRANDEIS UNIVERSITY, WALTHAM, MA 02254, USA
E-mail address: ruberman@binah.cc.brandeis.edu