

1-1-2022

On the existence for parametric boundary value problems of a coupled system of nonlinear fractional hybrid differential equations

YIGE ZHAO

YIBING SUN

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

ZHAO, YIGE and SUN, YIBING (2022) "On the existence for parametric boundary value problems of a coupled system of nonlinear fractional hybrid differential equations," *Turkish Journal of Mathematics*: Vol. 46: No. 2, Article 18. <https://doi.org/10.3906/mat-2107-19>
Available at: <https://journals.tubitak.gov.tr/math/vol46/iss2/18>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

On the existence for parametric boundary value problems of a coupled system of nonlinear fractional hybrid differential equations

Yige ZHAO* , Yibing SUN 

School of Mathematical Sciences, University of Jinan, Jinan, Shandong, P.R. China

Received: 08.07.2021

Accepted/Published Online: 10.12.2021

Final Version: 21.01.2022

Abstract: In this paper, we consider the existence and uniqueness for parametric boundary value problems of a coupled system of nonlinear fractional hybrid differential equations. By the fixed point theorem in Banach algebra, an existence theorem for parametric boundary value problems of a coupled system of nonlinear fractional hybrid differential equations is given. Further, a uniqueness result for parametric boundary value problems of a coupled system of nonlinear fractional hybrid differential equations is proved due to Banach's contraction principle. Further, we give three examples to verify the main results.

Key words: Boundary value problems, coupled fractional hybrid differential systems, Dhage fixed point theorem, existence.

1. Introduction

In this paper, we discuss the following parametric boundary value problems (in short PBVP) of a coupled system of nonlinear fractional hybrid differential equations (in short FHDS)

$$\begin{cases} {}^C D_{0+}^{\alpha} \left[\frac{y(\zeta) - k_1(\zeta, y(\zeta))}{f_1(\zeta, y(\zeta))} \right] = g_1(\zeta, y(\zeta), z(\zeta)), & 0 < \zeta < 1, \\ {}^C D_{0+}^{\beta} \left[\frac{z(\zeta) - k_2(\zeta, z(\zeta))}{f_2(\zeta, z(\zeta))} \right] = g_2(\zeta, y(\zeta), z(\zeta)), & 0 < \zeta < 1, \\ a_1 \left[\frac{y(\zeta) - k_1(\zeta, y(\zeta))}{f_1(\zeta, y(\zeta))} \right]_{\zeta=0} + b_1 \left[\frac{y(\zeta) - k_1(\zeta, y(\zeta))}{f_1(\zeta, y(\zeta))} \right]_{\zeta=1} = c_1, \\ a_2 \left[\frac{z(\zeta) - k_2(\zeta, z(\zeta))}{f_2(\zeta, z(\zeta))} \right]_{\zeta=0} + b_2 \left[\frac{z(\zeta) - k_2(\zeta, z(\zeta))}{f_2(\zeta, z(\zeta))} \right]_{\zeta=1} = c_2, \end{cases} \quad (1.1)$$

where $0 < \alpha, \beta \leq 1$, ${}^C D_{0+}^{\alpha}$ and ${}^C D_{0+}^{\beta}$ are the Caputo derivatives, $\Delta = [0, 1]$, $f_i \in C(\Delta \times \mathbb{R}, \mathbb{R} \setminus \{0\})$, $k_i \in C(\Delta \times \mathbb{R}, \mathbb{R})$, $g_i \in C(\Delta \times \mathbb{R} \times \mathbb{R}, \mathbb{R})$ ($i = 1, 2$), and a_i, b_i, c_i ($i = 1, 2$) are real constants with $a_i + b_i \neq 0$ ($i = 1, 2$).

The theory and the applications of the FDE have gained many researchers' attention, see the study in [10]. Many papers on the solvability of the nonlinear FDE and FDS, see [6, 12, 14, 15, 17]. In recent years, the theory of the HDE has been a hot research topic; see [1–5, 7, 8, 11, 13, 16, 18]. Dhage [4] discussed the following first order hybrid differential equation

$$\begin{cases} \frac{d}{d\zeta} \left[\frac{x(\zeta) - k(\zeta, x(\zeta))}{f(\zeta, x(\zeta))} \right] = g(\zeta, x(\zeta)), & \zeta \in [\zeta_0, \zeta_0 + a], \\ x(\zeta_0) = x_0 \in \mathbb{R}, \end{cases}$$

*Correspondence: zhaoeager@126.com

2010 AMS Mathematics Subject Classification: 34A08 34B08

and developed the theory of the HDE and gave some original and interesting results. Dumitru Baleanu et al. [2] investigated sufficient conditions for existence and uniqueness of solutions for a coupled system of fractional order hybrid differential equations with multi-point hybrid boundary conditions due to Dhage and Banach's contraction principle. Amjad Ali [1] established necessary and sufficient conditions for existence of positive solutions for coupled systems of nonlinear hybrid differential equations subject to nonhomogeneous boundary conditions using hybrid fixed point theorem due to Dhage. Dhage et al. [5] studied a system of two nonhomogeneous boundary value problems of coupled hybrid integro-differential equations of fractional order by a hybrid fixed point theorem due to Dhage in Banach algebras. You and Sun [13] discussed a class of impulsive coupled hybrid fractional differential system due to a new hybrid fixed point theorem in Banach algebra.

To the best of our knowledge, there are no results for the PBVP (1.1) of the nonlinear coupled FHDS. From the above works, we consider the existence and uniqueness of the BVP (1.1) of the nonlinear coupled FHDS. An existence theorem and a uniqueness result for the PBVP (1.1) of the nonlinear coupled FDS are obtained. Further, three examples are given to verify the main results. To some extent, our work fills the gap on some basic theory for BVP of the nonlinear coupled FHDS.

The paper is organized as follows: Section 2 gives some theory of fractional calculus. Section 3 establishes an existence theorem for the PBVP (1.1) of the nonlinear coupled FHDS by the fixed point theorem in Banach algebra. Section 4 considers a uniqueness result for the PBVP (1.1) of the nonlinear coupled FHDS by Banach's contraction principle. Section 5 presents three examples to verify the existence theorem.

2. Preliminary

In this section, we give some basic theory from fractional calculus, see [3, 9, 13, 18].

Definition 2.1 ([9]) *The Caputo fractional derivative of order $0 < \alpha < 1$ of a continuous function $f : (0, +\infty) \rightarrow \mathbb{R}$ is given by*

$${}^C D_{0+}^\alpha f(\zeta) = \frac{1}{\Gamma(1-\alpha)} \int_0^\zeta \frac{f'(\omega)}{(\zeta-\omega)^\alpha} d\omega.$$

Definition 2.2 ([9]) *The Riemann-Liouville fractional integral of order $\alpha > 0$ of an integrable function $f : (0, +\infty) \rightarrow \mathbb{R}$ is given by*

$$I_{0+}^\alpha f(\zeta) = \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta-\omega)^{\alpha-1} f(\omega) d\omega,$$

provided that the right side is pointwise defined on $(0, +\infty)$.

Lemma 2.3 ([9]) *Let $\alpha > 0$. If we assume $y \in C(0, 1) \cap L(0, 1)$, then the fractional differential equation*

$${}^C D_{0+}^\alpha y(\zeta) = 0$$

has $y(\zeta) = c_0 + c_1\zeta + c_2\zeta^2 + \dots + c_{n-1}\zeta^{n-1}$, $c_i \in \mathbb{R}$, $i = 0, 1, 2, \dots, n-1$, as unique solutions, where n is the smallest integer greater than or equal to α .

Lemma 2.4 ([9]) *Assume that $y \in C^n[0, 1]$ with a fractional derivative of order $\alpha > 0$ that belongs to $C^n[0, 1]$. Then*

$$I_{0+}^\alpha {}^C D_{0+}^\alpha y(\zeta) = y(\zeta) + c_0 + c_1\zeta + c_2\zeta^2 + \dots + c_{n-1}\zeta^{n-1},$$

for some $c_i \in \mathbb{R}$, $i = 0, 1, 2, \dots, n - 1$, where n is the smallest integer greater than or equal to α .

Let $C(\Delta, \mathbb{R})$ be the space of all continuous functions defined on Δ . $\|\cdot\|$ denotes a supremum norm in $C(\Delta, \mathbb{R})$ by

$$\|y\| = \sup_{\zeta \in \Delta} |y(\zeta)|,$$

and a multiplication “ \cdot ” in $C(\Delta, \mathbb{R})$ by

$$(y \cdot z)(\zeta) = (yz)(\zeta) = y(\zeta)z(\zeta)$$

for $y, z \in C(\Delta, \mathbb{R})$. Clearly $C(\Delta, \mathbb{R})$ is a Banach algebra with respect to above norm and multiplication in it.

Lemma 2.5 ([18]) *Suppose that a, b, c are real constants with $a + b \neq 0$. Then for any $z \in L(\Delta, \mathbb{R})$, the function y is a solution of the PBVP*

$${}^c D_{0+}^\alpha \left[\frac{y(\zeta) - k(\zeta, y(\zeta))}{f(\zeta, y(\zeta))} \right] = z(\zeta), \quad 0 < \alpha \leq 1, \quad \zeta \in \Delta, \tag{2.1}$$

and

$$a \left[\frac{y(\zeta) - k(\zeta, y(\zeta))}{f(\zeta, y(\zeta))} \right]_{\zeta=0} + b \left[\frac{y(\zeta) - k(\zeta, y(\zeta))}{f(\zeta, y(\zeta))} \right]_{\zeta=1} = c, \tag{2.2}$$

if and only if y satisfies the integral equation

$$\begin{aligned} y(\zeta) &= f(\zeta, y(\zeta)) \left(\frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} z(\omega) d\omega \right. \\ &\quad \left. + \frac{1}{a+b} \left(c - \frac{b}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} z(\omega) d\omega \right) \right) + k(\zeta, y(\zeta)), \quad \zeta \in \Delta. \end{aligned} \tag{2.3}$$

Let $\tilde{U} = C(\Delta, \mathbb{R})$. Define multiplication and the sum on $X \times X$ as

$$(y_1, z_1) + (y_2, z_2) = (y_1 + y_2, z_1 + z_2),$$

$$p(y, z) = (py, pz), p \in \mathbb{R}.$$

Lemma 2.6 ([3]) *Let $\bar{U} = U \times U$. Define the product in X by*

$$(y_1, z_1)(y_2, z_2) = (y_1 y_2, z_1 z_2),$$

and

$$\|(y, z)\| = 2(\|y\| + \|z\|).$$

Then \bar{U} is a Banach algebra with respect to the above norm and multiplication.

Lemma 2.7 ([13]) *Let \tilde{S} be a nonempty, closed, convex and bounded subset of a Banach algebra X and $\bar{U} = \tilde{U} \times \tilde{S}, \bar{S} = \tilde{S} \times \tilde{S}$. Suppose $\tilde{A}_i, \tilde{C}_i : \tilde{U} \rightarrow \tilde{U}, \tilde{B}_i : \tilde{S} \rightarrow \tilde{U} (i = 1, 2)$ are operators satisfying*

(a) There exist $0 < \tilde{\rho}_i, \tilde{\delta}_i < 1$, such that, respectively,

$$\| \tilde{A}_i y - \tilde{A}_i z \| \leq \tilde{\rho}_i \| y - z \|, \| \tilde{C}_i y - \tilde{C}_i z \| \leq \tilde{\delta}_i \| y - z \|,$$

for all $y, z \in \tilde{U}$, $i = 1, 2$, $\tilde{\rho} = \max \{ \tilde{\rho}_1, \tilde{\rho}_2 \}$, $\tilde{\delta} = \max \{ \tilde{\delta}_1, \tilde{\delta}_2 \}$,

(b) \tilde{B}_i is completely continuous, $i = 1, 2$,

(c) $y = \tilde{A}_1 y \tilde{B}_1 z + \tilde{C}_1 y$ for all $z \in \tilde{S} \Rightarrow y \in \tilde{S}$, and $z = \tilde{A}_2 z \tilde{B}_2 y + \tilde{C}_2 z$ for all $y \in \tilde{S} \Rightarrow z \in \tilde{S}$,

(d) $4\rho \| \tilde{B}(\tilde{S}) \| + \tilde{\delta} < 1$, $\| \tilde{B}(\tilde{S}) \| = \max \{ \sup \{ \| \tilde{B}_1(y) \| : y \in \tilde{S} \}, \sup \{ \| \tilde{B}_2(y) \| : y \in \tilde{S} \} \}$.

Then the operator equation $(\tilde{T}_1(y, z), \tilde{T}_2(y, z)) = (y, z)$ has a fixed point in \bar{S} , where $\tilde{T}_1, \tilde{T}_2 : \bar{U} \rightarrow \bar{U}$ are defined by

$$\tilde{T}_1(y, z) = \tilde{A}_1 y \tilde{B}_1 z + \tilde{C}_1 y, \quad \tilde{T}_2(y, z) = \tilde{A}_2 z \tilde{B}_2 y + \tilde{C}_2 z, \quad (y, z) \in \bar{U}.$$

3. Existence result

In this section, we discuss the existence results for the PBVP (1.1) of the nonlinear coupled FHDS.

We present the following hypotheses.

(H₁) There exist constants $L_i > 0$ ($i = 1, 2$) and $\bar{L}_i > 0$ ($i = 1, 2$) such that

$$|f_i(\zeta, y) - f_i(\zeta, z)| \leq L_i |y - z|$$

and

$$|k_i(\zeta, y) - k_i(\zeta, z)| \leq \bar{L}_i |y - z|$$

for all $\zeta \in \Delta$ and $y, z \in \mathbb{R}$.

(H₂) There exist two nonnegative functions $d_i(\zeta) \in L(0, 1)$ ($i = 1, 2$) such that $g_i(\zeta, y, z) \leq d_i(\zeta) + m_i |y|^{\rho_i} + n_i |z|^{\theta_i}$, where $m_i, n_i \geq 0$, $0 < \rho_i, \theta_i < 1$, for $i = 1, 2$.

Now we will give the following existence theorem for the PBVP (1.1) of the nonlinear coupled FHDS.

Theorem 3.1 Suppose that (H₁) and (H₂) hold. Then the PBVP (1.1) of the nonlinear coupled FHDS has a solution.

Proof. By Lemma 2.5, the system (1.1) is equivalent to the following integral system

$$\begin{cases} y(\zeta) = f_1(\zeta, y(\zeta)) \left(\frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right. \\ \quad \left. + \frac{1}{a_1+b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right) \right) \\ \quad + k_1(\zeta, y(\zeta)), \quad \zeta \in \Delta. \\ z(\zeta) = f_2(\zeta, z(\zeta)) \left(\frac{1}{\Gamma(\beta)} \int_0^\zeta (\zeta - \omega)^{\beta-1} g_2(\omega, y(\omega), z(\omega)) d\omega \right. \\ \quad \left. + \frac{1}{a_2+b_2} \left(c_2 - \frac{b_2}{\Gamma(\beta)} \int_0^1 (1 - \omega)^{\beta-1} g_2(\omega, y(\omega), z(\omega)) d\omega \right) \right) \\ \quad + k_2(\zeta, z(\zeta)), \quad \zeta \in \Delta. \end{cases} \tag{3.1}$$

Set $U = C(\Delta, \mathbb{R})$ and define a subset S of U by

$$S = \{ y \in U \mid \|y\| \leq R \},$$

where

$$R \geq \max \left\{ 5l_1, 5\bar{l}_1, \frac{5|c_1|}{|a_1 + b_1|}, 5l_2, 5\bar{l}_2, \frac{5|c_2|}{|a_2 + b_2|}, \right. \\ \left. \left(\frac{5m_1(|b_1| + |a_1 + b_1|)}{|a_1 + b_1|\Gamma(\alpha + 1)} \right)^{\frac{1}{1-\rho_1}}, \left(\frac{5n_1(|b_1| + |a_1 + b_1|)}{|a_1 + b_1|\Gamma(\alpha + 1)} \right)^{\frac{1}{1-\theta_1}}, \right. \\ \left. \left(\frac{5m_2(|b_2| + |a_2 + b_2|)}{|a_2 + b_2|\Gamma(\beta + 1)} \right)^{\frac{1}{1-\rho_2}}, \left(\frac{5n_2(|b_2| + |a_2 + b_2|)}{|a_2 + b_2|\Gamma(\beta + 1)} \right)^{\frac{1}{1-\theta_2}}, \right. \\ \left. \frac{F_0M_1 + K_0}{1 - L_1M_1 - \bar{L}_1}, \frac{F_0M_2 + K_0}{1 - L_2M_2 - \bar{L}_2} \right\},$$

and

$$l_1 = \max_{\zeta \in \Delta} \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} d_1(\omega) d\omega, \quad \bar{l}_1 = \frac{|b_1|}{|a_1 + b_1|\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} d_1(\omega) d\omega, \\ l_2 = \max_{\zeta \in \Delta} \frac{1}{\Gamma(\beta)} \int_0^\zeta (\zeta - \omega)^{\beta-1} d_2(\omega) d\omega, \quad \bar{l}_2 = \frac{|b_2|}{|a_2 + b_2|\Gamma(\beta)} \int_0^1 (1 - \omega)^{\beta-1} d_2(\omega) d\omega, \\ M_1 = \frac{K}{\Gamma(\alpha + 1)} \left(1 + \frac{|b_1|}{|a_1 + b_1|} \right) + \frac{|c_1|}{|a_1 + b_1|}, \\ M_2 = \frac{K}{\Gamma(\beta + 1)} \left(1 + \frac{|b_2|}{|a_2 + b_2|} \right) + \frac{|c_2|}{|a_2 + b_2|}, \\ K = \max_{\zeta \in \Delta} \{ |g_1(\zeta, y(\zeta), z(\zeta))|, |g_2(\zeta, y(\zeta), z(\zeta))| \}, \\ F_0 = \max_{\zeta \in \Delta} \{ \sup |f_1(\zeta, 0)|, \sup |f_2(\zeta, 0)| \}, \\ K_0 = \max_{\zeta \in \Delta} \{ \sup |k_1(\zeta, 0)|, \sup |k_2(\zeta, 0)| \}, \quad L = \max\{L_1, L_2\}, \quad \bar{L} = \max\{\bar{L}_1, \bar{L}_2\}, \\ M = \max\{M_1, M_2\}, \quad 4LM + 4\bar{L} < 1.$$

Clearly, S is a closed, convex and bounded subset of the Banach space U .

Define the operators $A_i, C_i : U \rightarrow U$ and $B_i : S \rightarrow U$ ($i = 1, 2$) by

$$A_1y(\zeta) = f_1(\zeta, y(\zeta)), \quad A_2z(\zeta) = f_2(\zeta, z(\zeta)), \quad \zeta \in \Delta, \tag{3.2}$$

$$\begin{cases} B_1z(\zeta) = \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \\ \quad + \frac{1}{a_1+b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right), \quad \zeta \in \Delta. \\ B_2y(\zeta) = \frac{1}{\Gamma(\beta)} \int_0^\zeta (\zeta - \omega)^{\beta-1} g_2(\omega, y(\omega), z(\omega)) d\omega \\ \quad + \frac{1}{a_2+b_2} \left(c_2 - \frac{b_2}{\Gamma(\beta)} \int_0^1 (1 - \omega)^{\beta-1} g_2(\omega, y(\omega), z(\omega)) d\omega \right), \quad \zeta \in \Delta. \end{cases} \tag{3.3}$$

and

$$C_1y(\zeta) = k_1(\zeta, y(\zeta)), \quad C_2z(\zeta) = k_2(\zeta, z(\zeta)), \quad \zeta \in \Delta. \tag{3.4}$$

Then the system (3.1) is transformed into the system of operator equation as

$$\begin{cases} y(\zeta) &= A_1y(\zeta)B_1z(\zeta) + C_1y(\zeta), \\ z(\zeta) &= A_2z(\zeta)B_2y(\zeta) + C_2z(\zeta). \end{cases}$$

Set $\bar{U} = U \times U$, $\bar{S} = S \times S$. Define operators $\bar{A}, \bar{C} : \bar{U} \rightarrow \bar{U}$ and $\bar{B} : \bar{S} \rightarrow \bar{U}$ by

$$\bar{A}(y(\zeta), z(\zeta)) = (A_1y(\zeta), A_2z(\zeta)), \quad \bar{B}(y(\zeta), z(\zeta)) = (B_1z(\zeta), B_2y(\zeta)),$$

$$\bar{C}(y(\zeta), z(\zeta)) = (C_1y(\zeta), C_2z(\zeta)).$$

and define the operators $T_1 : \bar{U} \rightarrow U$, $T_2 : \bar{U} \rightarrow U$ by

$$T_1(y, z) = A_1yB_1z + C_1y, \quad T_2(y, z) = A_2zB_2y + C_2z.$$

Thus, we just prove that $(T_1(y, z), T_2(y, z)) = (y, z)$ has one solution in \bar{S} .

Next, we prove the operators A_i , B_i , and C_i ($i = 1, 2$) satisfy all the conditions of Lemma 2.7.

Firstly, we prove that (a) of Lemma 2.7 is satisfied. Let $y, z \in U$. Then by (H_1) ,

$$|A_iy(\zeta) - A_iz(\zeta)| = |f_i(\zeta, y(\zeta)) - f_i(\zeta, z(\zeta))| \leq L_i|y(\zeta) - z(\zeta)| \leq L_i\|y - z\|, \quad i = 1, 2,$$

for all $\zeta \in \Delta$. Taking supremum over ζ , then we have

$$\|A_iy - A_iz\| \leq L_i\|y - z\|, \quad i = 1, 2,$$

for all $y, z \in U$. Similarly, it can be implied that

$$\|C_iy - C_iz\| \leq \bar{L}_i\|y - z\|, \quad i = 1, 2,$$

for all $y, z \in U$. Thus, A_i and C_i ($i = 1, 2$) satisfy (a) of Lemma 2.7.

Next, we prove B_i ($i = 1, 2$) are compact and continuous operators on S into U . Firstly, we prove B_i ($i = 1, 2$) are continuous on S . Let $\{(y_n, z_n)\}$ be a sequence in \bar{S} converging to $(y, z) \in \bar{S}$. Then by the

Lebesgue dominated convergence theorem,

$$\begin{aligned}
 & \lim_{n \rightarrow \infty} B_1 z_n(\zeta) \\
 &= \lim_{n \rightarrow \infty} \left(\frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y_n(\omega), z_n(\omega)) d\omega \right. \\
 & \quad \left. + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y_n(\omega), z_n(\omega)) d\omega \right) \right) \\
 &= \lim_{n \rightarrow \infty} \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y_n(\omega), z_n(\omega)) d\omega \\
 & \quad + \lim_{n \rightarrow \infty} \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y_n(\omega), z_n(\omega)) d\omega \right) \\
 &= \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \\
 & \quad + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right) \\
 &= B_1 z(\zeta),
 \end{aligned}$$

for all $\zeta \in \Delta$. This shows that B_1 is a continuous operator on S . Similarly, we can prove that B_2 is also a continuous operator on S .

Next we prove B_i ($i = 1, 2$) are compact operators on S . It is enough to show that $B_i(S)$ ($i = 1, 2$) are uniformly bounded and equicontinuous sets in U . On the one hand, let $y \in S$ be arbitrary. Then by (H_2) ,

$$\begin{aligned}
 |B_1 z(\zeta)| &= \left| \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right. \\
 & \quad \left. + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right) \right| \\
 &\leq \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} |g_1(\omega, y(\omega), z(\omega))| d\omega \\
 & \quad + \frac{|b_1|}{|a_1 + b_1| \Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} |g_1(\omega, y(\omega), z(\omega))| d\omega + \frac{|c_1|}{|a_1 + b_1|} \\
 &\leq l_1 + \frac{m_1 R^{\rho_1} + n_1 R^{\theta_1}}{\Gamma(\alpha + 1)} + \bar{l}_1 + \frac{|b_1|(m_1 R^{\rho_1} + n_1 R^{\theta_1})}{|a_1 + b_1| \Gamma(\alpha + 1)} + \frac{|c_1|}{|a_1 + b_1|} \leq R,
 \end{aligned}$$

for all $\zeta \in \Delta$. Taking supremum over ζ , $\|B_1 z\| \leq R$ for all $y \in S$. Similarly, we can conclude that $\|B_2 z\| \leq R$ for all $z \in S$. This shows that B_i ($i = 1, 2$) are uniformly bounded on S .

On the other hand, let $\zeta_1, \zeta_2 \in \Delta$. Then for any $z \in S$, we get

$$\begin{aligned} & |B_1z(\zeta_1) - B_1z(\zeta_2)| \\ &= \left| \frac{1}{\Gamma(\alpha)} \int_0^{\zeta_1} (\zeta_1 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right. \\ &\quad \left. - \frac{1}{\Gamma(\alpha)} \int_0^{\zeta_2} (\zeta_2 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right| \\ &\leq \frac{1}{\Gamma(\alpha)} \left| \int_0^{\zeta_1} (\zeta_1 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right. \\ &\quad \left. - \int_0^{\zeta_1} (\zeta_2 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right. \\ &\quad \left. + \int_0^{\zeta_1} (\zeta_2 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega - \int_0^{\zeta_2} (\zeta_2 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right| \\ &= \frac{K}{\Gamma(\alpha + 1)} (\zeta_2^\alpha - \zeta_1^\alpha - (\zeta_2 - \zeta_1)^\alpha). \end{aligned}$$

Since the functions ζ^α is uniformly continuous on compact Δ , from the above analysis, $B_1(S)$ is an equicontinuous set in U . Thus, B_1 is completely continuous likewise B_2 .

Next, we show that (c) of Lemma 2.7 is satisfied. Let $y \in S$ and $z \in S$ be arbitrary such that $y = A_1yB_1z + C_1y$, $z = A_2zB_2y + C_2z$. Then, by assumption (H_1) , we have

$$\begin{aligned} & |y(\zeta)| \\ &\leq |A_1y(\zeta)||B_1z(\zeta)| + |C_1y(\zeta)| \\ &= |f_1(\zeta, y(\zeta))| \left| \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right. \\ &\quad \left. + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right) \right| + |k_1(\zeta, y(\zeta))| \\ &\leq [|f_1(\zeta, y(\zeta)) - f_1(\zeta, 0)| + |f_1(\zeta, 0)|] \cdot \left(\frac{K}{\Gamma(\alpha + 1)} \left(1 + \frac{|b_1|}{|a_1 + b_1|} \right) + \frac{|c_1|}{|a_1 + b_1|} \right) \\ &\quad + |k_1(\zeta, y(\zeta)) - k_1(\zeta, 0)| + |k_1(\zeta, 0)| \\ &\leq [L_1|y(\zeta)| + F_0]M_1 + \bar{L}_1|y(\zeta)| + K_0. \end{aligned}$$

Thus, we get

$$|y(\zeta)| \leq \frac{F_0M_1 + K_0}{1 - L_1M_1 - \bar{L}_1}.$$

Taking supremum over ζ ,

$$\|y\| \leq \frac{F_0M_1 + K_0}{1 - L_1M_1 - \bar{L}_1} \leq R.$$

Similarly, we can obtain that

$$\|z\| \leq \frac{F_0M_2 + K_0}{1 - L_2M_2 - \bar{L}_2} \leq R.$$

This shows that (c) of Lemma 2.7 is satisfied.

Finally, we obtain

$$\begin{aligned} & \|B(S)\| \\ &= \max\{\sup\{\|B_1(y)\| : y \in S\}, \sup\{\|B_2(y)\| : y \in S\}\} \\ &= \max\left\{\frac{K}{\Gamma(\alpha + 1)} \left(1 + \frac{|b_1|}{|a_1 + b_1|}\right) + \frac{|c_1|}{|a_1 + b_1|}, \right. \\ &\quad \left. \frac{K}{\Gamma(\beta + 1)} \left(1 + \frac{|b_2|}{|a_2 + b_2|}\right) + \frac{|c_2|}{|a_2 + b_2|}\right\} \\ &= \max\{M_1, M_2\} = M. \end{aligned}$$

and so,

$$4L\|B(S)\| + \bar{L} = 4LM + \bar{L} < 1.$$

This shows that (d) of Lemma 2.7 is satisfied.

Thus, all the conditions of Lemma 2.7 are satisfied and $(T_1(y, z), T_2(y, z)) = (y, z)$ has one solution in \bar{S} . Therefore, the PBVP (1.1) of the nonlinear coupled FHDS has a solution.

4. Uniqueness of solution

In this section, we give the uniqueness of the solution for the PBVP (1.1) of the nonlinear coupled FHDS.

We present the following hypotheses.

(H₃) There exist constants $h_i > 0$ ($i = 1, 2$) and $\bar{h}_i > 0$ ($i = 1, 2$) such that

$$|g_i(\zeta, y, z) - g_i(\zeta, y', z')| \leq h_i|y - y'| + \bar{h}_i|z - z'|$$

for all $\zeta \in \Delta$ and $y, y', z, z' \in \mathbb{R}$.

Next define the following notations

$$K' = \max_{\zeta \in \Delta} \{|f_1(\zeta, y(\zeta))|, |f_2(\zeta, z(\zeta))|\}, \quad \mu_1 = \frac{K'}{\Gamma(\alpha + 1)} \left(1 + \frac{|b_1|}{|a_1 + b_1|}\right), \quad \mu_2 = \frac{K'}{\Gamma(\beta + 1)} \left(1 + \frac{|b_2|}{|a_2 + b_2|}\right).$$

Now we will present the following uniqueness of solutions for the PBVP (1.1) of the nonlinear coupled FHDS.

Theorem 4.1 *Suppose that (H₁) and (H₃) hold. If*

$$2(\mu_1(h_1 + \bar{h}_1) + \mu_2(h_2 + \bar{h}_2) + 2M + 2\bar{L}) < 1.$$

Then the PBVP (1.1) of the nonlinear coupled FHDS has a unique solution.

Proof. Define $U = C(\Delta, \mathbb{R})$ and choose a subset S' of U by

$$S' = \{y \in U \mid \|y\| \leq R'\},$$

where

$$R' \geq \frac{4MF_0 + 4K_0}{1 - 2(LM + \bar{L})},$$

and F_0, K_0, L, \bar{L}, M are defined in Theorem 3.1. Thus, we just prove that $(T_1(y, z), T_2(y, z)) = (y, z)$ has one solution in $\bar{S}' = S' \times S'$.

For $(y, z) \in \bar{S}'$ and $\zeta \in \Delta$, we have

$$\begin{aligned} & |T_1(y(\zeta), z(\zeta))| \\ & \leq |A_1 y(\zeta)| |B_1 z(\zeta)| + |C_1 y(\zeta)| \\ & = |f_1(\zeta, y(\zeta))| \left| \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right. \\ & \quad \left. + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y(\omega), z(\omega)) d\omega \right) \right| + |k_1(\zeta, y(\zeta))| \\ & \leq [|f_1(\zeta, y(\zeta)) - f_1(\zeta, 0)| + |f_1(\zeta, 0)|] \cdot \left(\frac{K}{\Gamma(\alpha + 1)} \left(1 + \frac{|b_1|}{|a_1 + b_1|} \right) + \frac{|c_1|}{|a_1 + b_1|} \right) \\ & \quad + |k_1(\zeta, y(\zeta)) - k_1(\zeta, 0)| + |k_1(\zeta, 0)| \\ & \leq [L_1 |y(\zeta)| + F_0] M_1 + \bar{L}_1 |y(\zeta)| + K_0. \end{aligned}$$

Taking supremum over ζ , we have

$$\|T_1(y, z)\| \leq [L_1 \|y\| + F_0] M_1 + \bar{L}_1 \|y\| + K_0.$$

Similarly, we can obtain that

$$\|T_2(y, z)\| \leq [L_2 \|z\| + F_0] M_2 + \bar{L}_2 \|z\| + K_0.$$

Thus, we get

$$\begin{aligned} \|T(y, z)\| &= 2(\|T_1(y, z)\| + \|T_2(y, z)\|) \\ &\leq 2(L_1 M_1 + \bar{L}_1) \|y\| + 2(L_2 M_2 + \bar{L}_2) \|z\| + 2(M_1 + M_2) F_0 + 4K_0 \\ &\leq 2(LM + \bar{L}) R' + 4MF_0 + 4K_0 \leq R'. \end{aligned}$$

For $(y_1, z_1)(\zeta), (y_2, z_2)(\zeta) \in U \times U$ and any $\zeta \in \Delta$, we have

$$\begin{aligned}
 & |T_1(y_2, z_2)(\zeta) - T_1(y_1, z_1)(\zeta)| \\
 & \leq |A_1 y_2(\zeta) B_1 z_2(\zeta) + C_1 y_2(\zeta) - (A_1 y_1(\zeta) B_1 z_1(\zeta) + C_1 y_1(\zeta))| \\
 & = \left| f_1(\zeta, y_2(\zeta)) \left(\frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y_2(\omega), z_2(\omega)) d\omega \right. \right. \\
 & \quad \left. \left. + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y_2(\omega), z_2(\omega)) d\omega \right) \right) + k_1(\zeta, y_2(\zeta)) \right. \\
 & \quad \left. - f_1(\zeta, y_1(\zeta)) \left(\frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y_1(\omega), z_1(\omega)) d\omega \right. \right. \\
 & \quad \left. \left. + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y_1(\omega), z_1(\omega)) d\omega \right) \right) - k_1(\zeta, y_1(\zeta)) \right| \\
 & = \left| f_1(\zeta, y_2(\zeta)) \left(\frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y_2(\omega), z_2(\omega)) d\omega \right. \right. \\
 & \quad \left. \left. + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y_2(\omega), z_2(\omega)) d\omega \right) \right) + k_1(\zeta, y_2(\zeta)) \right. \\
 & \quad \left. - f_1(\zeta, y_2(\zeta)) \left(\frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y_1(\omega), z_1(\omega)) d\omega \right. \right. \\
 & \quad \left. \left. + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y_1(\omega), z_1(\omega)) d\omega \right) \right) \right. \\
 & \quad \left. + f_1(\zeta, y_2(\zeta)) \left(\frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y_1(\omega), z_1(\omega)) d\omega \right. \right. \\
 & \quad \left. \left. + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y_1(\omega), z_1(\omega)) d\omega \right) \right) \right. \\
 & \quad \left. - f_1(\zeta, y_1(\zeta)) \left(\frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y_1(\omega), z_1(\omega)) d\omega \right. \right. \\
 & \quad \left. \left. + \frac{1}{a_1 + b_1} \left(c_1 - \frac{b_1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y_1(\omega), z_1(\omega)) d\omega \right) \right) - k_1(\zeta, y_1(\zeta)) \right| \\
 & \leq |f_1(\zeta, y_2(\zeta))| \left| \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} (g_1(\omega, y_2(\omega), z_2(\omega)) - g_1(\omega, y_1(\omega), z_1(\omega))) d\omega \right| \\
 & \quad + \frac{|f_1(\zeta, y_2(\zeta))| |c_1|}{|a_1 + b_1|} \left| \int_0^1 (1 - \omega)^{\alpha-1} (g_1(\omega, y_2(\omega), z_2(\omega)) - g_1(\omega, y_1(\omega), z_1(\omega))) d\omega \right| \\
 & \quad + |f_1(\zeta, y_2(\zeta)) - f_1(\zeta, y_1(\zeta))| \left(\left| \frac{1}{\Gamma(\alpha)} \int_0^\zeta (\zeta - \omega)^{\alpha-1} g_1(\omega, y_1(\omega), z_1(\omega)) d\omega \right| \right. \\
 & \quad \left. + \frac{|c_1|}{|a_1 + b_1|} + \frac{|b_1|}{|a_1 + b_1|} \left| \frac{1}{\Gamma(\alpha)} \int_0^1 (1 - \omega)^{\alpha-1} g_1(\omega, y_1(\omega), z_1(\omega)) d\omega \right| \right) + |k_1(\zeta, y_2(\zeta)) - k_1(\zeta, y_1(\zeta))|
 \end{aligned}$$

$$\begin{aligned} &\leq \frac{K'}{\Gamma(\alpha + 1)}(h_1\|y_2 - y_1\| + \bar{h}_1\|z_2 - z_1\|) + \frac{K'}{\Gamma(\alpha + 1)} \cdot \frac{|c_1|}{|a_1 + b_1|}(h_1\|y_2 - y_1\| + \bar{h}_1\|z_2 - z_1\|) \\ &\quad + \|y_2 - y_1\| \left(\frac{K}{\Gamma(\alpha + 1)} + \frac{|c_1|}{|a_1 + b_1|} + \frac{|b_1|}{|a_1 + b_1|} \cdot \frac{K}{\Gamma(\alpha + 1)} \right) + \bar{L}\|y_2 - y_1\| \\ &\leq \mu_1(h_1\|y_2 - y_1\| + \bar{h}_1\|z_2 - z_1\|) + (M + \bar{L})\|y_2 - y_1\|. \end{aligned}$$

Taking supremum over ζ , we get

$$\begin{aligned} &\|T_1(y_2, z_2) - T_1(y_1, z_1)\| \\ &\leq (\mu_1 h_1 + M + \bar{L}) \|y_2 - y_1\| + \mu_1 \bar{h}_1 \|z_2 - z_1\| \\ &\leq (\mu_1 h_1 + M + \bar{L} + \mu_1 \bar{h}_1) \|y_2 - y_1\| + (\mu_1 h_1 + M + \bar{L} + \mu_1 \bar{h}_1) \|z_2 - z_1\| \\ &= (\mu_1(h_1 + \bar{h}_1) + M + \bar{L}) (\|y_2 - y_1\| + \|z_2 - z_1\|). \end{aligned}$$

Similarly, we can obtain that

$$\|T_2(y_2, z_2) - T_2(y_1, z_1)\| \leq (\mu_2(h_2 + \bar{h}_2) + M + \bar{L}) (\|y_2 - y_1\| + \|z_2 - z_1\|).$$

Thus, we get

$$\begin{aligned} &\|T(y_2, z_2) - T(y_1, z_1)\| \\ &\leq 2 (\mu_1(h_1 + \bar{h}_1) + \mu_2(h_2 + \bar{h}_2) + 2M + 2\bar{L}) (\|y_2 - y_1\| + \|z_2 - z_1\|) \\ &\leq (\|y_2 - y_1\| + \|z_2 - z_1\|), \end{aligned}$$

which implies that T is a contraction. By Banach's contraction principle, the operator T has a unique fixed point which is the unique solution of the PBVP (1.1) of the nonlinear coupled FHDS.

5. Examples

In this section, we will present three examples to illustrate the main results.

Example 5.1 Consider the following PBVP

$$\begin{cases} C D_{0+}^{\frac{1}{2}} \left[\frac{y(\zeta) - \frac{1}{32} \sin y(\zeta)}{\frac{1}{4} \sqrt{y^2(\zeta) + 1}} \right] = \zeta + (\zeta - \frac{1}{4})^4 (y(\zeta)^{\rho_1} + z(\zeta)^{\theta_1}), & 0 < \zeta < 1, \\ C D_{0+}^{\frac{1}{3}} \left[\frac{z(\zeta) - \frac{1}{24} \arctan z(\zeta)}{1 + \frac{1}{3} \cos z(\zeta)} \right] = \zeta^2 + (\zeta - \frac{1}{4})^4 (y(\zeta)^{\rho_2} + z(\zeta)^{\theta_2}), & 0 < \zeta < 1, \\ \left[\frac{y(\zeta) - \frac{1}{8} \sin y(\zeta)}{\sqrt{u^2(\zeta) + 1}} \right]_{\zeta=0} + \left[\frac{y(\zeta) - \frac{1}{8} \sin y(\zeta)}{\sqrt{u^2(\zeta) + 1}} \right]_{\zeta=1} = \frac{1}{16}, \\ \left[\frac{z(\zeta) - \frac{1}{24} \arctan z(\zeta)}{1 + \frac{1}{3} \cos z(\zeta)} \right]_{\zeta=0} + \left[\frac{z(\zeta) - \frac{1}{24} \arctan z(\zeta)}{1 + \frac{1}{3} \cos z(\zeta)} \right]_{\zeta=1} = \frac{1}{20}, \end{cases} \tag{5.1}$$

where $0 < \rho_i, \theta_i < 1$ ($i = 1, 2$).

Choose $d_1(\zeta) = \frac{\zeta}{2}$, $d_2(\zeta) = \zeta^2$ and $m_i = n_i = \frac{81}{256}$ ($i = 1, 2$). Then hypotheses (H_1) and (H_2) hold. Therefore, by Theorem 3.1, the PBVP (5.1) has a solution.

Example 5.2 Consider the following PBVP

$$\begin{cases} CD_{0+}^{\frac{1}{2}} \left[\frac{y(\zeta) - \frac{\zeta}{50e\zeta} |y(\zeta)| - \frac{\sin y(\zeta)}{3\zeta^4}}{1 + \frac{1}{10} \cos \zeta |y(\zeta)|} \right] = (\zeta - \frac{1}{2})^4 (y(\zeta)^{\rho_1} + z(\zeta)^{\theta_1}), & 0 < \zeta < 1, \\ CD_{0+}^{\frac{1}{3}} \left[\frac{z(\zeta) - \frac{\arctan \zeta}{40} |z(\zeta)| - \frac{\zeta^4}{7e\zeta}}{\frac{1}{16} (z(\zeta) + \sqrt{z^2(\zeta) + 1})} \right] = (\zeta - \frac{1}{2})^4 (y(\zeta)^{\rho_2} + z(\zeta)^{\theta_2}), & 0 < \zeta < 1, \\ \left[\frac{y(\zeta) - \frac{\zeta}{50e\zeta} |y(\zeta)| - \frac{\sin y(\zeta)}{3\zeta^4}}{1 + \frac{1}{10} \cos \zeta |y(\zeta)|} \right]_{\zeta=0} + \left[\frac{y(\zeta) - \frac{\zeta}{50e\zeta} |y(\zeta)| - \frac{\sin y(\zeta)}{3\zeta^4}}{1 + \frac{1}{10} \cos \zeta |y(\zeta)|} \right]_{\zeta=1} = 0, \\ \left[\frac{z(\zeta) - \frac{\arctan \zeta}{40} |z(\zeta)| - \frac{\zeta^4}{7e\zeta}}{\frac{1}{16} (z(\zeta) + \sqrt{z^2(\zeta) + 1})} \right]_{\zeta=0} + \left[\frac{z(\zeta) - \frac{\arctan \zeta}{40} |z(\zeta)| - \frac{\zeta^4}{7e\zeta}}{\frac{1}{16} (z(\zeta) + \sqrt{z^2(\zeta) + 1})} \right]_{\zeta=1} = 0, \end{cases} \tag{5.2}$$

where $0 < \rho_i, \theta_i < 1$ ($i = 1, 2$).

Choose $d_1(\zeta) = d_2(\zeta) = 0$ and $m_i = n_i = \frac{1}{256}$ ($i = 1, 2$). Then hypotheses (H_1) and (H_2) hold. Therefore, by Theorem 3.1, the PBVP (5.2) has a solution.

Example 5.3 Consider the following PBVP

$$\begin{cases} CD_{0+}^{\frac{1}{2}} \left[\frac{y(\zeta) - \frac{1}{100} \cos y(\zeta)}{1 + \frac{1}{100} \sin y(\zeta)} \right] = \frac{1}{31} + \frac{1}{200} \frac{|y(\zeta)|}{1 + |y(\zeta)|} + \frac{1}{100} \frac{|z(\zeta)|}{1 + |z(\zeta)|}, & 0 < \zeta < 1, \\ CD_{0+}^{\frac{1}{2}} \left[\frac{y(\zeta) - \frac{1}{100} \sin y(\zeta)}{1 + \frac{1}{100} \cos y(\zeta)} \right] = \frac{1}{31} + \frac{1}{100} \frac{|y(\zeta)|}{1 + |y(\zeta)|} + \frac{1}{200} \frac{|z(\zeta)|}{1 + |z(\zeta)|}, & 0 < \zeta < 1, \\ \left[\frac{y(\zeta) - \frac{1}{100} \cos y(\zeta)}{1 + \frac{1}{100} \sin y(\zeta)} \right]_{\zeta=0} + \left[\frac{y(\zeta) - \frac{1}{100} \cos y(\zeta)}{1 + \frac{1}{100} \sin y(\zeta)} \right]_{\zeta=1} = 0, \\ \left[\frac{y(\zeta) - \frac{1}{100} \sin y(\zeta)}{1 + \frac{1}{100} \cos y(\zeta)} \right]_{\zeta=0} + \left[\frac{y(\zeta) - \frac{1}{100} \sin y(\zeta)}{1 + \frac{1}{100} \cos y(\zeta)} \right]_{\zeta=1} = 0. \end{cases} \tag{5.3}$$

It is easy to check that $\alpha = \beta = 0.5$, $a_1 = b_1 = a_2 = b_2 = 1$, $c_1 = c_2 = 0$, $L = \bar{L} = 0.01$, $h_1 = \bar{h}_2 = 0.005$, $\bar{h}_1 = h_2 = 0.01$, $K = 0.0473$, $K' = 1.01$, $M = 0.08$, $\mu_1 = \mu_2 = 1.709$. Then, we have

$$2(\mu_1(h_1 + \bar{h}_1) + \mu_2(h_2 + \bar{h}_2) + 2M + 2\bar{L}) = 0.46254 < 1.$$

Thus, hypotheses (H_1) and (H_3) hold. Therefore, by Theorem 4.1, the PBVP (5.3) has a unique solution.

6. Conclusion

In this paper, we have studied the solvability for the PBVP (1.1) of the nonlinear coupled FHDS. We have presented an existence theorem and a uniqueness result for the PBVP (1.1) of the nonlinear coupled FHDS due to the fixed point theorem in Banach algebra and Banach’s contraction principle. The main results have been well illustrated with the help of three examples.

Acknowledgment

The authors sincerely thank the reviewers for their valuable suggestions and useful comments that have led to the present improved version of the original manuscript.

This research was funded by National Natural Science Foundation of China (61703180, 61803176), and Science and Technology Program of University of Jinan (1008399).

References

- [1] Ali A, Shah K, Khan RA. Existence of solution to a coupled system of hybrid fractional differential equations. *Bulletin of Mathematical Analysis and Applications* 2017; 9(1): 9-18.
- [2] Baleanu D, Khan H, Jafari H, Ali RK, Alipour M. On existence results for solutions of a coupled system of hybrid boundary value problems with hybrid conditions. *Advances in Difference Equations* 2015; 2015 (318): 1-14. doi: 10.1186/s13662-015-0651-z
- [3] Bashiri T, Vaezpour SM, Park C. Existence results for fractional hybrid differential systems in Banach algebra. *Advances in Difference Equations* 2016; 2016 (57): 1-13. doi: 10.1186/s13662-016-0784-8
- [4] Dhage BC. Basic results in the theory of hybrid differential equations with mixed perturbations of second type. *Functional Differential Equations* 2012; 19 (1-2): 87-106.
- [5] Dhage BC, Dhage SB, Buvaneswari K. Existence of mild solutions of nonlinear boundary value problems of coupled hybrid fractional integrodifferential equations. *Journal of Fractional Calculus and Applications* 2019; 10 (2): 191-206.
- [6] Doğan A. Existence results for a class of boundary value problems for fractional differential equations. *Turkish Journal of Mathematics* 2021; 45: 1149-1165. doi: 10.3906/mat-2012-33
- [7] Li S, Yin H, Li L. The solution of a cooperative fractional hybrid differential system. *Applied Mathematics Letters* 2019; 91: 48-54. doi: 10.1016/j.aml.2018.11.008
- [8] Lu H, Sun S, Yang D, Teng H. Theory of fractional hybrid differential equations with linear perturbations of second type. *Boundary Value Problems* 2013; 2013 (23): 1-16. doi: 10.1186/1687-2770-2013-23
- [9] Kilbas AA, Srivastava HH, Trujillo JJ. *Theory and Applications of Fractional Differential Equations*. Amsterdam, Netherlands: Elsevier Science Publisher B.V., 2006.
- [10] Podlubny I. *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications*. Cambridge, MA, USA: Academic Press Inc., 1999.
- [11] Sun S, Zhao Y, Han Z, Li Y. The existence of solutions for boundary value problem of fractional hybrid differential equations. *Communications in Nonlinear Science and Numerical Simulation* 2012; 17 (12): 4961-4967. doi: 10.1016/j.cnsns.2012.06.001
- [12] Xu M, Sun S. Positivity for integral boundary value problems of fractional differential equations with two nonlinear terms. *Journal of Applied Mathematics and Computing* 2019; 59: 271-283. doi: 10.1007/s12190-018-1179-7
- [13] You J, Sun S. On impulsive coupled hybrid fractional differential systems in Banach algebras. *Journal of Applied Mathematics and Computing* 2019; 62: 189-205. doi: 10.1007/s12190-019-01280-z
- [14] Zhao Y. On the existence for a class of periodic boundary value problems of nonlinear fractional hybrid differential equations. *Applied Mathematics Letters* 2021; 121 (107368): 1-6. doi: 10.1016/j.aml.2021.107368
- [15] Zhao Y, Hou X, Sun Y, Bai Z. Solvability for some class of multi-order nonlinear fractional systems. *Advances in Difference Equations* 2019; 2019 (23): 1-10. doi: 10.1186/s13662-019-1970-2
- [16] Zhao Y, Sun S, Han Z, Feng W. Theory of fractional hybrid differential equations. *Computers & Mathematics with Applications* 2011; 62 (3): 1312-1324. doi: 10.1016/j.camwa.2011.03.041
- [17] Zhao Y, Sun S, Han Z, Zhang M. Positive solutions for boundary value problems of nonlinear fractional differential equations. *Applied Mathematics and Computation* 2011; 217 (16): 6950-6958. doi: 10.1016/j.amc.2011.01.103
- [18] Zhao Y, Sun Y, Liu Z, Wang Y. Solvability for boundary value problems of nonlinear fractional differential equations with mixed perturbations of the second type. *AIMS Mathematics* 2020; 5 (1): 557-567. doi: 10.3934/math.2020037