

10-16-2023

## Neonicotinoid resistance in populations of the cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae) in Cotton Plantation Areas of Turkey

CAHİT KAYA

NACİYE SENA ÇAĞATAY

JOHN T. MARGARITPOULOS

JOHN VONTAS

REMZİ ATLIHAN

*See next page for additional authors*

Follow this and additional works at: <https://journals.tubitak.gov.tr/agriculture>



Part of the [Agriculture Commons](#), and the [Forest Sciences Commons](#)

### Recommended Citation

KAYA, CAHİT; ÇAĞATAY, NACİYE SENA; MARGARITPOULOS, JOHN T.; VONTAS, JOHN; ATLIHAN, REMZİ; and GÜZ, NURPER (2023) "Neonicotinoid resistance in populations of the cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae) in Cotton Plantation Areas of Turkey," *Turkish Journal of Agriculture and Forestry*. Vol. 47: No. 5, Article 3. <https://doi.org/10.55730/1300-011X.3114>  
Available at: <https://journals.tubitak.gov.tr/agriculture/vol47/iss5/3>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Agriculture and Forestry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

---

## Neonicotinoid resistance in populations of the cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae) in Cotton Plantation Areas of Turkey

### Authors

CAHİT KAYA, NACİYE SENA ÇAĞATAY, JOHN T. MARGARITOPOULOS, JOHN VONTAS, REMZİ ATLIHAN,  
and NURPER GÜZ

## Neonicotinoid resistance in populations of the cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae) in cotton plantation areas of Türkiye

Cahit KAYA<sup>1</sup> , Naciye Sena ÇAĞATAY<sup>2</sup> , John T. MARGARITOPOULOS<sup>3</sup> , John VONTAS<sup>4</sup> ,  
Remzi ATLIHAN<sup>5</sup> , Nurper GÜZ<sup>6,\*</sup> 

<sup>1</sup>Plant Protection Research Institute, Diyarbakır, Türkiye

<sup>2</sup>Molecular Entomology Laboratory, Department of Plant Protection, Faculty of Agriculture, Ankara University, Ankara, Türkiye

<sup>3</sup>Department of Plant Protection, Institute of Industrial and Fodder Crops, Hellenic Agricultural Organization–DEMETER, Volos, Greece

<sup>4</sup>Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology Hellas, Heraklion, Crete, Greece

<sup>5</sup>Department of Plant Protection, Faculty of Agriculture, Van Yüzüncü Yıl University, Van, Türkiye

<sup>6</sup>Biotechnology Institute, Ankara University, Ankara, Türkiye

Received: 21.02.2022

Accepted/Published Online: 12.04.2023

Final Version: 16.10.2023

**Abstract:** The cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae) is a polyphagous pest that could cause economic crop losses in various crops. Cotton production areas are under insecticide application pressure, and the possibility of insecticide resistance development is higher than in other crops. Chemical insecticides, especially neonicotinoids, are the most common instruments of Integrated Pest Management (IPM) strategies against *A. gossypii*. In this study, the resistance status of *A. gossypii* populations from the largest cotton plantation areas of Türkiye was analyzed. Nine field-collected aphid populations and a susceptible strain were examined in leaf-dip bioassays with three neonicotinoid insecticides. The resistance ratios of bioassays ranged from 22.6 to 82.6 for acetamiprid, 23.5 to 67.3 for imidacloprid, and 1.1 to 20.8 for thiamethoxam. Comparative sequence analysis between susceptible and resistant strains was analyzed to identify known mutations to confer resistance to neonicotinoids. The mean enzyme activity in some populations was significantly higher than in the susceptible strain. The enzyme activity ratios ranged from 1.9 to 3.9 for *CarE* and 1.5 to 3.1 for *GST*. The bioassay data revealed moderate to high resistance levels in acetamiprid and imidacloprid and low to medium levels in thiamethoxam. A partial sequence of the  $\beta 1$  subunit of the *nAChR* in specimens of the populations examined did not reveal any of the *V62I*, *L80S*, and *R81T* and point mutations. The lack of any correlation between the carboxylesterase or glutathione-S-transferase activity and the  $LC_{50}$  values of three insecticides suggested that these two detoxification enzymes were not involved in the resistance levels observed. However, the resistance levels observed in the present study could be attributed to metabolic resistance mechanisms. Another important point is the cross-resistance observed between the neonicotinoids in the present study. Their extensive use, especially in cotton, might select aphid genotypes resistant to more than one neonicotinoid.

**Key words:** *Aphis gossypii*, resistance, cotton, neonicotinoids, nicotinic acetylcholine receptors

### 1. Introduction

The cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae), is one of the most important aphids, that feeds on various host plants in agricultural and nonagricultural areas worldwide. *A. gossypii* is highly polyphagous and can infest over 900 plant species worldwide (Satar et al., 1999; Blackman and Eastop 2000; Begum et al., 2018). The wide host range of *A. gossypii* includes various field and glasshouse crops, especially cotton [*Gossypium hirsutum* L. (Malvaceae)], cucurbits (Cucurbitaceae), citrus (Rutaceae), cocoa [*Theobroma cacao* L. (Malvaceae)], aubergine [*Solanum melongena* L. (Solanaceae)], pepper

[*Capsicum annum* L. (Solanaceae)], potato [*Solanum tuberosum* L. (Solanaceae)], okra [*Abelmoschus esculentus* L. (Malvaceae)], and many ornamental plants. Moreover, it also causes indirect damage by transmitting more than 50 plant viruses (Blackman and Eastop, 2000). *A. gossypii* reproduces parthenogenetically all year round (anholocyclic life-cycle category) in the warmer parts of the world (Blackman and Eastop, 2017). In many countries throughout the world, the aphid causes severe losses in cotton production together with significant qualitative degradation (sticky lint). Due to its economic importance, *A. gossypii* is a target of intense chemical control programs

\* Correspondence: nurperguz@agri.ankara.edu.tr

worldwide.

The cotton aphid is also a key pest in the cotton production areas of Türkiye (Anonymous, 2017). Many active ingredients have been used for the control of *A. gossypii* in Türkiye, such as organophosphates, pyrethroids, and neonicotinoids (Anonymous, 2022a<sup>1</sup>). Insecticides have been a valuable tool for pest control for many decades; however, as with other classes, their extensive and widespread use has led to cases of resistance in many pest species, which can compromise successful control (Bass et al., 2015).

The first case of insecticide resistance in *A. gossypii* was reported against the organophosphate compound Demeton (Ghong et al., 1964). Later resistance to carbamate (Furk et al., 1980) and pyrethroid insecticides (Zil'bermint and Zhuravela, 1984) has been documented. Recent studies showed that the resistance of *A. gossypii* to various insecticides was quite common worldwide (Gubran et al., 1992; Nauen and Elbert, 2003; Wang et al., 2007; Carletto et al., 2010; Herron and Wilson, 2011; Foster et al., 2017). Especially many cases of resistance to neonicotinoids have been well documented in *A. gossypii* clones/samples from various regions, such as the Far East, USA, Australia, and Iran which focused on the underlying resistance mechanisms (Herron and Wilson, 2011; Gore et al., 2013; Koo et al., 2014; Matsuura and Nakamura, 2014; Seyedbrahimi et al., 2015; Hirata et al., 2017; Chen et al., 2020; Wang et al., 2021). Furthermore, target site neonicotinoid resistance in *A. gossypii* and three mutations in  $\beta 1$  subunit of nAChR (L80S, R81T, V62I and K264E) have been found to compromise neonicotinoid efficacy (Koo et al., 2014; Kim et al., 2015; Chen et al., 2017). Specifically, a higher resistance to nitro-substituted neonicotinoids have been shown than to cyano-substituted neonicotinoids in *A. gossypii* (Hirata et al., 2015; 2017). Metabolic resistance has also been reported in some *A. gossypii* genotypes and particularly the involvement of P450 monooxygenases and carboxylesterases (Chen et al., 2015; Wei et al., 2017). Moreover, transcriptome data and RNAi experiments showed an up-regulation in UDP-glucuronosyl transferases might be associated with thiamethoxam tolerance in a resistant strain of *A. gossypii* from China (Pan et al., 2015; 2018).

Cotton production areas in Türkiye are about 432,279 ha, and the production amount is about 2.250.000 t. The Southeastern Anatolian Region is one of the most important cotton production areas in Türkiye, consisting of 58.2% production areas and 56.5% production amount (Anonymous, 2022b<sup>2</sup>). The control of *A. gossypii* is based on chemical insecticides, particularly neonicotinoids (Ulusoy et al., 2018). Farmers and pesticide dealers

stated that neonicotinoid insecticides were the most common insecticide group used in cotton production against *A. gossypii* in Southeastern Anatolian Region, and insensitivity has developed over time against this group (personal interviews). Therefore, the present study aimed to examine resistance levels of the neonicotinoids including acetamiprid, imidacloprid, and thiamethoxam against *A. gossypii* field-collected samples in dose-response bioassays, as well as to determine the presence and frequency of known resistance mechanisms. The data obtained from this study will provide scientific evidence for improving the current Integrated Pest Management (IPM)/Insecticide Resistance Management (IRM) strategies.

## 2. Material and methods

### 2.1. Aphid material

A susceptible clone maintained during 20 years under in vitro conditions was obtained from Bayer (Leverkusen, Germany). Nine samples of *A. gossypii* were collected from cotton fields in various localities of Southeastern Anatolia in Türkiye, during the summer of 2017 (Table 1 and Figure 1). The aphid samples and the susceptible strain were reared on cotton plants [*Gossypium hirsutum* L. (Progen BA 119)] in cages in the laboratory at  $23 \pm 1$  °C,  $65\% \pm 5\%$  relative humidity, and 16: 8 h L: D photoperiod. Field samples were reared for 2–3 generations under laboratory conditions until bioassays.

Adult apterous females of approximately the same age were used in dose-response bioassays and biochemical and molecular analyses. To obtain equal-aged aphid individuals, 50 apterous adult individuals of females were transferred to each leaf of cotton plants at the stage of 5–8 permanent leaves. The plants were kept in plexiglass cages (60 × 60 × 80 cm dimensions) with two ventilation holes covered with fine muslin at three sidewalls of each cage. All adults were removed after 48 h, and the nymphs were reared until adulthood. The newly emerged adults were used either in dose-response bioassays or stored at  $-80$  °C until biochemical and molecular analyses.

### 2.2. Insecticides


Commercial formulations of acetamiprid (Mospilan, SP formulation, %20 Active ingredient, Sumi Agro Türkiye), imidacloprid (Confidor, SC formulation, 350 g/L Active ingredient, Bayer CropScience AG), and thiamethoxam (Actara, SC formulation, 240g/L Active ingredient, Syngenta Crop Protection AG) were used in bioassays.

1 <https://bku.tarimorman.gov.tr/Kullanım/TavsiyeArama>

2 <https://biruni.tuik.gov.tr/medas/?kn=92&locale=tr>

**Table 1.** The field samples of *Aphis gossypii* collection sites and map coordinates in Southeastern Anatolia, Türkiye.

Sample name	Sampling date	Sample location
Susceptible	1998	Germany
Babahaki	02.06.2017	37°51'49.3"N 40°45'16.7"E
Şekerören	06.06.2017	37°44'21.4"N 40°26'05.4"E
Yukarıkılıçtaşı	31.05.2017	37°56'28.9"N 40°15'20.7"E
Develi	12.06.2017	37°15'03.8"N 40°03'11.9"E
Altıntoprak	26.05.2017	37°06'05.3"N 40°35'31.7"E
Çamlıca	14.06.2017	37°10'02.9"N 40°42'20.6"E
Çatalhurma	30.06.2017	36°50'59.6"N 38°56'02.5"E
Yazılıkavak	20.06.2017	37°04'14.4"N 39°04'30.7"E
Yarımsu	03.07.2017	37°08'30.5"N 39°04'20.6"E

**Figure 1.** Sampling sites of *Aphis gossypii* from Southeastern Anatolia, Türkiye.

### 2.3. Dose-response bioassays

The bioassay method IRAC No 019 (IRAC, 2015) was used after modifications. A 4-cm diameter cotton leaf disc was dipped into at least eight different concentrations of water dispersions (including 0.2%, Triton X-100) with insecticides for 5 s. The control solution consisted of ddH<sub>2</sub>O and 0.2% Triton X-100. Leaf discs were placed

upside down in plastic dishes (4-cm diameter) containing 1% agar. The lid had a 2.4 cm diameter hole covered with fine muslin to allow ventilation. A total of 20 adult females were carefully brushed onto the leaf disc. Treated aphids were kept under constant conditions (20 °C, 60 ± 5% RH, and 16:8 h L: D photoperiod). Mortality was assessed after 72 h according to the IRAC (IRAC, 2015) test method. A.

*gossypii* individuals were turned down with a soft brush, and the individuals who could not change their position were considered as dead. The field-collected populations were tested against 1–250 ppm of the acetamiprid and imidacloprid, and 1–400 ppm of the thiamethoxam, while the susceptible population was tested against 0.1–20 ppm of the insecticides. The bioassays were performed with four replicates for each dose. About 720 apterous *A. gossypii* individuals were used in acetamiprid and imidacloprid bioassays and about 800 apterous individuals were used in thiamethoxam bioassays.

#### 2.4. Metabolic enzyme activities

The method described by Stumpf and Nauen (2002) was used with slight modifications to determine carboxylesterase (CarE) activity. Ten apterous adult females per sample were homogenized in 100  $\mu$ L sodium phosphate buffer (0.1 M, pH: 7.5). The homogenate was centrifuged at 10,000 rpm and 4 °C and the supernatant was used as enzyme source. Each well of plate was filled with 25  $\mu$ L of 0.2 M pH: 6.0 sodium phosphate buffer, 25  $\mu$ L of the enzyme source and 200  $\mu$ L of substrate solution. Enzyme activity was read at 23 °C for 10 min at 450 nm wavelength with the Versamax kinetic microplate reader spectrophotometer using SoftMAX PRO (Molecular Devices, CA, United States). Three replicates per sample were performed, and 25  $\mu$ L of 0.1% Triton X-100 containing 0.1 M pH: 7.5 sodium phosphate buffer was used as control.

Glutathione-S-transferase (GST) activity was determined using modified version of the method described by Stumpf and Nauen (2002). Thirty apterous adult females were homogenized in 300  $\mu$ L Tris-HCL buffer (0.05 M, pH 7.5). Homogenate was centrifuged at 10,000 rpm and 4 °C and the supernatant was used as an enzyme source. Each well of plate was filled with 100  $\mu$ L of the enzyme source, 100  $\mu$ L of 0.4 mM CDNB and 100  $\mu$ L of 4 mM GSH. The changes in absorbance were read at 25 °C for 5 min at 340 nm wavelength by using a Versamax kinetic microplate. The enzyme activity was analyzed with SoftMAX PRO software. Trials were replicated three times and nonenzymatic reaction of CDNB and GSH was quantified without homogenate as control.

The Bradford (Bradford, 1976) method was used to calculate the total proteins of each sample using bovine serum albumin (BSA) to build the standard curve. CarE activity was expressed as  $\mu$ mol naphthol/min/mg protein and GST activity as  $\mu$ mol conjugated glutathione/min/mg protein.

#### 2.5. RNA extraction, cDNA synthesis, nAChR $\beta$ 1 subunit sequencing

RNA was extracted from pools of adult *A. gossypii* individuals using the GeneJET RNA isolation kit (Thermo Fisher Scientific, USA) and quantified using NanoDrop® 1000 (Thermo Scientific, USA). One  $\mu$ g RNA was used

for cDNA synthesis using SuperScript III RT (Thermo Scientific, USA) and oligo-dT (Thermo Scientific, USA). PCR reactions (25  $\mu$ L total volume) contained Platinum II Taq DNA Polymerase (5 U/ $\mu$ L; Thermo Scientific, USA), 1  $\mu$ L of cDNA, and 10 pmol of each primer pair. Amplification conditions were as follows: 95 °C for 3 min; 35 cycle of 95 °C for 1 min, 50 °C for 1 min, and 72 °C for 10 min; and a final step at 72 °C for 5 min. A negative control was included in each run with nuclease free water. The primers used for the amplification of partial nAChR  $\beta$ 1 subunits that encompass R81T and V62I point mutations were nAChR  $\beta$ 1-F (5'-CAA CAA ACT AAT CAG ACC TGT CC-3') and nAChR  $\beta$ 1-R (5'-GGC AAG TAG AAC ACT AGC ACG C-3') (Hirata et al., 2015). PCR products were visualized on 1.5% TAE agarose gel electrophoresis. All PCR products were sequenced in both directions with same primers using the Sanger method. Sequencing results were analyzed using Bioedit version 7.2.5 (Hall, 1999). Multiple sequence alignment results were converted to the proteins with MegaX software (Kumar et al., 2018).

#### 2.6. Statistical analyses

The data from dose-response bioassays were analyzed by a probit analysis in order to calculate LC<sub>50</sub> values and the 95% fiducial limits using the software Polo-Plus 2.0 (LeOra Software, Berkeley, CA, USA). Resistance Factors (RF) were calculated by dividing the LC<sub>50</sub> value of each field sample to LC<sub>50</sub> of the susceptible population. In order to determine differences between field samples GST and EST enzyme test, the data were analyzed using a one-way analysis of variance and, the Tukey HSD was used to test differences among sample means for significance. The data did not show any departure from normal distribution (Shapiro-Wilk test,  $p > 0.05$ ), and the criterion of equality of variance was also met (Levene's test,  $p > 0.05$ ). Pearson's correlation analysis was applied to determine any relation between LC<sub>50</sub> values and enzyme activities. The analyses were performed using IBM SPSS Statistics Version 19.0 (IBM SPSS Statistics for Windows, IBM, Armonk, NY, USA).

### 3. Results

#### 3.1. Dose-response bioassays

The most of the bioassay data fit the probit model well ( $p < 0.05$ ). The significant heterogeneity observed in some cases could be attributed to the aphid samples consisting of a mixture of genotypes of different resistance levels.

The dose-response bioassays with acetamiprid and imidacloprid revealed medium to high RF values, ranging from 22.6–82.6 (median: 55.2) and 23.5–67.3 (median: 30.6), respectively. Five of the nine aphid samples examined with acetamiprid and three with imidacloprid showed relatively high RF values ( $>50$ ). By contrast, six out of the nine samples examined with thiamethoxam showed

relatively low RF values (<7.7), and in three samples, the RF values ranged from 11.3 to 20.8 (median: 7.7). The median and the range of the RF values showed that the resistance level was higher in acetamiprid, followed by imidacloprid and thiamethoxam, respectively. Interestingly, the sample from Yarımsu displayed the highest RF value for imidacloprid (RF = 67.3) and thiamethoxam (RF = 20.8) and the second highest value for acetamiprid (RF = 68.8). The most resistant sample to acetamiprid was detected from Şekerören (RF = 82.6) (Table 2, Figure 2).

A positive correlation for the  $LC_{50}$  values among three insecticides was observed. However, Pearson's correlation coefficient values were moderate (0.544 for acetamiprid-imidacloprid, 0.643 for acetamiprid-thiamethoxam, and 0.600 for imidacloprid-thiamethoxam), and the correlation was significant only between acetamiprid and thiamethoxam ( $r = 0.643$ ,  $N = 10$ ,  $p = 0.045$ ).

### 3.2. Metabolic enzyme activities

The field samples showed higher CarE (1.9 to 3.9-fold) and GST activity (1.8 to 3.1-fold) than the susceptible strain. Significant differences among the field samples were observed in CaRb ( $F = 9.769$ ,  $df = 9$ ,  $p < 0.001$ ) and GST ( $F = 21.827$ ,  $df = 9$ ,  $p < 0.001$ ) mean activities, and most of the samples showed significant higher activities than the susceptible strain (Tukey Test,  $p < 0.05$ ). However, no significant correlation has been detected ( $p > 0.378$ ) between CarE or GST activity and  $LC_{50}$  values of three neonicotinoids examined, with Pearson's correlation coefficient values being low (-0.026 to 0.313) (Table 3).

### 3.3. Screening for target site resistance

None of the L80S, R81T, and V62I mutations that confer resistance to neonicotinoids has been detected in partial sequences of the nAChR  $\beta 1$  subunit in *A. gossypii* samples and the susceptible strain (Figure 3).

## 4. Discussion

Due to the extensive use of the neonicotinoid insecticides for controlling cotton aphids in the field, we presented detailed data to understand resistance mechanisms against neonicotinoids of *A. gossypii* samples collected from important cotton growing regions of Türkiye. The bioassay data revealed different resistance ratios; while higher resistance ratio levels were observed for acetamiprid and imidacloprid, lower resistance ratio levels were observed for thiamethoxam. The development of resistance was more evident in acetamiprid than in imidacloprid as suggested by the median RF values (55.2 vs. 30.7) (Table 2). This might be due to the higher selection pressure by acetamiprid since it has a broad target spectrum in cotton, which is often used against *A. gossypii* and other cotton pests in this region (Anonymous, 2018).

In general, the RF values observed in the present study were lower than those in *A. gossypii* resistant strains/

samples with the R81T mutation (Koo et al., 2014; Hirata et al., 2015; Kim et al., 2015; Chen et al., 2017), however, they were in the same order with the values reported in studies where metabolic resistance has been documented (Chen et al., 2015; Seyedebrahimi et al., 2015; Wei et al., 2017; Ulusoy et al., 2018). This is in accordance with the lack of R81T (or V62I) mutations in the examined samples. The resistance levels observed in the present study could be attributed to metabolic resistance mechanisms. Published studies which reported metabolic resistance in *A. gossypii* strains/samples (including from Türkiye), showed the involvement of cytochrome P450 monooxygenases and CarE in neonicotinoid resistance (Seyedebrahimi et al., 2015; Chen et al., 2017; Wei et al., 2017; Ulusoy et al., 2018). In our study, no significant correlation has been detected between CarE and GST activity and  $LC_{50}$  values of three neonicotinoids suggested that these enzymes families are not involved in the resistance observed. A possible resistance mechanism might be attributed to cytochrome P450 monooxygenases, which should be investigated in future studies. Furthermore, recent transcriptome studies presented the importance of phase II enzymes such as Uridine diphosphate (UDP) and glycosyltransferases (UGTs) in resistance of *A. gossypii* to the neonicotinoid insecticides (Pan et al., 2018), which should be addressed, as well.

Another point of discussion is the cross-resistance observed between the neonicotinoids in the present study, as revealed by the positive Pearson's correlation coefficients (0.543–0.643; significant between acetamiprid and thiamethoxam) and a few samples showed the highest RF values in two or three insecticides. Cross resistance among neonicotinoids in *A. gossypii* has been observed in various studies even in populations from Türkiye (Chen et al., 2015; Ulusoy et al., 2018). In Türkiye, there are many registered neonicotinoid insecticides for the control of insect pests in various crops (Anonymous, 2018). Their extensive use in Türkiye, especially in cotton and vegetable crops (hosts of *A. gossypii*), might select aphid genotypes resistant to more than one neonicotinoid which makes the control of cotton aphid difficult.

Interestingly, the sample from Yarımsu locality showed the highest RF value for imidacloprid and thiamethoxam and the second highest value for acetamiprid, which might be due to genetic variation in *A. gossypii* populations (Brévault et al., 2008; Carletto et al., 2009). The selection, under intense chemical control scenarios, and eventually the proliferation of asexual clones with resistant traits such as that observed Yarımsu locality might pose a threat for the chemical control programs in Southeastern Anatolia Region. Since widespread and predominant *A. gossypii* clones (known as super-clones), resistant to insecticides, have been detected in various countries (e.g., Chen et al., 2013).

Table 2. Response of *Aphis gossypii* clones to acetamiprid, imidacloprid, and thiamethoxam in leaf-dip bioassays.

Sample	Acetamiprid			Imidacloprid			Thiamethoxam		
	LC <sub>50</sub> (95% FL)	Slope±SE	RF	LC <sub>50</sub> (95% FL)	Slope±SE	RF	LC <sub>50</sub> (95% FL)	Slope±SE	RF
Susceptible	0.261 (0.20–0.34)	1.060 ± 0.067	–	0.482 (0.38–0.61)	1.076 ± 0.068	–	4.187 (3.43–5.10)	1.301 ± 0.084	–
Babahaki	15.692 (12.69–19.21)	1.388 ± 0.097	60.1	14.776 (11.50–18.75)	1.152 ± 0.095	30.7	61.516 (48.30–79.81)	1.089 ± 0.084	14.7
Şekerören	21.550 (17.31–26.79)	1.306 ± 0.103	82.6	11.336 (8.98–14.09)	1.322 ± 0.101	23.5	25.370 (19.29–33.12)	0.946 ± 0.076	6.1
Yukarıkılıçtaşı	5.910 (4.38–7.67)	1.160 ± 0.110	22.6	22.680 (17.23–30.73)	1.427 ± 0.127	47.1	4.522 (2.90–6.46)	0.834 ± 0.076	1.1
Altıntoprak	7.313 (5.70–9.13)	1.381 ± 0.108	28.1	12.259 (9.71–15.23)	1.296 ± 0.091	25.4	30.940 (24.63–38.86)	1.167 ± 0.084	7.4
Çamlıca	7.849 (6.15–9.77)	1.395 ± 0.108	30.1	14.555 (11.39–18.36)	1.192 ± 0.097	30.2	31.848 (24.43–40.40)	1.246 ± 0.102	7.6
Develi	14.406 (11.58–17.77)	1.484 ± 0.107	55.2	26.171 (21.69–31.43)	1.572 ± 0.103	54.3	32.056 (25.02–41.60)	1.213 ± 0.096	7.7
Çatalhurma	7.461 (5.87–9.31)	1.393 ± 0.118	28.6	13.920 (10.96–17.34)	1.403 ± 0.117	28.9	32.094 (25.80–40.41)	1.294 ± 0.104	7.7
Yarımsu	17.945 (14.87–21.72)	1.694 ± 0.141	68.8	32.449 (25.72–40.95)	1.136 ± 0.083	67.3	87.251 (69.69–111.69)	1.366 ± 0.102	20.8
Yazlıkavak	17.376 (14.44–20.76)	1.721 ± 0.127	66.6	25.661 (21.01–31.13)	1.454 ± 0.099	53.2	47.132 (36.74–61.25)	1.031 ± 0.080	11.3

LC<sub>50</sub> in ppm; FL, fiducial limits; RF, lethal dose ratios (resistance factors).






Figure 2. Logarithmic dose probit lines.

Table 3. Carboxylesterase (in  $\mu\text{mol}$  naphtol/min/mg protein) and glutathione-S-transferase (in  $\mu\text{mol}$  conjugated glutathione/min/mg protein) activities in the field samples and the susceptible strain.

Sample	CarE activity <sup>1</sup>		CarE ratio	GST activity <sup>1</sup>		GST ratio
Susceptible	4.570 ± 0.349	d	1.0	1.795 ± 0.022	c	1.0
Develi	10.616 ± 0.746	c	2.3	5.505 ± 0.111	a	3.1
Altıntoprak	9.553 ± 0.498	cd	2.1	4.813 ± 0.384	a	2.7
Çamlıca	17.053 ± 2.264	ab	3.7	2.830 ± 0.164	bc	1.6
Yukarıkılıçtaşı	9.160 ± 0.699	cd	2.0	2.670 ± 0.140	bc	1.5
Şekerören	8.605 ± 1.076	cd	1.9	3.475 ± 0.384	b	1.9
Babahaki	8.560 ± 1.923	cd	1.9	3.521 ± 0.356	b	2.0
Çatalhurma	17.788 ± 1.521	a	3.9	5.015 ± 0.142	a	2.8
Yarımsu	11.155 ± 0.770	bc	2.4	3.572 ± 0.316	b	2.0
Yazılıkavak	9.068 ± 1.426	cd	2.0	3.159 ± 0.106	b	1.8
<b>Correlations<sup>2</sup></b>						
Acetamiprid	r = -0.026, p = 0.942			r = 0.267, p = 0.456		
Imidacloprid	r = 0.209, p = 0.563			r = 0.313, p = 0.378		
Thiamethoxam	r = 0.188, p = 0.604			r = 0.246, p = 0.493		

<sup>1</sup>Means followed by a different letter differ significantly (Tukey test, p < 0.05).

<sup>2</sup>Pearson's correlation coefficients between enzyme activities and LC<sub>50</sub> values of the examined neonicotinoids.

Our study demonstrated the development of neonicotinoid resistance in *A. gossypii* in Türkiye, which was also confirmed by Ulusoy et al., (2018). Scientists and farmers likewise should be alert and close monitoring of the progress of resistance in the aphid populations is guaranteed. Surveys for the early detection of invaded

genotypes carrying the R81T mutation, which will further deteriorate the situation, are also needed. Measures also should be taken to alleviate the selection pressure from neonicotinoids. Insecticide Resistance Management based on rotation approaches, which use a 'window' or block strategy frequently defined by pest life cycle and




Figure 3. The results of sequence analyze of  $\beta 1$  subunit of *Aphis gossypii* nAChRs.

the crop growth stages should be widespread adopted in cotton grown areas of Türkiye. These strategies involve alternation of insecticides with different mode of actions, population monitoring and the timing of insecticide application according to pest biology and elimination of alternative host plants. Furthermore, natural enemies should be protected and supported, which is one of the most important instruments of IPM and IRM programs.

**Acknowledgement**

We would like to thank Prof. Dr. M. Oktay Gürkan and Assoc. Prof. Dr. Fariba Mehrkhou for help in performing biochemical assay. The study was supported by the Republic of Türkiye, Ministry of Agriculture and Forestry,

General Directorate of Agricultural Research and Policies with a project no. TAGEM/BSAD/16/2/03-03. N.S.Ç., was supported by 100/2000 YÖK PhD Scholarship and TÜBİTAK [The Scientific and Technological Research Council of Türkiye, project number: 1649B031801960] 2211-A PhD Scholarship Programs.

**Conflict of interest**

The authors declare no conflict of interest.

## References

- Anonymous (2017). Pamuk Entegre Mücadele Teknik Talimatı. Republic of Turkey Ministry of Food, Agriculture and Livestock, General Directorate of Agricultural Research and Policies, 108 pages (in Turkish).
- Anonymous (2018). Pamuk Entegre Mücadele Teknik Talimatı. Republic of Turkey Ministry of Food, Agriculture and Livestock, General Directorate of Food and Control Department of Plant Protection Products.(in Turkish).
- Anonymous (2022a). Republic of Turkey Ministry of Food Agriculture and Livestock, General Directorate of Food and Control, Department of Plant Protection Products database, PPP Database Application.
- Anonymous (2022b). The Turkish Statistical Institute database.
- Bass C, Denholm I, Williamson MS, Nauen R (2015). The global status of insect resistance to neonicotinoid insecticides. *Pesticide Biochemistry and Physiology* 121: 78-87. <https://doi.org/10.1016/j.pestbp.2015.04.004>
- Blackman RL, Eastop VF (2017). Taxonomic Issues. In: Van Emden H, Harrington R (editors). *Aphids as Crop Pests*. Harpenden, United Kingdom:CAB International, pp 1-36.
- Blackman RL, Eastop VF (2000). *Aphids on the World's Crops: An Identification and Information Guide*, 2nd Edition . Harpenden, United Kingdom: CAB International.
- Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248-254. [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
- Brévault T, Carletto J, Linderme D, Vanlerberghe-Masutti F (2008). Genetic diversity of the cotton aphid *Aphis gossypii* in the unstable environment of a cotton growing area. *Agricultural and Forest Entomology* 10: 215-223. <https://doi.org/10.1111/J.1461-9563.2008.00377.X>
- Carletto J, Lombaert E, Chavigny P, Brévault T, Lapchin L et al. (2009). Ecological specialization of the aphid *Aphis gossypii* Glover on cultivated host plants. *Molecular Ecology* 18: 2198-2212. <https://doi.org/10.1111/J.1365-294X.2009.04190.X>
- Carletto J, Martin T, Vanlerberghe-Masutti F, Brévault T (2010). Insecticide resistance traits differ among and within host races in *Aphis gossypii*. *Pest Management Science* 66: 301-307. <https://doi.org/10.1002/PS.1874>
- Chen A, Zhang H, Shan T, Shi X, Gao X (2020). The overexpression of three cytochrome P450 genes CYP6CY14, CYP6CY22 and CYP6UN1 contributed to metabolic resistance to dinotefuran in melon/cotton aphid, *Aphis gossypii* Glover. *Pesticide Biochemistry and Physiology* 167: 104601. <https://doi.org/10.1016/j.pestbp.2020.104601>
- Chen X, Li F, Chen A, Ma K, Liang P et al. (2017). Both point mutations and low expression levels of the nicotinic acetylcholine receptor  $\beta 1$  subunit are associated with Imidacloprid resistance in an *Aphis gossypii* (Glover) population from a Bt cotton field in China. *Pesticide Biochemistry and Physiology* 141: 1-8. <https://doi.org/10.1016/j.pestbp.2016.11.004>
- Chen X, Shi X, Wang H, Wang J, Wang et al. (2015). The cross-resistance patterns and biochemical characteristics of an Imidacloprid-resistant strain of the cotton aphid. *Journal of Pesticide Science* 40: 55-59. <https://doi.org/10.1584/jpestics.D14-031>
- Foster SP, Devine G, Devonshire A (2017). Insecticide resistance. In: Emden HF Van, Harrington R (editor) *Aphids as crop pests*. Oxfordshire, United Kingdom ,CAB International, pp. 426-447
- Furk C, Powell DF, Heyd S (1980). Pirimicarb resistance in the melon and cotton aphid, *Aphis gossypii* Glover. *Plant Pathology* 29: 191-196. <https://doi.org/10.1111/J.1365-3059.1980.TB01211.X>
- Ghong K, Zhang G, Zhai G (1964). Resistance of cotton aphid to demeton. *Journal of Entomology* 13: 1-9.
- Gore J, Cook D, Catchot A, Leonard BR, Stewart SD et al. (2013). Cotton Aphid (Heteroptera: Aphididae) Susceptibility to commercial and experimental insecticides in the Southern United States. *Journal of Economic Entomology* 106: 1430-1439. <https://doi.org/10.1603/EC13116>
- Gubran EME, Delorme R, Augé D, Moreau JP (1992). Insecticide resistance in cotton aphid *Aphis gossypii* (Glov.) in the Sudan Gezira. *Pesticide Science* 35: 101-107. <https://doi.org/10.1002/PS.2780350202/FORMAT/PDF>
- Hall T (1999). BioEdit : a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41: 95-98.
- Herron GA, Wilson LJ (2011). Neonicotinoid resistance in *Aphis gossypii* Glover (Aphididae: Hemiptera) from Australian cotton. *Austral Entomology* 50: 93-98. <https://doi.org/10.1111/J.1440-6055.2010.00788.X/FORMAT/PDF>
- Hirata K, Jouraku A, Kuwazaki S, Kanazawa J, Iwasa T (2017). The R81T mutation in the nicotinic acetylcholine receptor of *Aphis gossypii* is associated with neonicotinoid insecticide resistance with differential effects for cyano- and nitro-substituted neonicotinoids. *Pesticide Biochemistry and Physiology* 143: 57-65. <https://doi.org/10.1016/j.pestbp.2017.09.009>
- Hirata K, Kiyota R, Matsuura A, Toda S, Yamamoto A, et al. (2015). Association between the R81T mutation in the nicotinic acetylcholine receptor  $\beta 1$  subunit of *Aphis gossypii* and the differential resistance to Acetamiprid and Imidacloprid. *Journal of Pesticide Science* 40: 25-31. <https://doi.org/10.1584/jpestics.D14-092>
- IRAC (2015). IRAC Susceptibility Test Methods Series.
- Kim JI, Kwon M, Kim GH, Kim SY, Lee SH (2015). Two mutations in nAChR beta subunit is associated with Imidacloprid resistance in the *Aphis gossypii*. *Journal of Asia-Pacific Entomology* 18: 291-296. <https://doi.org/10.1016/j.aspen.2015.01.010>
- Koo HN, An JJ, Park SE, Kim JI, Kim GH (2014). Regional susceptibilities to 12 insecticides of melon and cotton aphid, *Aphis gossypii* (Hemiptera: Aphididae) and a point mutation associated with Imidacloprid resistance. *Crop Prot* 55: 91-97. <https://doi.org/10.1016/j.cropro.2013.09.010>

- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018). MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and Evolution* 35: 1547. <https://doi.org/10.1093/molbev/msy096>
- Markussen MD, Kristensen M (2010). Low expression of nicotinic acetylcholine receptor subunit Mda2 in neonicotinoid-resistant strains of *Musca domestica* L. *Pest Management Science* 66: 1257-1262. <https://doi.org/10.1002/PS.2007>
- Matsuura A, Nakamura M (2014). Development of neonicotinoid resistance in the cotton aphid *Aphis gossypii* (Hemiptera: Aphididae) in Japan. *Applied Entomology Zoology* 49: 535-540. <https://doi.org/10.1007/S13355-014-0289-4/TABLES/2>
- Nauen R, Elbert A (2003). European monitoring of resistance to insecticides in *Myzus persicae* and *Aphis gossypii* (Hemiptera: Aphididae) with special reference to Imidacloprid. *Bulletin of Entomological Research* 93: 47-54. <https://doi.org/10.1079/BER2002215>
- Nauen R, Stumpf N (2002). Fluorometric Microplate Assay to Measure Glutathione S-Transferase Activity in Insects and Mites Using Monochlorobimane. *Analytical Biochemistry* 303: 194-198. <https://doi.org/10.1006/ABIO.2002.5578>
- Pan Y, Peng T, Gao X, Zhang L, Yang C et al. (2015). Transcriptomic comparison of Thiamethoxam-resistance adaptation in resistant and susceptible strains of *Aphis gossypii* Glover. *Comparative Biochemistry and Physiology - Part D* 13: 10-15. <https://doi.org/10.1016/J.CBD.2014.11.001>
- Pan Y, Tian F, Wei X, Wu Y, Gao X et al. (2018). Thiamethoxam resistance in *Aphis gossypii* Glover relies on multiple UDP-glucuronosyltransferases. *Frontiers in Physiology* 9:322. <https://doi.org/10.3389/FPHYS.2018.00322/BIBTEX>
- Qu Y, Chen J, Li C, Wang Q, Guo W et al. (2016). The subunit gene *Lda1* of nicotinic acetylcholine receptors plays important roles in the toxicity of Imidacloprid and Thiamethoxam against *Leptinotarsa decemlineata*. *Pesticide Biochemistry Physiology* 127: 51-58. <https://doi.org/10.1016/J.PESTBP.2015.09.006>
- Seydebrahimi S, Jahromi K, Imani S, Hosseini Naveh V, Hesami S (2015). Characterization of Imidacloprid resistance in *Aphis gossypii* (Glover) (Hemiptera: Aphididae) in Southern Iran. *Turkish Journal of Entomology* 39: 413-423. <https://doi.org/10.16970/TED.67424>
- Shan Y, Yan S, Hong X, Zha J, Qin J (2020). Effect of Imidacloprid on the behavior, antioxidant system, multitaxenobiotic resistance, and histopathology of Asian freshwater clams (*Corbicula fluminea*). *Aquatic Toxicology* 218: 105333. <https://doi.org/10.1016/J.AQUATOX.2019.105333>
- Sparks TC, Nauen R (2015). IRAC: Mode of action classification and insecticide resistance management. *Pesticide Biochemistry and Physiology* 121: 122-128. <https://doi.org/10.1016/J.PESTBP.2014.11.014>
- Ulusoy S, Atakan E, Dinçer S (2018). Neonicotinoid resistance of *Aphis gossypii* Glover, 1877 (Hemiptera: Aphididae) in cotton fields of Çukurova Region, Turkey. *Turkish Journal of Entomology* 42:23-31. <https://doi.org/10.16970/ENTOTED.380010>
- Wang K-Y, Guo Q-L, Xia X-M, Wang HY, Liu TX (2007). Resistance of *Aphis gossypii* (Homoptera: Aphididae) to selected insecticides on cotton from five cotton production regions in Shandong, China. *Journal of Pesticide Science* 32: 372-378. <https://doi.org/10.1584/jpestics.G06-51>
- Wang R, Fang Y, Mu C, Qu C, Li F et al. (2018). Baseline susceptibility and cross-resistance of cycloxaprid, a novel cis-nitromethylene neonicotinoid insecticide, in *Bemisia tabaci* MED from China. *Crop Protection*, 110: 283-287. <https://doi.org/10.1016/J.CROPRO.2017.02.012>
- Wang ZJ, Liang CR, Shang ZY, Yu QT, Xue CB (2021). Insecticide resistance and resistance mechanisms in the melon aphid, *Aphis gossypii*, in Shandong, China. *Pesticide Biochemistry and Physiology* 172: 104768. <https://doi.org/10.1016/J.PESTBP.2020.104768>
- Wei X, Pan Y, Xin X, Zheng C, Gao X et al. (2017). Cross-resistance pattern and basis of resistance in a Thiamethoxam-resistant strain of *Aphis gossypii* Glover. *Pesticide Biochemistry and Physiology* 138: 91-96. <https://doi.org/10.1016/J.PESTBP.2017.03.007>
- Zil'bermints IV, Zhuravela LM (1984). Response of melon and greenhouse aphids to ambush and actellic. *Khimiya v Sel'skom Khozyaistva* (in Russian)3: 37-40.