

1-1-2022

## Solving nonlinear integro-differential equations using numerical method

NEDJEM EDDINE RAMDANI

SANDRA PINELAS

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

RAMDANI, NEDJEM EDDINE and PINELAS, SANDRA (2022) "Solving nonlinear integro-differential equations using numerical method," *Turkish Journal of Mathematics*: Vol. 46: No. 2, Article 22.

<https://doi.org/10.3906/mat-2108-72>

Available at: <https://journals.tubitak.gov.tr/math/vol46/iss2/22>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Solving nonlinear integro-differential equations using numerical method

Nedjem Eddine RAMDANI<sup>1,\*</sup>, Sandra PINELAS<sup>2</sup>

<sup>1</sup>Department of Civil Engineering, Faculty of Technology, University Saad Dahlab, Blida, Algeria,

<sup>2</sup>Peoples' Friendship University of Russia (RUDN University), Moscow, Russian Federation,

Received: 16.08.2021

Accepted/Published Online: 27.09.2021

Final Version: 21.01.2022

**Abstract:** The aim of this paper is to establish conditions for the existence and uniqueness of the solution of a nonlinear integro-differential equation. Moreover, it is to propose a quadrature method in order to find an approximate solution and establish the convergence of the method. We conclude by providing the algorithm and some numerical simulation to confirm our theoretical results.

**Key words:** Nonlinear integro-differential equation, existence and uniqueness, Gauss–Kronrod, quadrature method, backward difference

### 1. Introduction

Nowadays, integral equations (IE) and integro-differential equations (IDE) appear in many areas of real life problems such as: dynamics, mechanics [11, Chapter 2], physics and biology [12, 13] and many mathematicians are interested in studying this type of equations ([2, 4, 5, 7, 15, 16, 20, 21]). The classical methods for solving (IE) and (IDE) are valid for linear equations. On the other hand, for nonlinear equations, they are much more difficult. For instance, existence and uniqueness of the solution is easy to obtain for the linear equations, whereas for the nonlinear equations it is complicated and is one of the most studied problems. See for example [8, 17, 22] and the references therein. Consider the nonlinear integro-differential equation of the form

$$x'(s) = y(s) + \int_{-1}^1 k(s, t, x(t))dt, \quad (1.1)$$

$$x(-1) = 0,$$

where  $x(s)$  is an unknown function that occurs inside and outside the integral sign,  $k(., ., .)$  is the kernel and  $y(s)$  is the source term. In our case, we are interested in the numerical study of Equation (1.1), because of the crucial role that the solution of Equation (1.1) plays in science and engineering [1]. More precisely, we intend to present an iterative method based on the Gauss–Kronrod quadrature ([6, 14, 19]).

The existence and uniqueness of this kind of equations will be shown later.

This paper contains 5 sections and a brief conclusion. In Section 2, conditions for existence and uniqueness are given. In Section 3, we discretize Equation (1.1) using a numerical scheme. Then, we solve Equation (1.1) and prove the vanishing of the error. In Section 4, we introduce a numerical approach based on Gauss–Kronrod

\*Correspondence: nedjemeddine.ramdani@yahoo.com

2010 AMS Mathematics Subject Classification: 45G15; 45J05; 65R20

quadrature. Then we establish the convergence of the approximate solution. In Section 5, we give the algorithm of the method and we provide some numerical examples to illustrate our theoretical results.

## 2. Existence and uniqueness of the solution

The purpose of this section is to obtain the existence and uniqueness of the solution of Equation (1.1) in  $C([-1, 1], \mathbb{R})$ . We therefore apply the contraction principle and Schauder's theorem to reach our results.

### 2.1. Existence of solution in $C([-1, 1], \mathbb{R})$

We consider Equation (1.1) and assume that the following conditions are satisfied:

$$(H_1) \quad k \in C([-1, 1] \times [-1, 1] \times \mathbb{R}, \mathbb{R})$$

$$(H_2) \quad y \in C([-1, 1], \mathbb{R})$$

In addition, suppose that

$$(H_3) \quad \text{there exists } M > 0 \text{ such that}$$

$$|k(s, t, x)| \leq M, \quad \forall s, t \in [-1, 1], \quad x \in \mathbb{R}$$

and we have the following existence theorem

**Theorem 2.1** *Suppose that the conditions  $(H_1) - (H_3)$  are satisfied. Then Equation (1.1) has at least one solution  $x^* \in C([-1, 1], \mathbb{R})$ .*

**Proof** We follow the same steps as [8]. Firstly, we introduce  $D : C([-1, 1], \mathbb{R}) \rightarrow C([-1, 1], \mathbb{R})$ , defined by

$$D(x)(s) := \int_{-1}^s x(w)dw,$$

where  $D$  is a compact operator. Let  $A : C([-1, 1], \mathbb{R}) \rightarrow C([-1, 1], \mathbb{R})$ , such that

$$A(x)(s) := \int_{-1}^s y(w)dw + \int_{-1}^s \int_{-1}^1 k(w, t, x(t))dt dw, \quad s \in [-1, 1] \quad (2.1)$$

Now, by using the Chebyshev norm and the compactness property of  $D$ , we obtain:

$$\|A(x)\|_C \leq \|Dy\|_C + \|D\| \left\| \int_{-1}^1 k(s, t, x(t))dt \right\|_C \leq C_0 [\|y\|_C + 2M], \quad \forall x \in C([-1, 1], \mathbb{R})$$

where  $\|D\| = C_0$ . Thus,  $A(X)$  is equal bounded. In addition, let  $X \subset C([-1, 1], \mathbb{R})$  a bounded subset. Then  $A(X)$  is also a bounded subset. From the uniform continuity of  $k$  with respect to  $s$  (Heine–Cantor theorem), it follows that  $A(X)$  is equal continuous. Therefore,  $A(X)$  is compact subset. Let  $X = \overline{\text{conv}}A(C([-1, 1], \mathbb{R}))$ . On the other hand, by the Ascoli–Arzela's theorem it results that  $A$  is completely continuous operator. At the end, since  $X$  is an invariant subset by  $A$  i.e.  $X \in I(A)$ , it follows that the conditions of the Schauder's theorem are satisfied and the proof is complete.  $\square$

**Remark 2.2** *Equation (1.1) has at least one solution, if we have the compactness property of  $k$  instead of the conditions  $(H_1) - (H_3)$ .*

## 2.2. Existence and uniqueness of the solution in $C([-1, 1], \mathbb{R})$

Suppose now that the following conditions are hold:

( $H_4$ ) there exists  $\gamma > 0$  such that

$$|k(s, t, u) - k(s, t, v)| \leq \gamma|u - v|, \forall s, t \in [-1, 1], u, v \in \mathbb{R}$$

( $H_5$ )  $2C_0\gamma < 1$

and we have the following existence and uniqueness theorem.

**Theorem 2.3** *Suppose that the conditions ( $H_1$ ) – ( $H_2$ ) and ( $H_4$ ) – ( $H_5$ ) are satisfied. Then Equation (1.1) has a unique solution  $x^* \in C([-1, 1], \mathbb{R})$ .*

**Proof** We follow the same steps as [8]. We attach to Equation (1.1) the operator  $A : C([-1, 1], \mathbb{R}) \rightarrow C([-1, 1], \mathbb{R})$ , defined by the relation (2.1). The set of the solutions of integral equation (1.1) coincides with the set of the fixed points of the operator  $A$ . Furthermore, by using ( $H_4$ ) and Chebyshev norm we have

$$\|A(x_1) - A(x_2)\|_C \leq 2C_0\gamma \|x_1 - x_2\|_C.$$

Therefore, by ( $H_5$ ) it results that the operator  $A$  is an  $\nu$ -contraction with the coefficient  $\nu = 2C_0\gamma$ , which completes the proof.  $\square$

## 3. Solving the integro-differential equation

Firstly, we suppose that Equation (1.1) has at least one solution. In this section, we discretize Equation (1.1) by using a numerical method of integration called Gauss–Kronrod quadrature and the backward difference. Then, we study the convergence analysis. Let  $\Gamma = s_1, s_2, \dots, s_n$  be the Gauss points [9] (pp.59, 115), for  $n \geq 1$ .

We have the backward difference:

$$x'(s) = \frac{x(s) - x(s-h)}{h} + O(h),$$

where  $h = s_i - s_{i-1}$  is the discretization step (length between the nodes) with  $h$  assume to goes to 0 as  $n \rightarrow \infty$ ,  $s_i = s$  and  $s_{i-1} = s - h$ . In addition,  $O(h)$  is the truncation error.

Now if  $x^*(s)$  is an analytical solution of Equation (1.1), then for  $\Gamma$  on  $[-1, 1]$ , we have

$$x^*(s_i) - x^*(s_{i-1}) = hy_i + h \int_{-1}^1 k(s_i, t, x^*(t)) dt - O(h^2), \quad i = 1, 2, \dots, n. \quad (3.1)$$

The integral term in Equation (3.1) can be estimated by a numerical integration method. In our case, we use Gauss–Kronrod quadrature [6]. Thus, Equation (3.1) becomes

$$x_i^* - x_{i-1}^* = hy_i + h \sum_{j=1}^n w_j k(s_i, t_j, x_j^*) + hR(\varepsilon) - O(h^2), \quad i = 1, 2, \dots, n, -1 < \varepsilon < 1 \quad (3.2)$$

where  $x_i^* = x^*(s_i)$ ,  $y_i = y(s_i)$ ,  $w_i$  are the weight functions  $i = 1, 2, \dots, n$ , which satisfy the following requirements

$$\sum_{i=1}^n w_i = 2,$$

and  $R(\varepsilon)$  is the Gauss–Kronrod quadrature error [6, Chapter 5], which is expressed as follows, for  $n \geq 1$

$$R(\varepsilon) = \frac{2^{2n+1}(n!)^4}{(2n+1)[(2n)!]^3} k_\varepsilon^{(2n)}(s, \varepsilon, x^*), \quad -1 < \varepsilon < 1.$$

where  $k_\varepsilon^{(2n)}$  is the  $(2n)^{th}$  derivatives with respect to  $\varepsilon$ .

For  $\Gamma$ , we treat a nonlinear system obtained by neglecting the truncation errors of (3.1), as below,

$$\xi_i - \xi_{i-1} = hy_i + h \sum_{j=1}^n w_j k(s_i, t_j, \xi_j), \quad i = 1, 2, \dots, n \tag{3.3}$$

and let  $\xi^* = (\xi_1^*, \xi_2^*, \dots, \xi_n^*)^T$  be the exact solution of the nonlinear system (3.3).

Now, we look for the conditions of vanishing  $\|x^* - \xi^*\|_\infty$ , where  $x^* = (x_1^*, x_2^*, \dots, x_n^*)^T$ .

**Proposition 3.1** *Assume that  $(H_1) - (H_4)$  are hold and for  $n \geq 1$ ,*

(i)  $k \in C^{2n}([-1, 1]^2 \times \mathbb{R}, \mathbb{R}_+^*)$

(ii)  $y \in C([-1, 1], \mathbb{R}_+)$

(iii)  $\gamma < \frac{1-\alpha}{2h}$ , and  $0 < \alpha < 1$  where

$$\alpha = \max_{1 \leq p \leq n} \frac{|x_{p-1}^* - \xi_{p-1}^*|}{|x_p^* - \xi_p^*|}$$

Therefore,

$$\|x^* - \xi^*\|_\infty \leq \frac{1}{1 - (2\gamma h + \alpha)} \left[ \frac{2^{2n+1}(n!)^4 h}{(2n+1)[(2n)!]^3} |k_\varepsilon^{(2n)}(s, \varepsilon, x^*)| - |O(h^2)| \right], \quad -1 < \varepsilon < 1 \tag{3.4}$$

**Proof** Let

$$|x_p^* - \xi_p^*| = \|x^* - \xi^*\|_\infty, \quad 1 \leq p \leq n.$$

By substituting (3.2) and (3.3), we get

$$x_p^* - \xi_p^* = x_{p-1}^* - \xi_{p-1}^* + h \sum_{j=1}^n w_j [k(s_p, t_j, x_j^*) - k(s_p, t_j, \xi_j^*)] + E(\varepsilon, h),$$

where  $E(\varepsilon, h) = hR(\varepsilon) - O(h^2)$ . According to  $(H_4)$  there exist  $\gamma > 0$ , such that

$$|k(s_p, t_j, x_j^*) - k(s_p, t_j, \xi_j^*)| \leq \gamma |x_j^* - \xi_j^*|, \quad j = 1, 2, \dots, n.$$

and by using the above inequality, we obtain

$$\begin{aligned} |x_p^* - \xi_p^*| &\leq |x_{p-1}^* - \xi_{p-1}^*| + \left| h \sum_{j=1}^n w_j [k(s_p, t_j, x_j^*) - k(s_p, t_j, \xi_j^*)] \right| + |E(\varepsilon, h)| \\ &\leq |x_{p-1}^* - \xi_{p-1}^*| + \gamma h \sum_{j=1}^n w_j |x_j^* - \xi_j^*| + |E(\varepsilon, h)| \end{aligned}$$

As  $\sum_{j=1}^n w_j = 2$ . Then,

$$|x_p^* - \xi_p^*| \leq |x_{p-1}^* - \xi_{p-1}^*| + 2\gamma h |x_p^* - \xi_p^*| + |E(\varepsilon, h)|$$

We divide the previous inequality by  $|x_p^* - \xi_p^*|$ , we get

$$\frac{|x_p^* - \xi_p^*|}{|x_p^* - \xi_p^*|} \leq 2\gamma h \frac{|x_p^* - \xi_p^*|}{|x_p^* - \xi_p^*|} + \frac{|x_{p-1}^* - \xi_{p-1}^*|}{|x_p^* - \xi_p^*|} + \frac{|E(\varepsilon, h)|}{|x_p^* - \xi_p^*|}$$

Now, from (i) and (ii) we have

$$x' > 0,$$

which means that  $x$  is strictly increasing, then

$$\max_{1 \leq p \leq n} \frac{|x_{p-1}^* - \xi_{p-1}^*|}{|x_p^* - \xi_p^*|} = \alpha,$$

where  $0 < \alpha < 1$ . Hence,

$$\|x^* - \xi^*\|_\infty \leq \frac{1}{1 - (2\gamma h + \alpha)} \left[ \frac{2^{2n+1}(n!)^4 h}{(2n+1)[(2n!)]^3} \left| k_\varepsilon^{(2n)}(s, \varepsilon, x^*) \right| - O(h^2) \right], \quad -1 < \varepsilon < 1$$

which completes the proof. □

Next, a nonlinear system with a special form is given by (3.3). A numerical approach could now be found to obtain the approximate solution.

#### 4. The numerical approach

The purpose of this section is to find an approximate solution of nonlinear systems, using iterative methods for more details we refer to [19, Chapter 4, 5] and [18, Chapter 2]. As we previously mentioned, the nonlinear system (3.3) has a special form which allows us to approximate its solution using an iterative method.

In linear systems, Gauss–Seidel’s method [18, Chapter 2, Section5] is an effective method for solving such types of systems of equations. Thus, we apply a successive substitution similar to this one and define an iterative process which leads to the vector sequence  $\{\xi^{(k)}\}$ , where  $\xi^{(k)} = (\xi_1^{(k)}, \xi_2^{(k)}, \dots, \xi_n^{(k)})$  and each element of  $\xi^{(k)}$  satisfy the following iteration formula,

$$\xi_i^{(k+1)} - \xi_{i-1}^{(k+1)} = h y_i + h \sum_{j=1}^n w_j k(s_i, t_j, \xi_j^{(k)}), \quad i = 1, 2, \dots, n, \quad k = 0, 1, \dots \tag{4.1}$$

Here, we seek for the conditions that guarantee the convergence of the sequence  $\{\xi^{(k)}\}$ .

Accordingly, we yield the following theorem.

**Theorem 4.1** *Under the same assumption as Proposition 3.1 and for  $1 \leq n < \infty$  the sequence  $\{\xi^{(k)}\}$  comes from (4.1) tends to the exact solution  $\xi^*$  of (3.3), for any arbitrary  $\xi^{(0)}$ .*

**Proof** By substituting (3.3) and (4.1), we get

$$\left(\xi_i^{(k+1)} - \xi_{i-1}^{(k+1)}\right) - \left(\xi_i^* - \xi_{i-1}^*\right) = h \sum_{j=1}^n w_j \left[ k(s_i, t_j, \xi_j^{(k)}) - k(s_i, t_j, \xi_j^*) \right],$$

with  $i = 1, 2, \dots, n, k = 0, 1, \dots$ . From (i) and (ii) we get the increasing property of  $\xi$  and from  $(H_4)$  we get

$$\sum_{j=1}^n w_j \left| k(s_i, t_j, \xi_j^{(k)}) - k(s_i, t_j, \xi_j^*) \right| \leq \gamma \sum_{j=1}^n w_j \left| \xi_j^{(k)} - \xi_j^* \right|$$

Thus

$$\begin{aligned} \left| \xi_i^{(k+1)} - \xi_i^* \right| &\leq \left| \xi_{i-1}^{(k+1)} - \xi_{i-1}^* \right| + 2\gamma h \left\| \xi^{(k)} - \xi^* \right\|_\infty \\ (1 - \beta) \left| \xi_i^{(k+1)} - \xi_i^* \right| &\leq 2\gamma h \left\| \xi^{(k)} - \xi^* \right\|_\infty \\ \left| \xi_i^{(k+1)} - \xi_i^* \right| &\leq \frac{2\gamma h}{1 - \beta} \left\| \xi^{(k)} - \xi^* \right\|_\infty \end{aligned}$$

where  $\beta = \max_{1 \leq i \leq n} \frac{\left| \xi_{i-1}^{(k+1)} - \xi_{i-1}^* \right|}{\left| \xi_i^{(k+1)} - \xi_i^* \right|}$  and  $0 < \beta < 1$ . Now, let  $\lambda = \frac{2\gamma h}{1 - \beta}$ , we have

$$\left\| \xi^{(k+1)} - \xi^* \right\|_\infty \leq \lambda \left\| \xi^{(k)} - \xi^* \right\|_\infty \leq \lambda^2 \left\| \xi^{(k-1)} - \xi^* \right\|_\infty \leq \dots \leq \lambda^k \left\| \xi^{(0)} - \xi^* \right\|_\infty, k = 0, 1, \dots$$

Since, we have  $0 < 2\gamma h < 1 - \beta$  and  $0 < 1 - \beta < 1$ , then  $0 < \lambda < 1$ . Hence,  $\left\| \xi^{(k+1)} - \xi^* \right\|_\infty$  goes to zero, when  $k$  goes to  $\infty$ . The proof is complete. □

## 5. Algorithm of the method and numerical simulations

This section is devoted to numerical tests to verify the efficiency of the method.

### 5.1. Algorithm of the method

The purpose of this section is to define an algorithm for the nonlinear integro-differential equation (1.1) and by taking in consideration the assumptions of Proposition 3.1.

The algorithm is as follows: [3]

**The initial data:**

- Choose any  $\varepsilon > 0$ .
- Let  $\Gamma = \{-1 = s_0 = t_0, s_1 = t_1, \dots, s_{n-1} = t_{n-1}, s_n = t_n = 1\}$  on  $[-1, 1]$ .

- Let  $h = s_i - s_{i-1}$ ,  $i = 1, \dots, n$ , be the step size.
- The initial vector  $\xi^{(0)} = (\xi_0^{(0)}, \xi_1^{(0)}, \dots, \xi_n^{(0)})^T$  and  $\xi_0^{(k)}$  is given.
- Set  $k = 0$  and go to the main steps

**Main steps:**

- (1) Compute  $\xi^{(k+1)}$  using Equation (4.1) and go to the next step.
- (2) Compute the error  $\|\xi^{(k+1)} - \xi^{(k)}\|_\infty$  and go the next step.
- (3) If  $\|\xi^{(k+1)} - \xi^{(k)}\|_\infty \leq \epsilon$ , stop; otherwise, set  $k = k + 1$  and go to step (1).

**5.2. Numerical simulation**

Let  $x^*$  be the exact solution and  $\xi^*$  be the approximate solution using Gauss–Kronrod quadrature with the following error function  $\delta(s_i) = \|x^*(s_i) - \xi^*(s_i)\|_\infty$ .

**5.2.1. Example 1**

Consider the nonlinear integro-differential equations

$$x'(s) = 2s + \int_{-1}^1 4ts \exp(x(t)) dt$$


$$x(-1) = 0,$$

with the exact solution

$$x(s) = s^2 - 1.$$


Where, the initial vector  $\xi^{(0)} = 0$ ,  $\xi_0^{(k)} = (1, \dots, 1)^T$ ,  $n = 199$  and  $\epsilon = 4.4409e - 8$ .

Figure 1 shows the comparison between the exact and the approximate solution and Figure 2 (Figure 3) shows the error function  $\delta(s)$  for  $n = 199$  ( $n = 900$ , respectively).




**Figure 1.** The exact and approximate solution of Example 5.2.1, with  $n = 199$ .





**Figure 2.** The error function for Example 5.2.1, with  $n = 199$ .



**Figure 3.** The error function for Example 5.2.1, with  $n = 900$ .

**5.2.2. Example 2**

Consider the nonlinear Fredholm integral equation of the second kind

$$x'(s) = -\pi \sin(\pi s) + \int_{-1}^1 \sin(t) s x(t)^2 dt$$


$$x(-1) = 0,$$

with the exact


$$x(s) = \cos(\pi s) + 1.$$

Where the initial vector  $\xi^{(0)} = 0$ ,  $\xi_0^{(k)} = 0$ ,  $n = 250$  and  $\epsilon = 4.4409e - 8$ .


Figure 4 shows the comparison between the exact and the approximate solution and Figure 5 (Figure 6) shows the error function  $\delta(s)$  for  $n = 250$  ( $n = 1200$ , respectively).



**Figure 4.** The exact and approximate solution of Example 5.2.2, with  $n = 250$ .



**Figure 5.** The error function for Example 5.2.2, with  $n = 250$ .



**Figure 6.** The error function for Example 5.2.2, with  $n = 1200$ .

### 5.2.3. Example 3


Consider the linear Fredholm integral equation of the second kind

$$x'(s) = 50s^{49} - 58s^{57} + \frac{16}{3009}s^{11} + 9s^8 - \frac{257}{273} + \int_{-1}^1 (s^{11} - t^{11})x(t)dt$$


$$x(-1) = 0,$$

with the initial vector  $\xi^{(0)} = 0$ ,  $\xi_0^{(k)} = 0$ ,  $n = 65$  and  $\epsilon = 4.4409e - 16$ .

Figure 7 shows the comparison between the exact and the approximate solution and Figure 8 shows the error function  $\delta(s)$ .



**Figure 7.** The exact and approximate solution of Example 5.2.3, with  $n = 65$ .



**Figure 8.** The error function for Example 5.2.3, with  $n = 65$ .

**Remark 5.1** *The main reason for Example 5.2.3 is that the proposed method can be applied to both linear and nonlinear equations and the figures show the similarity between exact and approximate solutions.*

**5.2.4. Example 4**

Suppose the following Fredholm integro-differential equation.

$$x'(s) = se^s + e^s - s + \frac{1}{2} \int_0^1 tx(t)dt$$

$$x(0) = 0$$

with exact solution  $x(s) = se^s$ . The errors of the variation of parameters method (VPM) in [10] and Gauss–Kronrod method (GKM) are given in Table .

**Table .** Error analysis of VPM and GKM for Example 5.2.4.

s	GKM	VPM
0	0.002	0.000000000
0.1	0.002213	0.0036943633
0.2	0.002455	0.0145247634
0.3	0.002728	0.0320381598
0.4	0.003038	0.0556708598
0.5	0.003387	0.0847318684
0.6	0.003783	0.1183839981
0.7	0.004229	0.155622457
0.8	0.004733	0.195250603
0.9	0.005302	0.235852468
1	0.005937	0.275761712

**Remark 5.2** *The aim of Example 5.2.4 is to compare the method (GKM) proposed above with (VPM) proposed by Ul Haq et al. [10] and Table shows the efficiency of this new method.*

**6. Conclusion**

In this paper, we have investigated the existence and uniqueness of the solution of a nonlinear integro-differential equation. We have presented a numerical approach named Gauss–Kronrod method with backward difference to obtain an approximate solution and we have proved the vanishing of the error. Furthermore, we have obtained precise results and the numerical simulations in Examples 5.2.1–5.2.4 show the accuracy of the proposed approach. Based on Table , the approximate solution of the proposed method is a better method compared to the VPM.

**Acknowledgment**

The authors are very grateful to the editors and anonymous reviewers for carefully reading the paper and for their comments and suggestions which have improved the paper. The first author has been partially supported by The General Directorate of Scientific Research and Technological Development (DGRSDT) and the second author has been supported by the RUDN University Strategic Academic Leadership Program.

## References

- [1] Arikoglu A, Ozkol I. Solutions of integral and integro-differential equation systems by using differential transform method, *Computers & Mathematics with Applications* 2008; 56 (9): 2411-2417. doi.org/10.1016/j.camwa.2008.05.017
- [2] Babolian E, Shamsavaran A. Numerical solution of nonlinear Fredholm integral equations of the second kind using Haar wavelets, *Journal of Computational and Applied Mathematics* 2009; 225: 87-95. doi.org/10.1016/j.cam.2008.07.003
- [3] Borzabadi AH, Fard OS. A numerical scheme for a class of nonlinear Fredholm integral equations of the second kind, *Journal of Computational and Applied Mathematics* 2009; 232: 449-454. doi:10.1016/j.cam.2009.06.038
- [4] Borzabadi AH, Kamyad AV, Mehne HH. A different approach for solving the nonlinear Fredholm integral equations of the second kind, *Applied Mathematics and Computation* 2006; 173 (2): 724-735. doi:10.1016/j.amc.2005.04.008
- [5] Chebbah H, Mennouni A, Ramdani NE. Numerical Solution of Generalized Logarithmic Integral Equations of the Second Kind by Projections, *Malaysian Journal of Mathematical Sciences* 2018; 12 (3): 349-367.
- [6] David K, Moler C, Nash S. *Numerical methods and software*, Englewood Cliffs, Prentice Hall, 1989.
- [7] Dobritoiu M. An Application of the Admissibility types in b-Metric Spaces, *Transylvanian Journal of Mathematics and Mechanics* 2020; 12 (1): 11-16.
- [8] Dobritoiu M. A Nonlinear Fredholm Integral Equation, *Transylvanian Journal of Mathematics and Mechanics* 2009; 1 (1-2): 25-32.
- [9] Golberg MA. *Numerical Solution of Integral Equations*, Mathematical Concepts and Methods in Science and Engineering, Springer, Boston, MA, 1990. doi:10.1007/978-1-4899-2593-0
- [10] Haq EU, Khan AS, Ali M. An Efficient Approach for Solving the Linear and Nonlinear Integro-differential Equation, *Computational Research Progress in Applied Science & Engineering* 2020; 06 (02).
- [11] Jerri AJ. *Introduction to Integral Equations with Applications*, 2<sup>nd</sup> edition, Clarkson University, 1986.
- [12] Kheybari S, Darvishi MT, Wazwaz AM. A semi analytical algorithm to solve systems of integro-differential equations under mixed boundary conditions, *Journal of Computational and Applied Mathematics* 2017; 317: 72-89. doi: 10.1016/j.cam.2016.11.029
- [13] Kheybari S, Darvishi MT, Wazwaz AM. A semi analytical approach to solve integro-differential equations, *Journal of Computational and Applied Mathematics* 2017; 317: 17-30. doi: 10.1016/j.cam.2016.11.011
- [14] Laurie DP. Calculation of Gauss-Kronrod Quadrature Rules, *Mathematics of Computation* 1997; 66 (219). doi: 10.1090/S0025-5718-97-00861-2
- [15] Mennouni A, Ramdani NE. Collocation method to solve second order Cauchy integro-differential equations, in *Differential and Difference Equations with Applications 2018*; ICDDEA 2017 proceedings, Editors: Pinelas S, Caraballo T, Kloeden P, Graef JR. (Eds.), Springer, ISBN 978-3-319-75647-9. doi: 10.1007/978-3-319-75647-9\_25
- [16] Mennouni A, Ramdani NE, Zennir K. A New Class of Fredholm Integral Equations of the Second Kind with Non Symmetric Kernel: Solving by Wavelets Method, *Boletim da Sociedade de Paranaense de Matemática* 2021; 6 (39): 67-80. doi:10.5269/bspm.41734
- [17] Moradi S, Anjedani MM, Analoei E. On existence and uniqueness of solutions of a nonlinear Volterra-Fredholm integral equation, *International Journal of Nonlinear Analysis and Applications* 2015; 6 (1): 62-68. doi:10.22075/ijnaa.2015.179
- [18] Sauer T. *Numerical Analysis*, 2<sup>nd</sup> Edition, Pearson Education, 2006.
- [19] Stoer J, Bulirsch R. *Introduction to Numerical Analysis*, Springer Verlag, New York, 1993.

- [20] Tuan T, Hong NT. A class of Fredholm equations and systems of equations related to the Kontorovich-Lebedev and the Fourier integral transforms, Turkish Journal of Mathematics 2020; 44: 643-655. doi:10.3906/mat-2001-24
- [21] Wazwaz AM. Nonlinear Fredholm Integro-Differential Equations, Linear and Nonlinear Integral Equations 2011; 517-546. doi: 10.1007/978-3-642-21449-3\_16
- [22] Zhang P, Hao X. Existence and uniqueness of solutions for a class of nonlinear integro-differential equations on unbounded domains in Banach spaces, Advances in Difference Equations 2018; 247. doi: 10.1186/s13662-018-1681-0