

1-1-1998

Crystal and molecular structure of tris(2,6-dichlorophenyl) thiophosphate

Veysel T. YILMAZ

Mustafa ODABAŞOĞLU

R. Alan HOWIE

Follow this and additional works at: <https://journals.tubitak.gov.tr/chem>

 Part of the [Chemistry Commons](#)

Recommended Citation

YILMAZ, Veysel T.; ODABAŞOĞLU, Mustafa; and HOWIE, R. Alan (1998) "Crystal and molecular structure of tris(2,6-dichlorophenyl) thiophosphate," *Turkish Journal of Chemistry*. Vol. 22: No. 4, Article 15. Available at: <https://journals.tubitak.gov.tr/chem/vol22/iss4/15>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Chemistry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Crystal and molecular structure of tris(2,6-dichlorophenyl) thiophosphate

Veysel T. YILMAZ, Mustafa ODABAŞOĞLU

19 Mayıs University, Department of Chemistry,
55139, Kurupelit, Samsun-TURKEY

R. Alan HOWIE

University of Aberdeen, Department of Chemistry,
Meston Walk, Aberdeen, AB9 2UE, UK

Received 14.11.1997


The crystal structure of tris(2,6-dichlorophenyl)thiophosphate was determined by single crystal X-ray diffractometry. The structure consists of isolated molecules and each unit cell contains 8 molecules of the compound. The crystal system is monoclinic with a space group of C2/c. The P atom is bonded to three O atoms and one S atom in a distorted tetrahedron environment.

Introduction

Phosphorus compounds are used in the plastics industry, food manufacture, water treatment, and also as insecticides, detergent and oil additives¹. In the course of studying sulfur-containing organo-phosphorus compounds, Odabaşoğlu and Gümrükçüoğlu synthesized some novel arylthiophosphates and arylphosphorothiochloridates². Single crystal X-ray analyses of some of these crystalline compounds were studied by Odabaşoğlu et al.³ and Büyükgüngör et al.⁴. In this paper, we report the molecular and crystal structure of tris(2,6-dichlorophenyl)thiophosphate from the same family.

Experimental

The compound was synthesized using a method described by Kamai and Bogdanov⁵. 0.15 mol PCl₃ and 0.1 mol 2,6-dichlorophenol were mixed and stirred in an oil bath at 150 °C for 2 hours. After the removal of the excess PCl₃, the synthesized product, tris 2,6-dichlorophenyl phosphite, was purified by vacuum distillation. An equivalent amount of methyl iodide was mixed with tris 2,6-dichlorophenyl phosphite in a sealed glass tube and heated to 100 °C. The reaction product was reacted with S₈ and refluxed in C₆H₆ for 12 hours. The synthesis reaction is formulated below:



Scheme 1.

In order to obtain crystals suitable for X-ray analysis, the compound was recrystallized from a mixture of n-hexane and dioxane (1:1).

Single crystal X-ray diffraction data were obtained from a colourless crystal on a Nicolet P3 four-circle diffractometer with MoK α radiation and graphite monochromator at 298 K. The cell parameters were determined using 14 reflections in the θ range of 7.5-11.4. Data collection used ω scan rates of 1.0 ($I_p < 150$) to 29.3 ($I_p > 2500$) $^\circ \text{ min}^{-1}$ where I_p was the prescan intensity. The scan width was $0.6^\circ \omega$. Background counts were taken at $\mp 1.0^\circ$ in ω from the Bragg angle of each reflection. Atomic scattering factors were taken from the Int. Tables for X-ray crystallography⁶. The RDNIC program⁷ was used for data reductions and SHELXS86⁸ and SHELX76⁹ were used for the structure solution and refinement, respectively. Further detail concerning data collection and refinement are given in Table 1. All non-H atoms were refined anisotropically. H atoms were placed in calculated positions with C-H=0.95Å. The molecular diagram was obtained using the ORTEX graphics program¹⁰. All computations were performed on the SUN SPARCServer (UNIX operating system) of the computing centre of University of Aberdeen.

Table 1. Crystallographic data and refinement parameters for tris(2,6-dichlorophenyl)thiophosphate

Formula	C ₁₈ H ₁₉ Cl ₆ O ₃ PS	Temperature (K)	298
Mw	549.03	Radiation, $\lambda(\text{Å})$	MoK α 0.71073
crystal size(mm)	0.5×0.4×0.4	Range of hkl	\mp 33; 0-9;0-22
Crystal colour	Colourless	Scan type	ω
Crystal system	Monoclinic	$2\theta_{max}(^\circ)$	25
Lattice parameters		Standard reflection	2
a(Å)	28.443(23)	Max. intensity variation(%)	4.4
b(Å)	8.283(7)	Internal	50
c(Å)	18.659(10)	No. of reflections measured	4187
$\alpha(^\circ)$	90	No. of independent reflections	3116
$\beta(^\circ)$	99.54	No. of observed reflections	2075
$\gamma(^\circ)$	90	R	0.103
V(Å ³)	4335(5)	R _w	0.110
Space group	C2/c	S	1.341
Z	8	No. of reflections used	2075
D _C (gcm ⁻¹)	1.682	No. of reflect. paramtr. refined	263
Reflections for	lattice parameters	Weight scheme	1/($\sigma^2 F + 0.000511 F^2$)
Number	14	Max.A/ σ	0.017
θ range ($^\circ$)	7.5-11.4	Pmin-max (e Å ⁻³)	-1.15
μ (cm ⁻¹)	0.98		

Results and Discussion

The shape of the molecule with the atomic numbering scheme is shown in Figure 1. The fractional atomic coordinates and equivalent isotropic displacement parameters for non-H atoms are given in Table 2. The selected bond angles and distances are listed in Table 3.

The structure of the compound is quite similar to those of sulfur-containing organo-phosphorus compounds reported in previous studies^{3,4}. The asymmetric unit of the compound comprises one third of the formula unit. The P and S atoms are located on the threefold axis. The average values of the O-P-S and the O-P-O angles are 116.7(3) $^\circ$ and 101.3(3) $^\circ$, respectively. These values are significantly different from

the ideal tetrahedral angle of 109.5° , but suggest a distorted tetrahedral geometry around the central P atom. Similar observations were reported for tris(2-isopropyl-5-methylphenyl)phosphate³ and tris(o-4-ter-butylphenyl)-phosphate⁴.

Table 2. Fractional atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters (\AA^2 , $\times 10^3$) for non-hydrogen atoms with e.s.d. in parentheses.

$$U_{eq} = (1/3) \sum_i \sum_j j U_{ij} a_i a_j$$

	x/a	y/b	z/c	U_{eq}
P	3695.8(10)	755(5)	3300.4(13)	68.7(13)
C/1	3398.0(15)	-3313(5)	3813.8(19)	115.3(18)
C/2	3037.2(12)	2678(5)	4647.6(17)	96.3(16)
C/3	2670.9(14)	3200(6)	2632(2)	121.3(18)
C/4	4555.7(17)	3924(8)	2954(4)	207(3)
C/5	4288.2(15)	3104(5)	4931(3)	142(2)
C/6	4744.8(19)	-1901(10)	3372(3)	220(3)
S	3728.4(14)	-72(5)	2370(2)	106.5(18)
O1	3246(2)	187(10)	3626(3)	70(2)
O2	3668(3)	2695(10)	3337(4)	71(3)
O3	4092(2)	335(8)	3982(3)	54(2)
C1	3215(4)	-354(19)	4311(5)	65(4)
C2	3253(4)	-1967(19)	4460(6)	74(5)
C3	3189(4)	-2623(19)	5128(7)	86(5)
C4	3078(5)	-1604(23)	5637(7)	95(6)
C5	3.34(4)	36(24)	5519(6)	92(6)
C6	3096(4)	624(18)	4845(6)	75(5)
C7	3607(5)	3689(16)	2725(6)	78(5)
C8	3161(5)	4030(18)	2383(6)	91(5)
C9	3117(8)	5167(21)	1793(8)	121(8)
C10	3511(10)	5839(27)	1605(10)	155(14)
C11	3938(8)	5439(23)	1921(10)	149(11)
C12	4000(6)	4349(19)	2510(7)	110(6)
C13	4557(4)	766(20)	4098(6)	80(5)
C14	4700(4)	2086(16)	4533(9)	95(6)
C15	5174(7)	2584(21)	4663(12)	151(10)
C16	5500(8)	1668(35)	4318(18)	196(17)
C17	5355(8)	407(39)	3940(14)	193(17)
C18	4900(5)	-108(23)	3808(8)	112(7)

The P=S double bond length of the compound is $1.884(4)\text{\AA}$, which is almost the same as that of tris(2-isopropyl-5-methylphenyl)phosphate [$1.886(4)\text{\AA}$]³, but smaller than that of tris(o-4-ter-butylphenyl)-phosphate [$1.908(1)\text{\AA}$]⁴. The mean P-O bond distance is $1.593(7)\text{\AA}$, which is also similar to the reported compounds [$1.588(6)\text{\AA}$]³ and [$1.579(2)\text{\AA}$]⁴ and agrees with that of the P-OH bonds of orthophosphoric acid [1.576\AA]⁶.

The phenyl ring do not exhibit any unexpected geometrical features. However, as a result of the effect of both the aromatic ring and the Cl atom, the mean C-O single bond is $1.373(12)\text{\AA}$, which is somewhat

shorter than a praffinic C-O bond of 1.43Å⁶.




Figure 1. The molecule of tris(2,6-dichlorophenyl)thiophosphate showing atom labels and 20% probability ellipsoids.

Table 3. Selected geometric parameters

Bond distances (Å)			
S-P	1.884(4)	C1-O1	1.371(11)
O1-P	1.577(7)	C7-O2	1.395(12)
O2-P	1.611(9)	C13-O3	1.353(12)
O3-P	1.592(6)	C2-C1	1.466(17)
C2-C11	1.740(13)	C6-C1	1.369(16)
C6-C12	1.743(14)	C3-C2	1.399(16)
C8-C13	1.688(15)	C4-C3	1.347(18)
C12-C14	1.695(16)	C5-C4	1.379(19)
C14-C15	1.710(16)	C6-C5	1.387(15)
C18-C16	1.715(19)		
Bond angles (°)			
O1-P-S	114.4(3)	O3-P-O2	102.5(4)
O2-P-S	114.3(3)	C1-O1-P	129.6(6)
O3-P-S	121.4(3)	C7-O2-P	123.7(7)
O2-P-O1	103.4(4)	C13-O3-P	127.9(6)
O3-P-O1	98.0(3)		

References

1. B.J. Walker, "Organophosphorus Chemistry", Penguin Library of Physical Sciences, Middlesex (1972).
2. M. Odabaşođlu and İ.E. Gümrükçüođlu, Turkish J. Chem., 17, 29-32 (1993).
3. M. Odabaşođlu, İ.E. Gümrükçüođlu, V.T. Yılmaz and R.A. Howie, Turkish J. Chem., 16, 293-298 (1992).
4. O. Büyükgüngör, M. Odabaşođlu, İ.E. Gümrükçüođlu, K. Eichkron and G. Mattern, Acta Cryst., C51, 1207-1209 (1995).
5. G. Kamai and A.P. Bogdanov, Chem. Abstr., 51, 5721 (1957).
6. "International Tables for X-ray Crystallography", Kynoch Press, Birmingham, UK, 1974.
7. R.A. Howie, "RDNIC Data Reduction Program for Nicolet P3 Diffractometer", Univ. of aberdeen, Scotlad, 1980.
8. G.M. Sheldrick, Acta Cryst., A46, 467 (1990).
9. M. Sheldrick, "SHELX76, Program for Crystal Structure Determination", Univ. of Cambridge, England, 1976.
10. P.Mcardle, J. Appl. Cryst., 27, 438 (1994).

CONTENTS

<i>Electron Spectroscopy for Material Characterization</i>	309
Ş. SÜZER	
<i>Fatty Acid Composition and Cholesterol Content of Some Turkish Fish Species</i>	321
S. İMRE, S. SAĞLIK	
<i>Voltammetric Determination of Thioridazine Hydrochloride</i>	325
İ. BİRYOL, S. DERMİŞ	
<i>Polarographic Studies on the Kinetics of the Electrode Reaction of 4-Benzoyl-1,5 Diphenyl Pyrazole-3-Carboxylic Acid</i>	335
M. VERİRŞEN, S. ERTURAN	
<i>Determination of Structure-Toxicity Relationship of Amphiprotic Compounds by Means of the Inhibition of the Dehydrogenase Activity of Pseudomonas putida</i>	341
Ş. GÜL, D. ÖZTÜRK	
<i>Comparison of Different Production Processes for Bioethanol</i>	351
B. ÇAYLAK, F. VARDAR SUKAN	
<i>The Effect of Temperature on the Dynamic Viscosity of Acetone Sunflower-Seed Oil Mixtures</i>	361
H. TOPALLAR, Y. BAYRAK	
<i>Further Studies on the Reaction of Unsaturated Acids with o-Phenylenediamine and 4-Substituted o-Phenylenediamines in Acid Medium</i>	367
M. HASAN, S. MUNAWAR, N. KHAN, Z. MAQBOOL	
<i>Solvent Extraction of La³⁺ with Ethylenediamine-N,N'-ditetradecyl-N,N'-diacetic Acid(EDDAT) in Dichloromethane</i>	373
M. YURDAKOÇ, H. HOŞGÖREN	
<i>Separation of Molybdenum, Vanadium and Nickel by Liquid-Liquid Extraction</i>	379
T. COŞAR, R. ZİYADANOĞULLARI	
<i>Potentiometric Investigation of the Effects of Several Substituents on the Basicity of Benzilidene-o-hydroxyaniline</i>	387
E. KILIÇ, O. ATAKOL, E. CANEL, Z. ALİBEŞEOĞLU, T. GÜNDÜZ, F. KÖSEOĞLU	
<i>Kinetic Studies of the Thermal Decomposition of di- and tri-Organotin Derivatives of 1-Nitroso-2-Naphthol</i>	393
S. ALI, M. H. BHATTI, S. MAHMOOD, M. MAZHAR, M. S. KHALID, N. HUSSAIN	
<i>On the Türker Angles in the Theory of Total π-Electron Energy</i>	399
I. GUTMAN	
<i>Thermal Behaviour of Some Azo Dyes Containing Sterically Hindered and Water-Soluble Groups</i>	403
H. KOCAOKUTGEN, Z. HEREN	
<i>Crystal and molecular structure of tris(2,6-dichlorophenyl) thiophosphate</i>	409
V. T. YILMAZ, M. ODABAŞOĞLU, R. A. HOWIE	

CORRECTIONS

Ayşe KARAGÖLGE, Nazan DEMİR, Yaşar DEMİR & İrfan KÜFREVOĞLU

Apocarbonic Anhydrase-Enzymatic Determination of Zinc in Fruit Juices Volume 21, Number 3, 165 (1997).

1. The following corrections is in order in Table 2: Z_n^{2+} concentration (μ g/100 g fruit)
2. in Figure 3: Absorbance values have to be multiplied by (0.1).