

1-1-2013

## Hepatitis G virus and its prevalence and genotypes in patients with hepatitis B and C in Ahvaz, southwestern Iran

JAVANMARD DAVOD

MAKVANDI MANOOCHHR

HAJIANI ESKANDAR

KHALAFKHANY DAVOD

SAMARBAF ZADEH ALI REZA

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

---

### Recommended Citation

DAVOD, JAVANMARD; MANOOCHHR, MAKVANDI; ESKANDAR, HAJIANI; DAVOD, KHALAFKHANY; and REZA, SAMARBAF ZADEH ALI (2013) "Hepatitis G virus and its prevalence and genotypes in patients with hepatitis B and C in Ahvaz, southwestern Iran," *Turkish Journal of Medical Sciences*: Vol. 43: No. 3, Article 22. <https://doi.org/10.3906/sag-1203-55>

Available at: <https://journals.tubitak.gov.tr/medical/vol43/iss3/22>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Hepatitis G virus and its prevalence and genotypes in patients with hepatitis B and C in Ahvaz, southwestern Iran

Javanmard DAVOD<sup>1</sup>, Makvandi MANOOCHEHR<sup>1,2</sup>, Hajiani ESKANDAR<sup>3</sup>,  
Khalafkhany DAVOD<sup>1</sup>, Samarbaf Zadeh ALI REZA<sup>1,\*</sup>

<sup>1</sup>Department of Virology, Faculty of Medicine, Jondishapour University of Medical Science, Ahvaz, Iran

<sup>2</sup>Infectious and Tropical Disease Research Center, Jondishapour University of Medical Science, Ahvaz, Iran

<sup>3</sup>Division of Gastroenterology and Hepatology, Department of Internal Medicine, Emam Hospital, Jondishapour University of Medical Science, Ahvaz, Iran

Received: 14.03.2012 • Accepted: 14.08.2012 • Published Online: 29.05.2013 • Printed: 21.06.2013

**Aim:** The aim of this study was to determine the prevalence and genotype distribution of hepatitis G virus (HGV, or GB virus C) in patients with hepatitis B and C in Ahvaz, southeastern Iran.

**Materials and methods:** A total of 100 patients were selected; 50 were positive for hepatitis C virus (HCV) and 50 for hepatitis B virus (HBV). HGV status was examined among the HCV- and HBV-positive patients. An enzyme-linked immunosorbent assay (ELISA) kit was used to detect anti-E2 antibody. Nested reverse transcription polymerase chain reaction (RT-PCR) was used for RNA detection from serum samples of chronic hepatitis patients. Sequencing was done to determine genotypes

**Results:** Using the ELISA method, co-infection of HGV with HCV and HBV was determined at 6% and 8%, respectively. RT-PCR showed that co-infection of HGV with HCV and HBV was 22% and 10%, respectively. Based on nucleotide sequencing of PCR products, the predominant genotype of HGV among the samples was 2a.

**Conclusion:** Our study showed that the co-infection rate of HGV in patients with hepatitis B and C was somewhat high and was higher in HCV- than in HBV-infected patients. As our findings and other reports from Iran and neighboring countries indicate, genotype 2 of HGV may be the most common genotype of HGV in Middle Eastern countries.

**Key words:** Hepatitis G virus, hepatitis B virus, hepatitis C virus, prevalence, genotype, reverse transcription polymerase chain reaction

### 1. Introduction

Hepatitis G virus (HGV, or GB virus C) is an enveloped positive-sense, single-stranded virus belonging to the family *Flaviviridae* (1). This virus is transmitted via blood and blood products, like the hepatitis C virus (HCV) (2). HGV has a worldwide distribution among those with multiple sexual partners and drug abusers (3). Based on analysis of the 5' noncoding region (NCR) or E2 sequence, there are 5 major genotypes and a recently identified sixth genotype, all distributed distinctly in different geographical regions (4). Transfusion-related infections are of the life-threatening kind. Hepatitis was the first disease related to transfusion and was first identified by Beeson in 1943 (5). Although HGV belongs to *Flaviviridae*, unlike HCV, it seems that it is not involved in hepatitis (6). The rate of hepatitis B virus (HBV) infection is very high with approximately 350 million carriers in the world, constituting a serious public health

problem (7). HCV infection is the most important cause of chronic liver disease with a total of over 200 million cases of HCV worldwide (8). Co-infection of HGV with HCV is very common, and to a lesser degree with HBV (9). Several studies have shown that its co-infection with human immunodeficiency virus (HIV) is associated with a reduced mortality rate and enhanced reduction of HIV viral load in response to the highly active antiretroviral therapy regime (10,11).

HGV infection could remit because of the disappearance of RNA at the time of antibody production. Most patients that have anti-E2 antibody are negative for HGV RNA, which suggests an inverse correlation of these 2 viral markers (12,13). Anti-E2 antibody can be distinguished in the beginning of infection and also later, along with HGV RNA, suggesting a window period between clearance of RNA and reappearance of anti-E2 antibody (14). There is an enzyme-linked immunosorbent assay (ELISA) test for the

\* Correspondence: alirezasarbaf\_78@hotmail.com

detection of anti-E2 antibody and a reverse transcription polymerase chain reaction (RT-PCR) test for the detection of RNA to diagnose HGV infection (15,16). The aim of this study was to epidemiologically survey prevalence and genotype distribution of HGV in patients with hepatitis B and C in the southwest of Iran. The presence of anti-E2 antibody and RNA in sera was tested, and sequencing was performed to distinguish genotypes of HGV in HBV- and HCV-positive patients.

**2. Materials and methods**

**2.1. Study population**

This study was conducted between July 2011 and December 2012. A total of 100 patients were included in the study. First, the patients with chronic hepatitis B and C were detected, and then their sera were collected from different diagnostic laboratories. Altogether 100 samples were collected, 50 HCV- and 50 HBV-positive. Serum samples of each patient were aliquoted into 2 microtubes. One tube was stored at -20 °C for the ELISA test and the other at -80 °C for RT-PCR.

**2.2. Serological assays**

HCV-Ab and HBs-Ag were detected with ELISA kits (Dia.Pro, Italy) according to manufacturer instructions. Anti-E2 antibody was detected with an IgG ELISA kit (Cusabio, Japan).

**2.3. RNA extraction and cDNA synthesis**

RNA was extracted from 200 µL of each serum using the High Pure Nucleic Acid Kit (Roche, Germany). The extracted RNA was used to synthesize complementary DNA by cDNA preparation kit (Fermentas, Lithuania); the prepared cDNA was stored at -20 °C.

**2.4. PCR amplification of HGV RNA**

Nested PCR using 4 primers designed from the 5' untranslated region (5'-UTR) of HGV (Table 1) was used for HGV RNA amplification (17). In the first amplification step, 5 µL of cDNA sample was amplified in a 25-µL

reaction volume containing 2.5 µL of buffer for PCR reaction (Roche), 0.5 µL dNTP (10 mM), 2.5 µL of MgCl<sub>2</sub>, 0.15 Taq DNA polymerase (5unit, Roche), and 0.3 µL of 2 outer primers located at position 102 for sense and 457 for antisense; sterile water was used as the rest of the volume. The amounts of ingredients in the next round were similar.

**2.5. Polymerase chain reaction**

Thermal conditions of 35 cycles are shown in Table 2. The thermal conditions for the next amplification step were similar to those of the first, containing 5 µL of the first-round PCR product in 25-µL volumes for PCR reaction. For inner primers, the sense was located at position 134 and antisense at 376. The second-round PCR amplicon containing 261 nucleotides was round in 2% agarose gel, considered as positive for HGV RNA (Figure 1). In each round of nested PCR, RNase DNase-free water and control positive (obtained from Keivan Virology Laboratory, Iran) were used, as well as negative serum as the negative control.

**2.6. Phylogenetic analysis**

Sequencing of nested-PCR products was done by Bioneer Company, South Korea. Genotypes of all products were determined by BLAST with online sequences. The phylogenetic tree was constructed in MEGA 5 software (Biodesign Institute, USA) by the boot-strapping method.

**3. Results**

Table 3 shows the results of the serologic and molecular methods of HGV detection in this project. The mean age of patients with HBV was 35.24 ± 13.89 years with 48% male and 52% female, and 30.28 ± 12.03 years for HCV with 68% male and 32% female. These samples were first tested to determine presence of the anti-E2 antibody, with 7 out of 100 positive. Four of 50 patients with hepatitis B (8%) and 3 of 50 patients with hepatitis C (6%) had antibodies against the HGV E-2 envelope protein.


All samples, including those anti-E2 antibody-positive and -negative, were subjected to PCR testing. HGV RNA

**Table 1.** Four primers of 5'-UTR used in HGV PCR reaction and genotyping.

Outer	102-121	Sense	5'-GCCAAAAGGTGGTGGATGGG-3'
	457-477	Antisense	5'-CGGAGCTGGGTGGCCCCATGC-3'
Inner	134-153	Sense	5'-TGGTAGGTCGTAAATCCCGG-3'
	376-395	Antisense	5'-TGGTCCTTGTCAACTCGCCG-3'

**Table 2.** Temperature and time of PCR steps.

Step	Preheating	Denaturation	Annealing	Extension	Final extension
Temperature (°C)	94	94	45	72	72
Time (min)	5	1	1	2	10



**Figure 1.** The image of HGV RNA PCR product on agarose gel painted with safe staining, observed on a Gel Documenter; the first lane is the negative control, lane 2 is a 50-bp DNA ladder, and lanes 4 and 8 are 2 positive samples with 260-bp nucleic acids, while the others are also negative.

was not found in the negative control, but it was found in 16 out of 100 (16%) patients. In the patients with HBV, 5 out of 50 (10%) had HGV RNA, and in the case of HCV, 11 patients (22%) were positive for RNA. No one had HGV RNA and anti-E2 antibody simultaneously. The mean age of patients co-infected with HGV was 34.21 years. However, there was no significant difference between sexes ( $P > 0.05$ ).

The results of sequencing of 5'-UTR showed that all positive samples were genotype 2a, both co-infected with HCV and with HBV. The phylogenetic tree was constructed with the boot-strapping method using MEGA 5 software, showing 6 genotypes of HGV, including 2a, the most prevalent genotype in Iran (Figure 2).

**4. Discussion**

HGV was detected by RT-PCR in 16% of all cases. ELISA testing showed that 8% of HBV-infected patients and 6% of HCV-infected patients were positive for anti-E2 antibody. The values related to co-infection of HGV with HCV and HBV have been reported by other researchers in Iran as varying between 5% and 43% (9,18,19). Nevertheless, co-


infection of HGV has been studied only with HCV, not HBV, in Iran. Ghanbari et al. reported that its prevalence in HCV-infected patients was 43.6% (18), much higher than our findings. Amini et al. reported that 25% of HGV-infected patients also had HCV (9), a value that is close to our findings. Zali et al. reported 40% co-infection of HGV with HCV (19), higher than our findings. This result is consistent with the data reported earlier, but greater than the result of our report (22%). The present study represents the first investigation of HGV infection in patients with chronic hepatitis B living in Ahvaz, Iran. Yang et al. in Taiwan showed that co-infection of HGV with HBV and HCV was 18% and 55%, respectively (14), suggesting that HGV and HCV may share the same route of transmission, in contrast to HBV.

In some studies, co-infection of HGV with HCV and HBV was reported with lower values. Alvarado-Mora et al. in Colombia reported that 5.06% of HBsAg-positive samples were also HGV-positive, while 3.2% of HCV-positive cases were HGV-positive (4), which in the case of HBV samples is close to our findings but much lower than the co-infection rate of HGV and HCV in the present study. Co-infection of HGV with HCV was reported as 12.2% by Hofer et al. (20). In an investigation by Ziaee et al. on hemodialysis patients in Birjand, Iran, 5% of HGV-infected patients also had HCV (21), much lower than our results. Co-infection of this virus with HBV and HCV has been studied by Abo Odeh et al. in the United Arab Emirates, and the figures were 5.7% and 14.3%, respectively (22), close to our findings. Co-infection of HGV with HCV and HBV in Turkey was 7% and 29%, respectively (23), which is dissimilar from the results of the present study. There is a large variation and difference in the prevalence of HGV infection in different geographical regions. This difference may be due to the volume of the population involved in the study, methodology used to detect HGV infection, demographic and clinical features of patients, and different patterns of transmission of virus around the world (blood and blood components, sexual routes, intravenous injection, etc.)

Based on the nucleotide sequence of residue 134-395 of the 5'-UTR region of HGV samples, the genotype 2a was the only genotype of HGV in our city. Our findings are consistent with other studies reported from other

**Table 3.** Demographic data, hepatitis markers, and percentage of HGV RNA and anti-E2 antibody in patients with hepatitis B and C.

Virus	Patients	Male/female (%)	Mean age (years), male/female	Anti-E2 antibody	HGV RNA
HBV	50	48/52	34.12/36.73	4 (8%)	5 (10%)
HCV	50	68/32	32.5 / 28.06	3 (6%)	11 (22%)
Total	100	100	33.31 / 32.3	7 (7%)	16 (16%)



**Figure 2.** Phylogenetic tree of HGV genome constructed by boot-straping method. The GenBank accession numbers are given.

cities of Iran, as they have reported only genotype 2a (18). Reports from other Middle Eastern countries (United Arab Emirates, Turkey, and Saudi Arabia) showed that genotype 2 was the most prevalent genotype in these countries (22,24,25). Although the aim of this research was to identify the prevalence and genotype of HGV (or GB virus C) in patients with hepatitis B and C, to our best knowledge, this report is the first study to survey co-infection of HGV with HBV in the Iran. We found that the co-infection rate of HGV with HBV in Ahvaz was 10% by RT-PCR and 8% by ELISA, and its genotype was 2a.

Our findings show that co-infection of HGV with HCV is much higher than with HBV, and this result is consistent with other studies reported from other countries. High co-infection of HGV with HCV awaits more investigations, however.

**Acknowledgments**

The authors greatly acknowledge the Keivan Virology Laboratory for kindly donating the HGV positive control and Mr Kaydani and Mrs Neisi for their assistance.

## References

- Tucker TJ, Smuts HE. GBV-C/HGV genotypes: Proposed nomenclature for genotypes 1–5. *J Med Virol* 2000; 62: 82–3.
- Tang S, Lai KN. Chronic viral hepatitis in hemodialysis patients. *Hemodial Int* 2005; 9: 169–79.
- Krajden M, Yu A, Braybrook H, Lai AS, Mak A, Chow R et al. GBV-C/hepatitis G virus infection and non-Hodgkin lymphoma: a case control study. *Int J Cancer* 2010; 126: 2885–92.
- Alvarado-Mora MV, Botelho L, Nishiya A, Neto RA, Gomes-Gouvêa MS, Gutierrez MF et al. Frequency and genotypic distribution of GB virus C (GBV-C) among Colombian population with hepatitis B (HBV) or hepatitis C (HCV) infection. *Virology* 2011; 8: 1–7.
- Öner S, Yapıcı G, Şaşmaz CT, Kurt AÖ, Buğdaycı R. Hepatitis B, hepatitis C, HIV, and VDRL seroprevalence of blood donors in Mersin, Turkey. *Turk J Med Sci* 2011; 41: 335–41.
- Zhu WF, Yin LM, Li P, Huang J, Zhuang H. Pathogenicity of GB virus C on virus hepatitis and hemodialysis patients. *World J Gastroenterol* 2003; 9: 1739–42.
- Dikici B, Uzun H, Gözü A, Fidan M. Prevalence of hepatitis B infection among schoolchildren in Southeast Turkey. *Turk J Med Sci* 2009; 39: 289–93.
- Kandemir Ö, Polat G, Saraçoğlu G, Taşdelen B. The predictive role of AST level, prothrombin time, and platelet count in the detection of liver fibrosis in patients with chronic hepatitis C. *Turk J Med Sci* 2009; 39: 857–62.
- Amini S, Mahmoodabadi SA, Lamian S, Joulaie M, Farahani MM. Prevalence of hepatitis G virus (HGV) in high-risk groups and blood donors in Tehran, Iran. *Iran J Public Health* 2005; 34: 41–6.
- Souza IE, Zhang W, Diaz RS, Chaloner K, Klinzman D, Stapleton JT. Effect of GB virus C on response to antiretroviral therapy in HIV-infected Brazilians. *HIV Med* 2006; 7: 25–31.
- Polgreen PM, Xiang J, Chang Q, Stapleton JT. GB virus type C/hepatitis G virus: a non-pathogenic flavivirus associated with prolonged survival in HIV-infected individuals. *Microbes Infect* 2003; 5: 1255–61.
- Tacke M, Kiyosawa K, Stark K, Schlueter V, Ofenloch-Haehnle B, Hess G et al. Detection of antibodies to a putative hepatitis G virus envelope protein. *Lancet* 1997; 349: 318–20.
- Hwang SJ, Lu RH, Chan CY, Wang YJ, Wu JC, Lee SD. The role of hepatitis G virus infection in patients with acute post transfusion hepatitis in Taiwan. *Gastroenterology* 1997; 112: 1260–4.
- Yang J, Dai C, Chuang W, Lin W, Lin Z, Chen S et al. Prevalence and clinical significance of HGV/GBV-C infection in patients with chronic hepatitis B or C. *Jpn J Infect Dis* 2006; 59: 25–30.
- Tan D, Matsumoto A, Conry-Cantilena C, Melpolder JC, Shih JW, Leuther M et al. Analysis of hepatitis G virus (HGV) RNA, antibody to HGV envelope protein, and risk factors for blood donors co-infected with HGV and hepatitis C virus. *J Infect Dis* 1999; 179: 1055–63.
- Ramezani A, Gachkar L, Eslamifard A, Khoshbaten M, Jalilvand S, Adibi L et al. Detection of hepatitis G virus envelope protein E2 antibody in blood donors. *Int J Infect Dis* 2008; 12: 57–61.
- Handajani R, Lusida MI, Suryohudoyo P, Adi P, Setiawan PB, Nidom CA et al. Prevalence of GB virus C/hepatitis G virus infection among various populations in Surabaya, Indonesia, and identification of novel groups of sequence variants. *J Clin Microbiol* 2000; 38: 662–8.
- Ghanbari R, Ravanshad M, Hosseini SY, Yaghobi R, Shahzamani K. Genotyping and infection rate of GBV-C among Iranian HCV-infected patients. *Hepat Mon* 2010; 10: 80–7.
- Zali MR, Mayumi M, Haoufi MM, Nowroozi A. GBV-C infection among patients with hepatitis C virus in the Islamic Republic of Iran: a preliminary report. *East Mediterr Health J* 1999; 5: 1030–4.
- Hofer H, Aydin I, Neumueller-Guber S, Mueller C, Scherzer TM, Staufer K et al. Prevalence and clinical significance of GB virus type C/hepatitis G virus coinfection in patients with chronic hepatitis C undergoing antiviral therapy. *J Viral Hepat* 2011; 18: 513–7.
- Ziaee M, Zarban A, Malekinejad P, Akhbary H. Evaluation of HGV viremia prevalence and its co-infection with HBV, HCV, HIV and HTLV-1 in hemophilic patients of Southern Khorassan, Iran. *Hepat Mon* 2007; 7: 11–4.
- Abu Odeh RO, Al-Moslih MI, Al-Jokhdar MW, Ezzeddine SA. Detection and genotyping of GBV-C virus in the United Arab Emirates. *J Med Virol* 2005; 76: 534–40.
- Akcali S, Sanlidag T, Ozbakkaloglu B. Prevalence of GBV-C/hepatitis G virus viremia among chronic hepatitis B, chronic hepatitis C and hemodialysis patients in Turkey. *Ann Saudi Med* 2006; 26: 68–9.
- Kalkan A, Ozdarendeli A, Bulut Y, Saral Y, Ozden M, Kelestimur N et al. Prevalence and genotypic distribution of hepatitis GB-C/HG and TT viruses in blood donors, mentally retarded children and four groups of patients in eastern Anatolia, Turkey. *Jpn J Infect Dis* 2005; 58: 222–7.
- Al-Ahdal MN, Rezeig MA, Kessie G, Chaudhry F, Al-Shammary FJ. GB virus C/hepatitis G virus infection in Saudi Arabian blood donors and patients with cryptogenic hepatitis. *Arch Virol* 2000; 145: 73–84.