

1-1-2022

On (a,d) -edge local antimagic coloring number of graphs

RAJKUMAR SUNDARAMOORTHY

NALLIAH MOVIRI CHETTIAR

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

SUNDARAMOORTHY, RAJKUMAR and CHETTIAR, NALLIAH MOVIRI (2022) "On (a,d) -edge local antimagic coloring number of graphs," *Turkish Journal of Mathematics*: Vol. 46: No. 5, Article 26.
<https://doi.org/10.55730/1300-0098.3247>

Available at: <https://journals.tubitak.gov.tr/math/vol46/iss5/26>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

On (a, d) -edge local antimagic coloring number of graphs

Rajkumar SUNDARAMOORTHY , Nalliah MOVIRI CHETTIAR* 

Department of Mathematics, School of Advanced Sciences, Vellore Institute of Technology, Vellore, India

Received: 08.12.2021

Accepted/Published Online: 11.05.2022

Final Version: 20.06.2022

Abstract: For any graph $G = (V, E)$, the order and size of G are p and q . A bijection l from $V(G)$ to $\{1, 2, \dots, p\}$ is called (a, d) -edge local antimagic labeling if for any two adjacent edges are not received the same edge-weight (color) and the set of all edge-weights are formed an arithmetic progression $\{a, a + d, a + 2d, \dots, a + (c - 1)d\}$, for some integers $a, d > 0$ and c is the number of distinct colors used in the proper coloring. An edge-weight (color) $w(uv)$ is the sum of two end vertices labels, $w(uv) = f(u) + f(v), uv \in E(G)$. The (a, d) -edge local antimagic coloring number is the least color (edge-weight) used in any (a, d) -edge local antimagic labeling. In the present study, we introduce a new type of labeling and a parameter, also we obtain the (a, d) -edge local antimagic coloring number for paths and wheel graph $W_n, n = 3, 4, 5$. Moreover, we obtain an upper bound of the (a, d) -edge local antimagic coloring number for wheel $W_n, n \geq 6$.

Key words: (a, d) -edge local antimagic labeling, (a, d) -edge local antimagic coloring number, paths and wheel graph

1. Introduction

A graph $G = (V, E)$ is a finite and undirected graph without loops and multiple edges. Let p and q be the number of vertices and edges of G . For graph-theoretic terminology, we refer to Chartrand and Lesniak [2].

Hartsfield and Ringel [5] introduced an antimagic labeling in 1990. An antimagic labeling is a bijection from the set of edges to $\{1, 2, \dots, q\}$ such that all the vertex weights are distinct, where a weight of the vertex v is $w(v) = \sum_{e \in E(v)} f(e)$, and $E(v)$ is the set of edges incident to v . A graph G is called antimagic if G has an antimagic labeling. Several authors studied and obtained several results based on the conjectures, every connected graph is antimagic except K_2 and all trees are antimagic. For further study see in [3–5, 9, 12]. Still, these two conjectures are open. These antimagic labeling and vertex coloring concepts are motivated to introduce a local version of an antimagic labeling.

The local vertex antimagic labeling was introduced by Arumugam et al. [1] in 2017. A local vertex antimagic is a bijection from the set of edges to $\{1, 2, 3, \dots, q\}$ such that for any two adjacent vertices are not received the same weight (color), where the weight of the vertex v is $w(v) = \sum_{e \in E(v)} f(e)$, and $E(v)$ is the set of edges incident to v . The local vertex antimagic chromatic number $\chi_{la}(G)$ is the least number of colors used in any local vertex antimagic labeling of G . They proved some basic results in [1]. For more study see in [4, 6, 8, 11].

*Correspondence: nalliahklu@gmail.com

2020 AMS Mathematics Subject Classification: 05C78, 05C15

The local edge antimagic labeling was introduced by Agustin et al. [7] in 2017. The local edge antimagic labeling is a bijection from the set of vertices to $\{1, 2, 3, \dots, p\}$ such that for any two adjacent edges are not received the same edge-weight (color), where the edge-weight $w(e = uv)$ is the sum of two end vertices labels, $w(uv) = f(u) + f(v), uv \in E(G)$. The local edge antimagic chromatic number $\chi'_{lea}(G)$ is the least number of colors used in any local edge antimagic labeling of G . The following results are obtained in [7].

Theorem 1.1 [7] If $\Delta(G)$ is maximum degree of G , we have $\chi'_{lea}(G) \geq \Delta(G)$.

Theorem 1.2 [7] For a path graph P_n on $n \geq 3$ vertices, we have $\chi'_{lea}(P_n) = 2$.

Theorem 1.3 [7] For a complete graph K_n on $n \geq 3$ vertices, we have $\chi'_{lea}(K_n) = 2n - 3$.

Theorem 1.4 [7] For a wheel graph W_n on $n \geq 3$ vertices, $\chi'_{lea}(W_n) = n + 2$.

Rajkumar and Nalliah [10] found the correct local edge antimagic chromatic number for the Agustin et al.'s [7] result of wheel graph W_n . The correct result is given below. Also, they obtained local edge antimagic chromatic number for fan T_n and friendship F_n graphs as follows.

Theorem 1.5 [10] For the wheel graph W_n on $n \geq 3$ vertices, we have

$$\chi'_{lea}(W_n) = \begin{cases} 5, & \text{if } n = 3, 4 \\ n, & \text{if } n \geq 5. \end{cases}$$

Theorem 1.6 [10] For the fan graph T_n on $n + 1$ vertices, we have

$$\chi'_{lea}(T_n) = \begin{cases} n + 1 & \text{if } n = 2, 3 \\ n & \text{if } n \geq 4. \end{cases}$$

Theorem 1.7 [10] For the friendship graph F_n , we have

$$\chi'_{lea}(F_n) = \begin{cases} 3 & \text{if } n = 1, \\ 2n & \text{if } n \geq 2. \end{cases}$$

2. Main results

The local edge antimagic labeling and the chromatic number of a graph are motivated to study a new type of labeling and a parameter.

Definition 2.1 A (a, d) -edge local antimagic labeling is a bijection l from the set of vertices to $\{1, 2, 3, \dots, p\}$ such that for any two adjacent edges are not received the same edge-weight (color), where an edge-weight $w(e = uv)$ is the sum of two end vertices labels, $w(uv) = f(u) + f(v), uv \in E(G)$ and the set of all edge-weights are formed an arithmetic progression $\{a, a + d, a + 2d, \dots, a + (c - 1)d\}$, for some integers $a, d > 0$ and c is the number of distinct colors are used in the proper coloring. A (a, d) -edge local antimagic labeling is denoted by (a, d) -ELA labeling.

Definition 2.2 The (a, d) -edge local antimagic coloring number of a graph G is the least number of colors used in any (a, d) -edge local antimagic labeling of G and is denoted by $\chi'_{(a,d)\text{-ela}}(G)$.

Example 2.3 From Theorem 1.5[10], we get the local edge antimagic chromatic number of W_5 is 5 and we proved there is no $(a, 1)$ -ELA labeling with 5-colors in Theorem 2.14. Hence, the graph W_5 has $(4, 1)$ -ELA labeling with 6-colors, which is given in Figure 1.




Figure 1. The $(4, 1)$ -ELA labeling of W_5 with 6-colors.

Observation 2.4 If the graph G admits an (a, d) -ELA labeling f , then $\chi'_{(a,d)\text{-ela}}(G) \geq \chi'_{lea}(G) \geq \chi'(G) \geq \Delta(G)$.

Observation 2.5 If the graph G admits an (a, d) -ELA labeling with c -colors, then $d \leq \frac{2p-4}{c-1}$.

Proof Let G be a graph of order p . If G admits an (a, d) -ELA labeling f with c -colors, then the edge-weights are $a, a + d, \dots, a + (c - 1)d$. Then the maximum possible edge-weight of an edge e is $w(e) \leq p + p - 1$. The minimum possible edge-weight of an edge e is $a \geq 3$. Therefore, $a + (c - 1)d \leq 2p - 1$, which implies, we get $d \leq \frac{2p-4}{c-1}$. \square

Observation 2.6 Let G be a graph with order p and size q . If G admits an (a, d) -ELA labeling f with c -colors then

$$\sum_{v \in V(G)} \deg(v)f(v) = \sum_{i=1}^c a_i w_i, \quad \text{where} \quad \sum_{i=1}^c a_i = |E(G)| = q. \tag{2.1}$$

Now, we consider a path graph P_n . From Observation 2.5, if the (a, d) -ELA coloring number of P_n with 2-colors, then $d \leq 2n - 4$. The following theorems gives the (a, d) -ELA labeling with 2-colors, when $d = 1$ and 2.

Theorem 2.7 For a path graph P_n on $n \geq 3$ vertices. Then $\chi'_{(n+1,1)\text{-ela}}(P_n) = 2$.

Proof Let $V(P_n) = \{v_i, 1 \leq i \leq n\}$ and $E(P_n) = \{v_i v_{i+1}, 1 \leq i \leq n - 1\}$. Then $|V(P_n)| = n$ and $|E(P_n)| = n - 1$. Now, define a bijection $f_1 : V(P_n) \rightarrow \{1, 2, \dots, n\}$ by

$$f_1(v_i) = \begin{cases} \frac{i+1}{2}, & \text{if } i \text{ is odd} \\ \frac{2n+2-i}{2}, & \text{if } i \text{ is even.} \end{cases}$$

The edge-weights of P_n are

$$w_1(v_i v_{i+1}) = \begin{cases} n + 1, & \text{if } i \text{ is odd} \\ n + 2, & \text{if } i \text{ is even.} \end{cases}$$

It is easy to identify that f_1 proves a proper edge coloring of P_n and hence $\chi'_{(n+1,1)\text{-ela}}(P_n) \leq 2$. Since $\chi'_{lea}(P_n) = 2$, it follows, we get $\chi'_{(n+1,1)\text{-ela}}(P_n) \geq 2$. Hence $\chi'_{(n+1,1)\text{-ela}}(P_n) = 2$. \square

Example 2.8 The graph P_7 has $(8, 1)$ -ELA labeling with 2-colors, which is given in Figure 2.



Figure 2. The $(8, 1)$ -ELA labeling of P_7 with 2-colors.

Theorem 2.9 For a path graph P_n on $n \geq 3$ vertices. Then $\chi'_{(a,2)\text{-ela}}(P_n) = 2$. Moreover,

$$a = \begin{cases} n, & n \text{ is odd} \\ n + 1, & n \text{ is even.} \end{cases}$$

Proof Let $V(P_n) = \{v_i, 1 \leq i \leq n\}$ and $E(P_n) = \{v_i v_{i+1}, 1 \leq i \leq n - 1\}$. Then $|V(P_n)| = n$ and $|E(P_n)| = n - 1$.

Case(i) n is odd.

Now, define a bijection $f_2 : V(P_n) \rightarrow \{1, 2, \dots, n\}$ by

$$f_2(v_i) = \begin{cases} i, & \text{if } i \text{ is odd} \\ n + 1 - i, & \text{if } i \text{ is even.} \end{cases}$$

The edge-weights of P_n are

$$w_2(v_i v_{i+1}) = \begin{cases} n, & \text{if } i \text{ is odd} \\ n + 2, & \text{if } i \text{ is even.} \end{cases}$$

Case(ii) n is even.

Now, define a bijection $f_3 : V(P_n) \rightarrow \{1, 2, \dots, n\}$ by

$$f_3(v_i) = \begin{cases} i, & \text{if } i \text{ is odd} \\ n + 2 - i, & \text{if } i \text{ is even.} \end{cases}$$

The edge-weights of P_n are

$$w_3(v_i v_{i+1}) = \begin{cases} n + 1, & \text{if } i \text{ is odd} \\ n + 3, & \text{if } i \text{ is even.} \end{cases}$$

It is easy to identify that f_2 and f_3 proves a proper edge coloring of P_n and hence $\chi'_{(a,2)\text{-ela}}(P_n) \leq 2$. Since $\chi'_{lea}(P_n) = 2$, it follows, we get $\chi'_{(a,2)\text{-ela}}(P_n) \geq 2$. Hence $\chi'_{(a,2)\text{-ela}}(P_n) = 2$. \square

Example 2.10 The graphs P_7 and P_8 admit $(a, 2)$ -ELA labeling with 2-colors, where $a = 7$ and 9, which is given in Figures 3 and 4.



Figure 3. The $(7, 2)$ -ELA labeling of P_7 with 2-colors.




Figure 4. The $(9, 2)$ -ELA labeling of P_8 with 2-colors.

Theorem 2.11 There is no (a, d) -ELA labeling of the path P_n with 2-colors, where $d \geq 3$.

Proof Suppose $\chi'_{(a,d)\text{-ela}}(P_n) = 2$, where $d \geq 3$. Then there exists a (a, d) -ELA labeling f with 2-colors. The minimum possible edge-weight is $a \geq 3$ and the maximum possible edge-weight is $a + (2 - 1)d \leq n + n - 1$. Therefore, we get $3 \leq a \leq 2n - 1 - d$. If $d \geq 3$ then the edge-weights are a and $a + d$. If $n \geq 3$ is odd then the edge-weights a and $a + d$ must be used $\frac{n-1}{2}$ times and hence an edge-weight $w = n$ or $n - 1$ has only $\frac{n-1}{2}$ possibilities of two elements sets with their sum is w . Therefore, there is no edge with weight $a + d, d \geq 3$, which is a contradiction.

If n is even then the edge-weights a and $a + d$ must be used $\frac{n}{2}$ and $\frac{n}{2} - 1$ times and hence an edge-weight $w = n + 1$ has only $\frac{n}{2}$ possibilities of two elements sets with their sum is w . Similarly, an edge weight $w' = n - 2$ or $n - 1$ or n or $n + 2$ or $n + 3$ is only have $\frac{n}{2} - 1$ possibilities of two elements sets with their sum is w' . Therefore, there is no edge with weight $a + d, d \geq 3$, which is a contradiction. \square

Theorem 2.12 Let G be the graph of wheel graph $W_n, n \geq 5$, fan graph $T_n, n \geq 4$ and friendship graph $F_n, n \geq 2$ with p vertices. If the graph G admits an (a, d) -ELA labeling with $c = \chi'_{lea}(G)$ -colors, then $d \leq 2$.

Proof Let $G \cong W_n$ and T_n be the wheel and fan graph on $p = n + 1$ vertices. From Theorem 1.5 [10] and Theorem 1.6 [10], we get $c = \chi'_{lea}(G) = n$ and Observation 2.5, we get $d \leq \frac{2(n+1)-4}{n-1} = 2$. Let G be the friendship graph F_n with $p = 2n + 1$. From Theorem 1.7 [10], we get $c = \chi'_{lea}(F_n) = 2n$, and Observation 2.5, we get $d \leq \frac{2(2n+1)-4}{2n-1} = 2$. \square

Theorem 2.13 For a wheel graph W_n on $n + 1$ vertices, where $n = 3, 4$. Then $\chi'_{(a,1)\text{-ela}}(W_n) = 5$, where

$$a = \begin{cases} 3, & n = 3 \\ 4, & n = 4. \end{cases}$$

Proof For $n = 3$, clearly, $W_3 \cong K_4$, it follows from Theorem 1.3 [7], we get $\chi'_{(3,1)\text{-ela}}(W_3) = 5$. For $n = 4$, let $V(W_4) = \{c, v_1, v_2, v_3, v_4\}$. Now, define a labeling $f_4 : V(W_4) \rightarrow \{1, 2, 3, 4, 5\}$ by $f_4(c) = 3, f_4(v_1) = 1, f_4(v_2) = 5, f_4(v_3) = 2$, and $f_4(v_4) = 4$. Then the edge-weights are $w_4(cv_1) = 4, w_4(cv_2) = 8, w_4(cv_3) = 5, w_4(v_1v_2) = 6, w_4(v_1v_3) = 3, w_4(v_1v_4) = 5, w_4(v_2v_3) = 7$, and $w_4(v_2v_4) = 6, w_4(v_3v_4) = 6$. It is easy to identify that f_4 proves a

proper edge coloring of W_n and hence $\chi'_{(4,1)\text{-ela}}(W_4) \leq 5$. From Theorem 1.5 [10], we get $\chi'_{(4,1)\text{-ela}}(W_4) \geq 5$. Thus $\chi'_{(4,1)\text{-ela}}(W_4) = 5$. □

Theorem 2.14 *Let W_5 be the wheel graph on 6 vertices. Then there is no $(a, 1)$ -ELA labeling of W_5 with 5-colors.*

Proof Suppose the graph W_5 has $\chi'_{(a,1)\text{-ela}}(W_5) = 5$. Then there exists an $(a, 1)$ -ELA labeling f with 5-colors. The minimum possible edge-weight is $a \geq 3$ and the maximum possible edge-weight is $a + (5 - 1)1 \leq 11$, which implies $a \leq 7$. Hence $3 \leq a \leq 7$. From Equation (2.1), we get

$$3 \left[\frac{(6)(7)}{2} - i \right] + 5i = \sum_{i=1}^5 a_i w_i, \quad \text{where } f(c) = i \quad \text{and} \quad \sum_{i=1}^5 a_i = 10.$$

This implies, we get the equation

$$63 + 2i = \sum_{i=1}^5 a_i w_i, \quad \text{where } f(c) = i \quad \text{and} \quad \sum_{i=1}^5 a_i = 10. \tag{2.2}$$

Since $3 \leq a \leq 7$, it follows that, there are five possible edge-weight sets: $W'_1 = \{3, 4, 5, 6, 7\}$, $W'_2 = \{4, 5, 6, 7, 8\}$, $W'_3 = \{5, 6, 7, 8, 9\}$, $W'_4 = \{6, 7, 8, 9, 10\}$ and $W'_5 = \{7, 8, 9, 10, 11\}$. Since the edge-weights 3, 4, 10 and 11 are only one possible set of two elements set, the edge-weights 5, 6, 8 and 9 are two possible sets of two elements sets, and an edge-weight 7 is three possible sets of two elements sets.

Case(i) $a = 3$ and 7

If $a = 3$ then the edge-weight set is $W'_1 = \{3, 4, 5, 6, 7\}$ and hence $a_1 = a_2 = 1$, $a_3 = a_4 = 2$, $a_5 = 3$. Therefore, we get $\sum_{i=1}^5 a_i = 9$, which is a contradiction. A similar contradiction arise for the case $a = 7$.

Case(ii) $a = 4$

Then the edge-weight set is $W'_2 = \{4, 5, 6, 7, 8\}$. Now, we substitute the values $w_1 = 4, w_2 = 5, w_3 = 6, w_4 = 7, w_5 = 8$ and $a_1 = 1, a_2 = a_3 = a_5 = 2, a_4 = 3$ in Equation (2.2), we get $63 + 2i = 1(4) + 2(5) + 2(6) + 3(7) + 2(8)$ with $\sum_{i=1}^5 a_i = 10$. This implies, we get $i = 0$, which is a contradiction.

Case(iii) $a = 5$

Then the edge-weight set is $W'_3 = \{5, 6, 7, 8, 9\}$. Since $\sum_{i=1}^5 a_i = 10$, it follows that, the possible 5-tuples $(a_1, a_2, a_3, a_4, a_5)$ are $(1, 2, 3, 2, 2)$, $(2, 1, 3, 2, 2)$, $(2, 2, 2, 2, 2)$, $(2, 2, 3, 1, 2)$ and $(2, 2, 3, 2, 1)$. If $(a_1, a_2, a_3, a_4, a_5) = (1, 2, 3, 2, 2)$, $(2, 2, 2, 2, 2)$ and $(2, 2, 3, 2, 1)$ then we substitute the values $w_1 = 5, w_2 = 6, w_3 = 7, w_4 = 8, w_5 = 9$ and the corresponding 5-tuples values of a_i in Equation (2.2), we get i is not an integer, which is a contradiction.

If $(a_1, a_2, a_3, a_4, a_5) = (2, 1, 3, 2, 2)$ then we substitute the values $w_1 = 5, w_2 = 6, w_3 = 7, w_4 = 8, w_5 = 9$ and the corresponding 5-tuples values of a_i in Equation (2.2), we get $i = 4$ and hence the central vertex label $f(c) = 4$ and other vertices are received the labels except 4. Let $e = cv$. If $f(v) = 6$ then $w(e) = 10 \notin W'_3$, which is a contradiction. A similar contradiction arise for the case $(a_1, a_2, a_3, a_4, a_5) = (2, 2, 3, 1, 2)$. □

Problem 2.15 Does there exist a $(a, 1)$ -ELA coloring number with n -colors for the wheel graph $W_n, n \geq 6$?

Theorem 2.16 For a wheel graph W_n on $n + 1$ vertices, where $n \geq 5$. Then $\chi'_{(a,1)-ela}(W_n) \leq n + 1$, where

$$a = \begin{cases} \frac{n+3}{2}, & \text{if } n \text{ is odd} \\ \frac{n}{2} + 2, & \text{if } n \text{ is even.} \end{cases}$$

Proof Let $V(W_n) = \{c, v_i, 1 \leq i \leq n\}$ and $E(W_n) = \{cv_i, 1 \leq i \leq n\} \cup \{v_i v_{i+1}, 1 \leq i \leq n - 1\} \cup \{v_n v_1\}$. Then $|V(W_n)| = n + 1$ and $|E(W_n)| = 2n$. Now, define a bijection $f_5 : V(W_n) \rightarrow \{1, 2, \dots, n + 1\}$ by

$$f_5(c) = \begin{cases} \frac{n+1}{2} & \text{if } n \text{ is odd} \\ \frac{n+2}{2} & \text{if } n \text{ is even} \end{cases}$$

$$f_5(v_i) = \begin{cases} \frac{i+1}{2}, & \text{if } n \text{ is odd, } i \text{ is odd, } 1 \leq i \leq n - 2 \\ n + 1 & \text{if } n \text{ is odd, } i=2 \\ \frac{2n+2-i}{2}, & \text{if } n \text{ is odd, } i \text{ is even, } 4 \leq i \leq n - 1 \\ n, & \text{if } n \text{ is odd, } i=n \\ \frac{i+1}{2}, & \text{if } n \text{ is even, } i \text{ is odd, } 1 \leq i \leq n - 1 \\ n + 1 & \text{if } n \text{ is even, } i=2 \\ \frac{2n+2-i}{2}, & \text{if } n \text{ is even, } i \text{ is even, } 4 \leq i \leq n - 2 \\ n, & \text{if } n \text{ is even, } i=n. \end{cases}$$

The edge-weights of W_n are

$$w_5(cv_i) = \begin{cases} \frac{n+2+i}{2}, & \text{if } n \text{ is odd, } i \text{ is odd, } 1 \leq i \leq n - 2 \\ \frac{3n+3}{2} & \text{if } n \text{ is odd, } i=2 \\ \frac{3n+3-i}{2}, & \text{if } n \text{ is odd, } i \text{ is even, } 4 \leq i \leq n - 1 \\ \frac{3n+1}{2}, & \text{if } n \text{ is odd, } i=n \\ \frac{n+3+i}{2}, & \text{if } n \text{ is even, } i \text{ is odd, } 1 \leq i \leq n - 1 \\ \frac{3n+4}{2} & \text{if } n \text{ is even, } i=2 \\ \frac{3n+4-i}{2}, & \text{if } n \text{ is even, } i \text{ is even, } 4 \leq i \leq n - 2 \\ \frac{3n+2}{2}, & \text{if } n \text{ is even, } i=n \end{cases}$$

$$w_5(v_i v_{i+1}) = \begin{cases} n + 2, & \text{if } n \text{ is odd, } i=1, i \text{ is even, } 4 \leq i \leq n - 3 \\ n + 3, & \text{if } n \text{ is odd, } i=2 \\ n + 1, & \text{if } n \text{ is odd, } i \text{ is odd, } 3 \leq i \leq n - 2 \\ n + 2, & \text{if } n \text{ is even, } i=1, i \text{ is even, } 4 \leq i \leq n - 2 \\ n + 3, & \text{if } n \text{ is even, } i=2 \\ n + 1, & \text{if } n \text{ is even, } i \text{ is odd, } 3 \leq i \leq n - 3 \end{cases}$$

$$w_5(v_{n-1} v_n) = \begin{cases} \frac{3n+3}{2}, & \text{if } n \text{ is odd} \\ \frac{3n}{2}, & \text{if } n \text{ is even} \end{cases}$$

$$w_5(v_n v_1) = n + 1.$$

It is easy to identify that f_5 proves a proper edge coloring of W_n and hence $\chi'_{(a,1)\text{-ela}}(W_n) \leq n + 1$, where

$$a = \begin{cases} \frac{n+3}{2}, & n \text{ is odd} \\ \frac{n}{2} + 2, & n \text{ is even.} \end{cases}$$

□

Example 2.17 The graphs W_5 and W_6 admit $(a, 1)$ -ELA labeling with 6-colors and 7-colors, where $a = 4$ and 5, which is given in Figures 5 and 6.




Figure 5. The $(4, 1)$ -ELA labeling of W_5 with 6-colors.




Figure 6. The $(5, 1)$ -ELA labeling of W_6 with 7-colors.

3. Conclusion

In this paper, we have introduced the new type of (a, d) -ELA labeling and a parameter (a, d) -ELA coloring number of G . We obtained the (a, d) -ELA coloring number for paths when $d = 1$ and 2, and wheel graph $W_n, n = 3, 4, 5$ when $d = 1$. The problem of determining the (a, d) -ELA coloring number for the remaining graphs is still open.

Acknowledgment

The authors are thankful to the anonymous referees for helpful suggestions which led to substantial improvement in the presentation of the paper. S. Rajkumar is thankful to Vellore Institute of Technology, Vellore, India for providing a Teaching Research Associate Fellowship.

References

[1] Arumugam S, Premalatha K, Martin Bacá, Semaničová-Fečňovčíková A. Local Antimagic Vertex Coloring of a Graph. Graphs and Combinatorics 2017; 33: 275-285.

- [2] Chartrand G, Lesniak L. Graphs and Digraphs. Chapman and Hall, CRC, 4th edition, 2005.
- [3] Cheng Y. A new class of antimagic Cartesian product graphs. Discrete Mathematics 2008; 308: 6441-6448.
- [4] Gallian JA. A Dynamic Survey of Graph labeling. The Electronic Journal of Combinatorics #DS6 2020.
- [5] Hartsfield N, Ringel G. Pearls in graph theory. Academic Press, INC., Boston,1994.
- [6] Haslegrave J. Proof of a local antimagic conjecture. Discrete Mathematics and Theoretical Computer Science 2018; 20 (1): Article 18.
- [7] Ika Hesti A, Moh Hasan D, Ridho A, Prihandini RM. Local Edge Antimagic Coloring of Graphs. Far East Journal of Mathematical Sciences 2017; 102 (9): 1925-1941.
- [8] Lau GC Every graph is local antimagic total and its application to local antimagic (total) chromatic number. arXiv:1906.10332v23[math.co] 2020.
- [9] Martin Bača, Mirka M. Super Edge Antimagic Graphs A Wealth of Problems and Some Solutions, Brown Walker Press, Baco Raton, Florida, USA 2008.
- [10] Rajkumar S, Nalliah. M On Local edge antimagic chromatic number of graphs. Proyecciones 2022; Accepted.
- [11] Shankar R, Nalliah M. Local Vertex Antimagic Chromatic Number of Some Wheel Related Graphs. Proyecciones 2022; 41 (1): 319-334.
- [12] Wang TM, Zhang GH. Arumugam S, Smyth B. (Eds.) On antimagic labeling of odd regular graphs. Lecture notes in Computer Science(International Workshop on Combinatorial Algorithms) 2012; 7643: 162-168.