

1-1-2013

## Mean platelet volume as a fibrosis marker in patients with chronic hepatitis C

HATİCE KARAMAN

ÇİĞDEM KARAKÜKCÜ

AHMET KARAMAN

TUBA KAYMAN

SAMET YALÇIN

*See next page for additional authors*

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

---

### Recommended Citation

KARAMAN, HATİCE; KARAKÜKCÜ, ÇİĞDEM; KARAMAN, AHMET; KAYMAN, TUBA; YALÇIN, SAMET; TAŞDEMİR, ERDEM ARZU; ZARARSIZ, GÖKMEN; and POYRAZOĞLU, ORHAN KÜRŞAT (2013) "Mean platelet volume as a fibrosis marker in patients with chronic hepatitis C," *Turkish Journal of Medical Sciences*: Vol. 43: No. 1, Article 8. <https://doi.org/10.3906/sag-1204-17>

Available at: <https://journals.tubitak.gov.tr/medical/vol43/iss1/8>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

---

## Mean platelet volume as a fibrosis marker in patients with chronic hepatitis C

### Authors

HATİCE KARAMAN, ÇİĞDEM KARAKÜKCÜ, AHMET KARAMAN, TUBA KAYMAN, SAMET YALÇIN, ERDEM ARZU TAŞDEMİR, GÖKMEN ZARARSIZ, and ORHAN KÜRŞAT POYRAZOĞLU

## Mean platelet volume as a fibrosis marker in patients with chronic hepatitis C

Hatice KARAMAN<sup>1\*</sup>, Çiğdem KARAKÜKCÜ<sup>2</sup>, Ahmet KARAMAN<sup>3</sup>, Tuba KAYMAN<sup>4</sup>,  
Samet YALÇIN<sup>5</sup>, Erdem Arzu TAŞDEMİR<sup>1</sup>, Gökmen ZARARSIZ<sup>6</sup>, Orhan Kürşat POYRAZOĞLU<sup>3</sup>

<sup>1</sup>Department of Pathology, Education and Research Hospital, Kayseri, Turkey

<sup>2</sup>Department of Clinical Biochemistry, Education and Research Hospital, Kayseri, Turkey

<sup>3</sup>Department of Gastroenterology, Education and Research Hospital, Kayseri, Turkey

<sup>4</sup>Department of Clinical Microbiology, Education and Research Hospital, Kayseri, Turkey

<sup>5</sup>Department of General Surgery, Faculty of Medicine, Yildirim Beyazıt University, Ankara, Turkey

<sup>6</sup>Department of Biostatistics and Medical Informatics, Faculty of Medicine, Erciyes University, Kayseri, Turkey

Received: 03.04.2012

Accepted: 03.07.2012

Published Online: 18.01.2013

Printed: 18.02.2013

**Aim:** We designed this study to evaluate the role of mean platelet volume (MPV) as a fibrosis marker in patients with chronic hepatitis C (CHC).

**Materials and methods:** The study was designed at Kayseri Education and Research Hospital. Ninety-five patients with CHC were enrolled retrospectively into the study. The control group comprised 33 age- and sex-matched healthy individuals. Hepatitis C virus-RNA level, alanine aminotransferase (ALT), aspartate aminotransferase (AST), bilirubin, prothrombin time (PT), platelet count (PC), serum albumin, histological activity index (HAI), liver fibrosis score (LFS), and MPV were obtained from the patients' files and a computerized database.

**Results:** Statistically significant differences in MPV and PC were seen in patients with CHC compared to healthy controls (MPV:  $9.1 \pm 1.31$  fL vs.  $8.58 \pm 0.8$  fL,  $P = 0.008$ ; PC ( $\times 10^3/\mu\text{L}$ ):  $219.37 \pm 74.31$  vs.  $258.52 \pm 48.34$ ,  $P = 0.001$ ). In multivariate analysis, 4 variables remained as independent risk factors: AST (OR 1.11, 95% CI 1.02 to 1.21), ALT (OR 0.92, 95% CI 0.86 to 0.99), PT (OR 2.11, 95% CI 1.15 to 3.88), and MPV (OR 2.28, 95% CI 1.22 to 4.25). Cut-off values were calculated for diagnostic performance, and the cut-off value for MPV was 8.4 fL.

**Conclusion:** We suggest that high MPV levels (especially those over 8.4 fL) may help to predict advanced fibrosis in patients with CHC. However, it should not be forgotten that MPV is not a specific marker for fibrosis, and the negative predictive rate seems more valuable to exclude a high fibrosis ratio in patients with CHC.

**Key words:** MPV, chronic hepatitis C, fibrosis

### 1. Introduction

Hepatitis C virus (HCV) was discovered in 1989 (1). It was found to be responsible for the majority of chronic non-A and non-B hepatitis and cryptogenetic liver diseases (2,3). In a study from Turkey, it was found that the prevalence of HCV was 0.4% in blood donors (4). In adults, acute HCV infection leads to chronic infection in approximately 80% of cases. Approximately 120–130 million individuals are chronically infected with HCV worldwide. The risk of chronic hepatitis C (CHC) infection after an acute episode is high. In many studies, 80% to 100% of patients remain HCV-RNA positive, and 60% to 80% have persistently elevated liver enzymes (5). The mechanism responsible

for the high prevalence of CHC infection is unclear. It may be related to the genetic diversity of the virus and its tendency toward rapid mutation, allowing HCV to escape immune recognition (6). Host factors may also be involved in the ability of spontaneous clearance of the virus. One of the most influential factors appears to be certain polymorphisms of a site close to the interleukin-28B (IL28B) gene, and polymorphisms are also an important predictor of response to treatment (7). Chronic HCV infection is responsible for chronic hepatitis, which results in cirrhosis in approximately 20% of cases. Patients with cirrhosis are exposed to life-threatening complications, including end-stage liver disease, peritonitis, esophageal

\* Correspondence: htckaraman@yahoo.com

variceal hemorrhage, and the development of hepatocellular carcinoma, which occurs at an incidence of 4%–5% per year (8). It is essential to stop the progression of early fibrosis to cirrhosis. Liver biopsy is the gold standard for the assessment of liver histology and stage of fibrosis, but it has some problems, such as bleeding, poor patient compliance, sampling error, and coagulation problems. Because of these problems, there is always an effort to find a method or a laboratory test for predicting the stage of fibrosis stage in CHC patients.

Platelet volume is an indicator of platelet function and activation (9). Platelet activity and aggregation capacity can be easily determined by measuring mean platelet volume (MPV) (10). Large platelets have more granules, aggregate more rapidly with collagen, produce higher levels of thromboxane A<sub>2</sub>, and express more glycoprotein Ib and IIb/IIIa receptors than smaller ones (11). MPV has been reported as a risk factor for atherothrombosis. Elevated MPV values have been shown in atherothrombotic diseases like acute myocardial ischemia, acute myocardial infarction, coronary atherosclerosis, and cerebrovascular events (12–14). Additionally, MPV has been shown to be a sign of inflammation in ulcerative colitis, Crohn's disease, and rheumatoid arthritis (15,16). In a study from Turkey, MPV was reported to be independently associated with advanced fibrosis in patients with chronic hepatitis B (17). In the present study, we aimed to investigate the association of MPV and stage of liver fibrosis in patients with CHC.

## 2. Materials and methods

Ninety-five patients with biopsy-proven CHC were enrolled in the study. The control group comprised 33 age- and sex-matched healthy subjects. The following data were obtained from a computerized patient registry database: HCV-RNA level, alanine aminotransferase (ALT), aspartate aminotransferase (AST), bilirubin, prothrombin time (PT), platelet count (PC), serum albumin (ALB), histological activity index (HAI), liver fibrosis score (LFS), and MPV. Laboratory findings of patients were screened before treatment. Exclusion criteria were atherosclerotic heart disease, any medication use that can influence platelet function (e.g., aspirin), diabetes mellitus, asthma, chronic obstructive pulmonary disease, peripheral and cerebral vascular disease, hematological disorders, and malignancies.

### 2.1. Histopathological evaluation

Fibrosis staging was assessed according to the METAVIR system. LFSs were defined as follows: score for fibrosis (F): F0, no fibrosis; F1, portal fibrosis without septa; F2, portal fibrosis with rare septa; F3, numerous septa without cirrhosis; and F4, cirrhosis (18). According to the METAVIR scoring system, patients were divided into 2 groups: patients without significant fibrosis (F0, F1, or F2)

(Group 1) and patients with advanced fibrosis (F3 or F4) (Group 2).

### 2.2. Statistical analysis

The Shapiro–Wilk test was used to assess the normality of the data. Accordingly, either an independent samples t-test or Mann–Whitney U tests and one-way analysis of variance were used to compare the differences of continuous variables between groups. For RNA values, a logarithmic transformation was applied because of high skewness. Chi-square analysis was used to compare the differences of categorical variables. Results were expressed as frequencies and percentages, mean  $\pm$  standard deviation, or median (25th and 75th percentiles). Univariate and multivariate logistic regression analyses were also performed and odds ratios with 95% confidence intervals (CIs) were calculated in order to identify risk factors of liver fibrosis in patients with CHC. Statistically significant variables in univariate analysis were taken to multivariate analysis, and backward stepwise elimination was used at a  $P < 0.05$  stringency level to identify the independent risk factors of liver fibrosis. Receiver operating characteristic (ROC) curves were drawn for these risk factors, and the areas under the ROC curve (AUC) values with 95% CIs were calculated and compared with each other. A cut-off value was determined for each factor. Sensitivity, specificity, positive predictive rate (PPR), negative predictive rate (NPR), and accuracy rate (AR) diagnostic measures were calculated, and Kappa tests were performed for each factor with the given cut-off value. Matthew's correlation coefficient (also referred to as the phi coefficient and similar to the Pearson correlation coefficient in its interpretation) was also calculated because the fibrosis classes were of different sizes (19). A probability level of  $P < 0.05$  was considered statistically significant. MedCalc (Version 9.2.0.1) and SPSS 15.0 (SPSS Inc., Chicago, IL, USA) software were used for all analyses.

## 3. Results

Demographic data and laboratory findings of patient and control groups are reported in Table 1. Age and sex were similar among the groups. Statistically significant differences in MPV and PC were seen in patients with CHC compared to healthy controls (MPV:  $9.1 \pm 1.31$  fL vs.  $8.58 \pm 0.8$  fL,  $P = 0.008$ ; PC ( $\times 10^3/\mu\text{L}$ ):  $219.37 \pm 74.31$  vs.  $258.52 \pm 48.34$ ,  $P < 0.001$ ).

Serum AST, ALT, PT, bilirubin, ALB, PC, and MPV were significantly different between groups 1 and 2. Univariate logistic regression analysis revealed an association between fibrosis in CHC patients and serum AST, ALT, ALB, PT, PLT, and MPV values (Table 2). In multivariate analysis, 4 variables remained as independent risk factors: AST (OR 1.11, 95% CI 1.02 to 1.21), ALT (OR 0.92, 95% CI 0.86 to 0.99), PT (OR 2.11, 95% CI 1.15 to 3.88), and MPV (OR 2.28, 95% CI 1.22 to 4.25) (Table 2).

**Table 1.** Demographic parameters of control and patient groups.

	Control (n = 33)	CHC (n = 95)	P
Age (years)	55.12 ± 8.12	54.28 ± 10.72	0.56
Sex (female/male)	18 (54.5%)/15 (45.5%)	63 (66.3%)/52 (33.7%)	0.318
Platelet count (×10 <sup>3</sup> µL)	258.52 ± 48.34	219.37 ± 74.31	0.001
MPV (fL)	8.58 ± 0.80	9.10 ± 1.31	0.008

Values are expressed as n (%) or mean ± standard deviation.

**Table 2.** Demographic characteristics and laboratory findings of CHC patients.

	Between-group comparisons			Logistic regression analysis	
	Group 1 (n = 72) F0, F1, F2	Group 2 (n = 23) F3, F4	P	Univariate OR (95% CI)	Multivariate OR (95% CI)
Age (years)	53.89 ± 10.48	55.52 ± 11.62	0.528	1.02 (0.97–1.07)	-
Sex (female*/male)	50 (69.4%)/22 (30.6%)	13 (56.5%)/10 (43.5%)	0.374	1.75 (0.67–4.59)	-
AST (IU/L)	33.00 (25.00–48.00)	51.00 (34.50–81.75)	0.002	1.03 (1.01–1.05)	1.11 (1.02–1.21)
ALT (IU/L)	42.00 (28.25–50.75)	63.00 (29.50–96.50)	0.050	1.02 (1.00–1.03)	0.92 (0.86–0.99)
PT (s)	13.27 ± 1.33	14.37 ± 1.18	0.001	1.97 (1.28–3.04)	2.11 (1.15–3.88)
BIL (mg/dL)	0.70 (0.45–0.90)	0.90 (0.60–1.18)	0.037	2.99 (0.88–10.13)	-
ALB (mg/dL)	4.10 (4.00–4.33)	3.90 (3.45–4.20)	0.018	0.14 (0.04–0.56)	-
log (RNA)	5.78 ± 0.72	5.42 ± 0.96	0.083	0.58 (0.31–1.09)	-
PC (×10 <sup>3</sup> µL)	235.14 ± 66.13	170.00 ± 78.24	<0.001	0.98 (0.97–0.99)	-
MPV (fL)	8.91 ± 1.24	9.67 ± 1.39	0.015	1.56 (1.07–2.27)	2.28 (1.22–4.25)

AST, aspartate aminotransferase; ALT, alanine aminotransferase; PT, prothrombin time; BIL, bilirubin; ALB, albumin; RNA, ribonucleic acid; PC, platelet count; MPV, mean platelet volume. Values are expressed as n (%), mean ± SD or median (25th and 75th percentiles).


\*Reference category for logistic regression analysis.

Comparison of ROC curves for the diagnostic performance of AST, ALT, PT, and MPV in identifying fibrosis in CHC is shown in Figure 1a. AUC values for AST, ALT, PT, and MPV were 0.74 (0.63–0.82), 0.66 (0.55–0.76), 0.73 (0.63–0.82), and 0.66 (0.55–0.76), respectively, and the differences of these values between parameters were not statistically significant ( $P > 0.05$ ). A plot to obtain a cut-off value for MPV that displays sensitivity and specificity variation for each MPV value and a dot diagram that plots the distribution of CHC samples with mild and severe fibrosis around an 8.4-fL cut-off value are shown in Figures 1b and 1c. In addition, sensitivity, specificity, PPR, NPR, and AR were demonstrated for the independent risk factors AST, ALT, PT, and MPV (Table 3).

#### 4. Discussion

CHC infection often follows a progressive course over years and may result in cirrhosis, hepatocellular carcinoma, and the need for liver transplantation. Because of the long course of the disease, the natural history of CHC has not been clearly defined. Many studies have provided estimates of the proportion of patients with CHC infection who develop cirrhosis within 20 years (20,21).

The MPV is the geometric mean of the transformed lognormal platelet volume data in impedance technology systems. In some optical systems, MPV is the mode of the measured platelet volume (22). There is an inverse relationship between platelet size and number. Therefore, the total platelet mass, the product of the MPV and



**Figure 1.** a) Comparison of ROC curves for the diagnostic performance of AST, ALT, PT, and MPV in identifying fibrosis in CHC. AUC values were 0.74 (0.63–0.82), 0.66 (0.55–0.76), 0.73 (0.63–0.82), and 0.66 (0.55–0.76), respectively, and the differences of these values between parameters were not statistically significant ( $P > 0.05$ ). b) A plot to obtain a cut-off value for MPV that displays sensitivity and specificity variation for each MPV value. c) A dot diagram that plots the distribution of CHC samples with mild and severe fibrosis around an 8.4-fL cut-off value.

**Table 3.** Diagnostic measures and Kappa test results of parameters in the detection of liver fibrosis.

Parameters	Diagnostic measures						Kappa test	
	SEN (95% CI)	SPE (95% CI)	PPR (95% CI)	NPR (95% CI)	AR (95% CI)	MCC (95% CI)	$\kappa$	P
AST (42.0 IU/L)	72.7 (49.8–89.3)	66.7 (54.6–77.3)	40.0 (24.9–56.7)	88.9 (77.4–95.8)	68.1 (57.7–77.3)	33.7 (14.8–52.7)	0.307	0.001
ALT (51.0 IU/L)	63.6 (40.7–82.8)	75.0 (63.4–84.5)	43.8 (26.4–62.3)	87.1 (76.2–94.3)	72.3 (62.2–81.1)	34.5 (14.2–54.8)	0.334	<0.001
PT (14.7 s)	50.0 (28.2–71.8)	85.1 (74.3–92.6)	52.4 (29.8–74.3)	83.8 (72.9–91.6)	76.4 (66.2–84.8)	35.6 (13.4–57.9)	0.356	<0.001
MPV (8.4 fL)	91.3 (72.0–98.9)	43.7 (31.9–56.0)	34.4 (22.7–47.7)	93.9 (79.8–99.3)	55.3 (44.7–65.6)	31.5 (16.9–46.1)	0.224	0.002

SEN, sensitivity; SPE, specificity; PPR, positive predictive rate; NPR, negative predictive rate; AR, accuracy rate; MCC, Matthew's correlation coefficient.

platelet count, is closely regulated. When platelets decrease in number, bone marrow megakaryocytes are stimulated by thrombopoietin and their nucleus becomes hyperlobulated, with much higher DNA content. These stimulated megakaryocytes produce larger platelets. Thus, platelets with a higher MPV are expected to be seen in destructive thrombocytopenia when megakaryocytic stimulation is present. Conversely, platelets with a lower MPV are expected in thrombocytopenic states associated with marrow hypoplasia or aplasia. An exception to this relationship occurs with splenic sequestration, in which a low MPV is seen because the spleen sequesters large platelets. In hyposplenic states a higher MPV is seen, since there is no spleen to sequester the larger platelets (23–25).

There are many studies on the relationship between MPV and some thrombotic and cardiac disorders, but there are few on inflammatory diseases. Beyazit et al. (26) reported that overall accuracy of MPV in predicting disease severity according to computerized tomography severity index was not superior compared with other inflammation markers in patients with acute pancreatitis. Zubcevic et al. (22) reported that one of the most reliable indicators of activity of Crohn's disease was MPV; however, it was not sensitive enough to distinguish the relationship between moderate and severe disease. A decrease in MPV reflects the activation and participation of platelets in the inflammatory process of colon mucosa, and so MPV may be a useful marker of active ulcerative colitis (27). Korkmaz et al. (28) reported that higher MPV may reflect increased atherosclerotic burden and cardiovascular risk. Yazici et al. (29) reported that MPV may prove to be useful as a prognostic marker in patients with metabolic syndrome and ST elevation in myocardial infarction. Köşüş et al. (30) reported that mild glycemic disorders are associated with increased MPV. Increased MPV might be associated with elevated baseline cardiovascular risk factors. Individuals with these glycemic disorders might be more aggressively targeted with strategies to lower cardiovascular disease risk.

In this study, our aim was to investigate whether MPV could be a marker for fibrosis in CHC patients. We found that MPV was significantly higher in patients with CHC when compared to control subjects. In contrast, PC was significantly lower in CHC patients. Portal hypertension and hypersplenism in some of the subjects with advanced fibrosis may be the cause of this significant difference. Patients were divided into 2 groups according to severity of fibrosis. AST, ALT, ALB, PT, PC, and MPV were significantly different between the groups. Therefore, we suggest that these parameters are affected by the degree of fibrosis in CHC. The independent risk factors for fibrosis were AST, ALT, PT, and MPV in this study. The cut-off

value for MPV to predict advanced fibrosis in patients with CHC was 8.4 fL. At this cut-off value, sensitivity was 91.3% and specificity was 43.7%. The predictive role of some laboratory tests in the detection of liver fibrosis was previously investigated in patients with CHC. Kandemir et al. (31) reported that PC and the GUCI (Goteborg University Cirrhosis Index, calculated using AST, PC, and PT) can discriminate to some degree of accuracy patients with severe fibrosis. We also found AST, ALT, PT, and MPV to have diagnostic values in identifying severity of fibrosis in this study.

The question of this study is why MPV is related to advanced fibrosis. It is well known that a high MPV in a thrombocytopenic patient indicates active marrow production of platelets, whereas a low MPV is indicative of bone marrow suppression (32). The PC of patients with advanced fibrosis was significantly lower in our study. This finding could be one reason for the higher MPV ratio in patients with advanced fibrosis.

Hepatic steatosis and insulin resistance as a part of metabolic syndrome in CHC patients are associated with hepatic fibrosis (33). A high MPV ratio has been observed in some diseases associated with metabolic syndrome (28,34,35). The relationship between CHC infection and metabolic syndrome and its components may be another reason for the higher MPV ratio in patients with CHC.

The higher MPV and lower PCs in patients with advanced fibrosis can lead to thrombocytopenia and platelet-endothelial adhesion dysfunction, which affect clot formation and contribute to a relative hypocoagulable situation in patients with advanced fibrosis. The surface phospholipids of platelets provide the platform for factor complexes, amplification, and propagation of clot formation. Many theories exist regarding the genesis of thrombocytopenia in patients with liver disease. Decreased thrombopoietin levels (36), splenic sequestration of platelets due to portal hypertension, auto-antibody destruction of platelets (37), and bone marrow suppression due to underlying liver disease can all contribute to thrombocytopenia. Decreased platelet count may mobilize young platelets (with high MPV) in bone marrow in patients with CHC. Here we want to point out the quite high NPR at this cut-off value (93.9%), which reveals the absence of severe fibrosis in CHC under this value.

In conclusion, we suggest that high MPV levels (especially those over 8.4 fL) may help to predict advanced fibrosis in patients with CHC. However, it should not be forgotten that MPV is not a specific marker for fibrosis, and a high NPR seems to be more important in helping to exclude a high fibrosis ratio in patients with CHC.



## References

1. Choo QL, Kuo G, Weiner AJ, Overby LR, Bradley DW, Houghton M. Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science* 1989; 244: 359–62.
2. Alter HJ, Purcell RH, Shih JW, Melpolder JC, Houghton M, Choo QL et al. Detection of antibody to hepatitis C virus in prospectively followed transfusion recipients with acute and chronic non-A, non-B hepatitis. *N Engl J Med* 1989; 321: 1494–500.
3. van der Poel CL, Reesink HW, Lelie PN, Leentvaar-Kuypers A, Choo QL, Kuo G et al. Anti-hepatitis C antibodies and non-A, non-B post-transfusion hepatitis in the Netherlands. *Lancet* 1989; 2: 297–8.
4. Öner S, Yapıcı G, Şaşmaz CT, Kurt AÖ, Buğdaycı R. Hepatitis B, hepatitis C, HIV, and VDRL seroprevalence of blood donors in Mersin, Turkey. *Turk J Med Sci* 2011; 41: 335–41.
5. Barrera JM, Bruguera M, Ercilla MG, Gil C, Celis R, Gil MP et al. Persistent hepatitis C viremia after acute self-limiting posttransfusion hepatitis C. *Hepatology* 1995; 21: 639–44.
6. Farci P, Shimoda A, Coiana A, Diaz G, Peddis G, Melpolder JC et al. The outcome of acute hepatitis C predicted by the evolution of the viral quasispecies. *Science* 2000; 288: 339–44.
7. Grebely J, Petoumenos K, Hellard M, Matthews GV, Suppiah V, Applegate T et al. Potential role for interleukin-28B genotype in treatment decision-making in recent hepatitis C virus infection. *Hepatology* 2010; 52: 1216–24.
8. Seeff LB. Natural history of chronic hepatitis C. *Hepatology* 2002; 36: S35–46.
9. Park Y, Schoene N, Harris W. Mean platelet volume as an indicator of platelet activation: methodological issues. *Platelets* 2002; 13: 301–6.
10. Bancroft AJ, Abel EW, McLaren M, Belch JJ. Mean platelet volume is a useful parameter: a reproducible routine method using a modified Coulter thrombocytometer. *Platelets* 2000; 11: 379–87.
11. Giles H, Smith RE, Martin JF. Platelet glycoprotein IIb-IIIa and size are increased in acute myocardial infarction. *Eur J Clin Invest* 1994; 24: 69–72.
12. Kiliçli-Camur N, Demirtunç R, Konuralp C, Eskiser A, Başaran Y. Could mean platelet volume be a predictive marker for acute myocardial infarction? *Med Sci Monit* 2005; 11: 387–92.
13. Pizzulli L, Yang A, Martin JF, Lüderitz B. Changes in platelet size and count in unstable angina compared to stable angina or non-cardiac chest pain. *Eur Heart J* 1998; 19: 80–4.
14. Bath P, Algert C, Chapman N, Neal B; PROGRESS Collaborative Group. Association of mean platelet volume with risk of stroke among 3134 individuals with history of cerebrovascular disease. *Stroke* 2004; 35: 622–6.
15. Yüksel O, Helvacı K, Başar O, Köklü S, Caner S, Helvacı N et al. An overlooked indicator of disease activity in ulcerative colitis: mean platelet volume. *Platelets* 2009; 20: 277–81.
16. Kisacık B, Tufan A, Kalyoncu U, Karadağ O, Akdoğan A, Öztürk MA et al. Mean platelet volume (MPV) as an inflammatory marker in ankylosing spondylitis and rheumatoid arthritis. *Joint Bone Spine* 2008; 75: 291–4.
17. Ekiz F, Yüksel O, Koçak E, Yılmaz B, Altınbaş A, Çoban S et al. Mean platelet volume as a fibrosis marker in patients with chronic hepatitis B. *J Clin Lab Anal* 2011; 25: 162–5.
18. Intraobserver and interobserver variations in liver biopsy interpretation in patients with chronic hepatitis C. The French METAVIR Cooperative Study Group. *Hepatology* 1994; 20: 15–20.
19. Baldi P, Brunak S, Chauvin Y, Andersen CA, Nielsen H. Assessing the accuracy of prediction algorithms for classification: an overview. *Bioinformatics* 2000; 16: 412–24.
20. Liang TJ, Rehermann B, Seeff LB, Hoofnagle JH. Pathogenesis, natural history, treatment, and prevention of hepatitis C. *Ann Intern Med* 2000; 132: 296–305.
21. Rodger AJ, Roberts S, Lanigan A, Bowden S, Brown T, Crofts N. Assessment of long-term outcomes of community-acquired hepatitis C infection in a cohort with sera stored from 1971 to 1975. *Hepatology* 2000; 32: 582–7.
22. Zubcevic N, Mesihovic R, Zubcevic S. Usefulness of laboratory data in estimation of Crohn's disease activity. *Med Arh* 2010; 64: 33–6.
23. Jackson SR, Carter JM. Platelet volume: laboratory measurement and clinical application. *Blood Rev* 1993; 7: 104–13.
24. Threatte GA. Usefulness of the mean platelet volume. *Clin Lab Med* 1993; 13: 937–50.
25. Corash L. Platelet sizing: techniques, biological significance, and clinical applications. *Curr Top Hematol* 1983; 4: 99–122.
26. Beyazıt Y, Sayılır A, Torun S, Suvak B, Yeşil Y, Purnak T et al. Mean platelet volume as an indicator of disease severity in patients with acute pancreatitis. *Clin Res Hepatol Gastroenterol* 2012; 36: 162–8.
27. Polińska B, Matowicka-Karna J, Kemonia H. Assessment of the influence of the inflammatory process on the activation of blood platelets and morphological parameters in patients with ulcerative colitis (colitis ulcerosa). *Folia Histochem Cytobiol* 2011; 49: 119–24.
28. Korkmaz L, Korkmaz AA, Akyüz AR, Ağaç MT, Acar Z, Kırış A et al. Association between mean platelet volume and coronary artery calcification in patients without overt cardiovascular disease: an observational study. *Anadolu Kardiyol Derg* 2012; 12: 35–9.
29. Yazıcı HU, Poyraz F, Sen N, Tavil Y, Turfan M, Tulmaç M et al. Relationship between mean platelet volume and left ventricular systolic function in patients with metabolic syndrome and ST-elevation myocardial infarction. *Clin Invest Med* 2011; 34: E330.



30. Köşüş N, Köşüş A, Turhan N. Mean platelet volume as a marker of future cardiovascular disease risk in pregnant women with impaired fasting glucose and impaired glucose tolerance. *Turk J Med Sci* 2012; 42: 245–51.
31. Kandemir O, Polat G, Saraçoğlu G, Taşdelen B. The predictive role of AST level, prothrombin time and platelet count in the detection of liver fibrosis in patients with chronic hepatitis C. *Turk J Med Sci* 2009; 39: 857–62.
32. Kaito K, Otsubo H, Usui N, Yoshida M, Tanno J, Kurihara E et al. Platelet size deviation width, platelet large cell ratio, and mean platelet volume have sufficient sensitivity and specificity in the diagnosis of immune thrombocytopenia. *Br J Haematol* 2005; 128: 698–702.
33. Hui JM, Sud A, Farrell GC, Bandara P, Byth K, Kench JG et al. Insulin resistance is associated with chronic hepatitis C virus infection and fibrosis progression [corrected]. *Gastroenterology* 2003; 125: 1695–704.
34. Purnak T, Efe C, Beyazit Y, Ozaslan E, Astan R, Milanoglu A et al. Recent insights into the relationship between inflammatory liver diseases and atherosclerosis. *J Investig Med* 2011; 59: 904–11.
35. Wree A, Kahraman A, Gerken G, Canbay A. Obesity affects the liver - the link between adipocytes and hepatocytes. *Digestion* 2011; 83: 124–33.
36. Goulis J, Chau TN, Jordan S, Mehta AB, Watkinson A, Rolles K et al. Thrombopoietin concentrations are low in patients with cirrhosis and thrombocytopenia and are restored after orthotopic liver transplantation. *Gut* 1999; 44: 754–8.
37. Kajihara M, Okazaki Y, Kato S, Ishii H, Kawakami Y, Ikeda Y et al. Evaluation of platelet kinetics in patients with liver cirrhosis: similarity to idiopathic thrombocytopenic purpura. *J Gastroenterol Hepatol* 2007; 22: 112–8.