

1-1-2022

## Solving a class of ordinary differential equations and fractional differential equations with conformable derivative by fractional Laplace transform

MOHAMMAD MOLAEI

FARHAD DASTMALCHI SAEI

MOHAMMAD JAVIDI

YAGHOUB MAHMOUDI

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

MOLAEI, MOHAMMAD; SAEI, FARHAD DASTMALCHI; JAVIDI, MOHAMMAD; and MAHMOUDI, YAGHOUB (2022) "Solving a class of ordinary differential equations and fractional differential equations with conformable derivative by fractional Laplace transform," *Turkish Journal of Mathematics*: Vol. 46: No. 7, Article 30. <https://doi.org/10.55730/1300-0098.3316>

Available at: <https://journals.tubitak.gov.tr/math/vol46/iss7/30>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Solving a class of ordinary differential equations and fractional differential equations with conformable derivative by fractional Laplace transform

Mohammad MOLAEI<sup>1</sup> , Farhad DASTMALCHI SAEI<sup>1,\*</sup> ,  
Mohammad JAVIDI<sup>1,2</sup> , Yaghoub MAHMOUDI<sup>1</sup> 

<sup>1</sup>Department of Mathematics, Tabriz Branch, Islamic Azad University, Tabriz, Iran

<sup>2</sup>Faculty of Mathematical Sciences, University of Tabriz, Tabriz, Iran

Received: 12.04.2021

Accepted/Published Online: 15.08.2022

Final Version: 05.09.2022

**Abstract:** In this paper, we use the fractional Laplace transform to solve a class of second-order ordinary differential equations (ODEs), as well as some conformable fractional differential equations (CFDEs), including the Laguerre conformable fractional differential equation. Specifically, we apply the transform to convert the differential equations into first-order, linear differential equations. This is done by using the fractional Laplace transform of order  $\alpha + \beta$  or  $\alpha + \beta + \gamma$ . Also, we investigate some more results on the fractional Laplace transform, obtained by Abdeljawad.

**Key words:** Fractional Laplace transform, conformable fractional derivative, conformable fractional differential equation, Laguerre conformable fractional differential equation

### 1. Introduction

Fractional calculus has been of interest to many researchers of the last and present centuries. Many researchers have dealt with the discrete version of fractional calculus, which benefits from the theory of time scales. However, in recent decades, fractional calculus and fractional differential equations have gained great development in both theory and applications, because of their powerful, potential applications. Fractional differential equations are sometimes referred to as extraordinary differential equations, because of their nature and the fact that they can be easily found in various fields of applied sciences [2, 10]. For example, fractional-order differential equations have been applied to the modeling of real-world phenomena in such diverse fields as physics, engineering, mechanics, control theory, economics, medical sciences, finance, etc. See [1–8, 15, 17].

In recent years, scientists have proposed many efficient and powerful methods to obtain exact or numerical solutions of fractional differential equations [15, 18]. In addition, many researchers have tried to propose new definitions of fractional derivatives. Such a definition usually uses an integral form for the fractional derivative. There are many types of differential derivatives in fractional calculus, including the derivatives of Grunwald–Letnikov, Riemann–Liouville, Caputo [7], Caputo–Fabrizio [11] and Atangana–Baleanu [2]. Recently, some authors introduced the concept of nonlocal derivative.

In [15], Khalil et al. proposed a new notion of derivative, prominently compatible with the classical one. This operator is called a "conformable derivative". The chain rule, which is an applicable and useful rule in calculus, holds authentically only for conformable fractional derivatives. This derivative satisfies some

\*Correspondence: E-mail: dastmalchi@iaut.ac.ir

2010 AMS Mathematics Subject Classification: 34A08, 44A10

conventional properties and can be used to solve conformable differential equations. In [18], the author used the conformable derivative to find the exact solutions of time heat differential equations. The conformable fractional derivative has many advantages in its properties. The impact of this fractional derivative in both pure and applied branches of sciences and engineering has increased substantially during the last decade. As a result, it is now widely used in many research fields.

This paper is organized as follows. In Section 2, we prove some important theorems based on the conformable fractional derivative. In Section 3, we present some conformable fractional Laplace theorems and provide some examples which are solved by taking the fractional Laplace transformation of order  $\alpha$ . In Section 4, we develop some new methods for solving a class of conformable fractional differential equations using fractional Laplace transformations of order  $\alpha + \beta$ . In Section 5, we present some new methods for solving a class of second-order ODEs using fractional Laplace transformations. Finally, in Section 6, we solve the Laguerre conformable fractional differential equation and some other conformable fractional differential equations by taking the fractional Laplace transformation of order  $\alpha + \beta$ .

## 2. Basic definitions and tools related to the conformable fractional derivative

Despite becoming a popular topic in recent years, the concept of the fractional derivative emerged in the late 17th century. There are several definitions for fractional derivatives. Recently, the conformable fractional derivative was introduced by Khalil et al. [15], using a limit operator. After that, Abdeljawad [17] has also presented fractional versions of the chain rule, exponential functions, Gronwall's inequality, Taylor power series expansions, and fractional Laplace transform for the conformable derivative.

In [15], Khalil et al. introduced a new kind of fractional derivatives as follows.

**Definition 2.1** *The left conformable fractional derivative of order  $0 < \alpha \leq 1$  of a function  $u : [a, +\infty) \rightarrow \mathbb{R}$ , starting from  $a \in \mathbb{R}$ , is defined by*

$$({}_t T_\alpha^a u)(t) = \lim_{\epsilon \rightarrow 0} \frac{u(t + \epsilon(t-a)^{1-\alpha}) - u(t)}{\epsilon}. \quad (2.1)$$

When  $a = 0$ ,

$$({}_t T_\alpha^0 u)(t) = {}_t T_\alpha u(t) = \lim_{\epsilon \rightarrow 0} \frac{u(t + \epsilon t^{1-\alpha}) - u(t)}{\epsilon}.$$

If  $({}_t T_\alpha^a u)(t)$  exists on  $(a, +\infty)$ , then  $({}_t T_\alpha^a u)(a) = \lim_{t \rightarrow a^+} ({}_t T_\alpha^a u)(t)$ . If  $({}_t T_\alpha^a u)(z_0)$  exists and is finite, then we say that  $u$  is left  $\alpha$ -differentiable at  $z_0$ . See [17].

**Theorem 2.2** *Suppose that  $0 < \alpha \leq 1$ , and that  $u_1$  and  $u_2$  are left  $\alpha$ -differentiable functions. Then, the following statements are true.*

- 1)  $\forall \omega_1, \omega_2 \in \mathbb{R}$ ,  $({}_t T_\alpha^a(\omega_1 u_1 + \omega_2 u_2))(t) = \omega_1 ({}_t T_\alpha^a u_1)(t) + \omega_2 ({}_t T_\alpha^a u_2)(t)$ .
- 2)  $\forall p \in \mathbb{R}$ ,  ${}_t T_\alpha^a((t-a)^p) = p(t-a)^{p-\alpha}$ .
- 3) If  $C$  is a constant, then  ${}_t T_\alpha^a(C) = 0$ .
- 4) (Product rule for the left CF derivatives)  $({}_t T_\alpha^a(u_1 u_2))(t) = u_1(t) ({}_t T_\alpha^a u_2)(t) + u_2(t) ({}_t T_\alpha^a u_1)(t)$ .

5) (Quotient rule for the left CF derivatives)  $\left({}_t T_\alpha^a \left(\frac{u_1}{u_2}\right)\right)(t) = \frac{u_2(t)({}_t T_\alpha^a u_1)(t) - u_1(t)({}_t T_\alpha^a u_2)(t)}{(u_2(t))^2}, \quad u_2(t) \neq 0.$

6)  $({}_t T_\alpha^a u)(t) = (t - a)^{1-\alpha} u'(t)$ , where  $u'(t) = \lim_{\epsilon \rightarrow 0} \left(\frac{u(t+\epsilon) - u(t)}{\epsilon}\right).$

**Proof** See [17]. □

The chain rule is valid for conformable fractional derivatives.

### 3. The fractional Laplace transform

In this section, we recall the basic definitions and some useful facts related to the fractional Laplace transform. The transform, introduced in [17], helps us to solve some conformable fractional differential equations and second-order ODEs. The conformable fractional Laplace transform is defined as follows.

**Definition 3.1** (Abdeljawad [17]) *The conformable fractional Laplace transform (CFLT) of order  $0 < \alpha \leq 1$  of a function  $u : [0, \infty) \rightarrow \mathbb{R}$ , starting from  $a$ , is defined by*

$$L_\alpha^a \{u(t)\} = \int_a^\infty e^{-s \frac{(t-a)^\alpha}{\alpha}} u(t) d_\alpha(t, a) = \int_a^\infty e^{-s \frac{(t-a)^\alpha}{\alpha}} u(t) (t-a)^{\alpha-1} dt = U_\alpha^a(s). \tag{3.1}$$

If  $a=0$ , then

$$L_\alpha^0 \{u(t)\} = \int_0^\infty e^{-s \frac{t^\alpha}{\alpha}} u(t) t^{\alpha-1} dt = U_\alpha^0(s) = U_\alpha(s). \tag{3.2}$$

In particular, if  $\alpha = 1$ , then equation (3.2) is reduced to the definition of the Laplace transform:

$$L \{u(t)\} = \int_0^\infty e^{-st} u(t) dt = U(s). \tag{3.3}$$

**Theorem 3.2** [14]. *If  $u(t)$  is piecewise continuous for  $t > t_0$  if  $|u(t)| \leq v(t)$  when  $t \geq t_0$  for some positive constant  $t_0$  and if  $\int_{t_0}^\infty v(t) d_\alpha t$  converges then  $\int_{t_0}^\infty u(t) d_\alpha t$  also converges. On the other hand if  $u(t) \geq v(t) \geq 0$  for  $t \geq t_0$  and if  $\int_{t_0}^\infty v(t) d_\alpha t$  diverges then  $\int_{t_0}^\infty u(t) d_\alpha t$  also diverges.*

**Theorem 3.3** [14]. *Let an exponential order function  $u(t)$  be piecewise continuous on the interval  $0 \leq t \leq A$  for any positive  $A$  and  $e^{-a \frac{t^\alpha}{\alpha}} |u(t)| < M$  when  $t > t_0$ . In this equality  $M, a, t_0$  are positive real constants and  $0 < \alpha \leq 1$ . Then the conformable Laplace transform defined by 3.1 exists for any  $s > a$ .*

**Theorem 3.4** *Let  $u : [a, \infty) \rightarrow \mathbb{R}$  be a differentiable real-valued function, and  $0 < \alpha \leq 1$ . Then,*

$$L_\alpha^a \{ {}_t T_\alpha^a (u)(t) \} = s U_\alpha^a(s) - u(a). \tag{3.4}$$

**Proof** See [17]. □

**Theorem 3.5** *If  $u$  is piecewise continuous on  $[0, \infty)$  and  $L_\alpha^a \{u(t)\} = U_\alpha^a(s)$ , then*

$$L_\alpha^0 \{ t^{n\alpha} u(t) \} = (-1)^n \alpha^n \frac{d^n}{ds^n} [U_\alpha^0(s)], \quad n \in \mathbb{N}. \tag{3.5}$$

**Proof** See [12]. □

**Example 3.6** We calculate the fractional Laplace transform of  $f(t) = \frac{1}{t}$ , for  $0 < \alpha \leq 1$  and  $\frac{1}{\alpha} \notin \mathbb{N}$ .  
By Definition 3.1,

$$L_{\alpha}^0 \left\{ \frac{1}{t} \right\} = \int_0^{\infty} e^{-s \frac{t^{\alpha}}{\alpha}} \frac{1}{t} t^{\alpha-1} dt.$$

Hence, using the change of variable  $z = \frac{t^{\alpha}}{\alpha}$  we obtain

$$L_{\alpha}^0 \left\{ \frac{1}{t} \right\} = \int_0^{\infty} e^{-sz} \frac{1}{\alpha^{\frac{1}{\alpha}} z^{\frac{1}{\alpha}}} dz = \frac{\alpha^{-\frac{1}{\alpha}}}{s^{1-\frac{1}{\alpha}}} \Gamma \left( 1 - \frac{1}{\alpha} \right).$$

**Example 3.7** If  $0 < \alpha \leq 1$ , then the following equalities hold.

$$1) \int_0^{\infty} e^{-2\sqrt{t}} \sin 2\sqrt{t} dt = \frac{1}{4}.$$

$$2) L_{\alpha}^0 \{ \ln(t) \} = -\frac{1}{\alpha s} \left( \ln\left(\frac{s}{\alpha}\right) + \gamma \right), \quad \gamma = .5772157\dots$$

Using Theorem 3.5 we can write

$$\int_0^{\infty} e^{-s \frac{t^{\alpha}}{\alpha}} f(t) t^{2\alpha-1} dt = \int_0^{\infty} e^{-s \frac{t^{\alpha}}{\alpha}} \left( t^{\alpha} f(t) \right) t^{\alpha-1} dt = -\alpha F'_{\alpha}(s).$$

So, letting  $f(t) = \sin(1 \frac{t^{\alpha}}{\alpha})$  we find that

$$\int_0^{\infty} e^{-s \frac{t^{\alpha}}{\alpha}} \sin\left(\frac{t^{\alpha}}{\alpha}\right) t^{2\alpha-1} dt = \frac{2\alpha s}{(s^2 + 1)^2}.$$

Finally, we let  $s = 1$  and  $\alpha = \frac{1}{2}$  to obtain

$$\int_0^{\infty} e^{-2\sqrt{t}} \sin 2\sqrt{t} dt = \frac{1}{4}.$$

By Definition 3.1 and the change of variable  $z = t^{\alpha}$  we can write

$$L_{\alpha}^0 \{ \ln(t) \} = \int_0^{\infty} e^{-s \frac{t^{\alpha}}{\alpha}} \ln(t) t^{\alpha-1} dt = \int_0^{\infty} e^{-\frac{s}{\alpha} z} \left( \frac{1}{\alpha} \ln(z) \right) \frac{1}{\alpha} dz = -\frac{1}{\alpha s} \left( \ln\left(\frac{s}{\alpha}\right) + \gamma \right).$$

#### 4. Basic properties of converting CFDEs to first order ODEs

In this section, we use the fractional Laplace transform for solving a class of conformable fractional differential equations. This is done by taking the fractional Laplace transform of such equations and converting them into first-order ODEs.

**Theorem 4.1** [13]. Let  $u : [a, \infty) \rightarrow \mathbb{R}$  be twice differentiable on  $(a, \infty)$ ,  $\alpha, \beta > 0$  and  $\alpha + \beta \leq 1$ . Then,

$${}_t T_{\beta} [ {}_t T_{\alpha} u(t) ] = (1 - \alpha) t^{1-(\alpha+\beta)} u'(t) + t^{2-(\alpha+\beta)} u''(t). \tag{4.1}$$

**Theorem 4.2** [13]. If  $u : [a, \infty) \rightarrow \mathbb{R}$  is twice differentiable on  $(a, \infty)$ ,  $\alpha, \beta > 0$  and  $\alpha + \beta \leq 1$ , then the following equalities hold.

- 1)  $\int_0^\infty e^{-s\frac{t^{\alpha+\beta}}{\alpha+\beta}} (tu''(t))dt = u(0) + s \int_0^\infty t^{\alpha+\beta} u'(t)e^{-s\frac{t^{\alpha+\beta}}{\alpha+\beta}} dt - sU_{(\alpha+\beta)}(s).$
- 2)  $\int_0^\infty (1-\alpha+st^{\alpha+\beta})u'(t)e^{-s\frac{t^{\alpha+\beta}}{\alpha+\beta}} dt = (\alpha-1)u(0) - (\alpha+\beta)sU_{(\alpha+\beta)}(s) + (1-\alpha)sU_{(\alpha+\beta)}(s) - (\alpha+\beta)s^2U'_{(\alpha+\beta)}(s).$
- 3)  $L^0_{(\alpha+\beta)} \left\{ {}_tT_\beta \left( {}_tT_\alpha u(t) \right) \right\} = \alpha u(0) - (2\alpha + \beta)sU_{(\alpha+\beta)}(s) - (\alpha + \beta)s^2U'_{(\alpha+\beta)}(s).$
- 4)  $L^0_{(\alpha+\beta)} \left\{ {}_tT_\beta \left( {}_tT_\alpha u(t) \right) + {}_tT_\alpha \left( {}_tT_\beta u(t) \right) \right\} = (\alpha + \beta)u(0) - (3\alpha + 3\beta)sU_{(\alpha+\beta)}(s) - (2\alpha + 2\beta)s^2U'_{(\alpha+\beta)}(s).$

**Theorem 4.3** [13]. If  $s, \alpha, \beta > 0$  and  $\alpha + \beta \leq 1$ , then the following are true.

- 1)  $L^0_{\alpha+\beta} \left\{ t^{-\beta} \left( {}_tT_\alpha u(t) \right) \right\} = -u(0) + sU_{(\alpha+\beta)}(s).$
- 2)  $L^0_{\alpha+\beta} \left\{ t^\alpha \left( {}_tT_\alpha u(t) \right) \right\} = L^0_{\alpha+\beta} \left\{ tu'(t) \right\} = L^0_{\alpha+\beta} \left\{ t^{\alpha+\beta} \left( {}_tT_{\alpha+\beta} u(t) \right) \right\} = -(\alpha+\beta)U_{(\alpha+\beta)}(s) - (\alpha+\beta)sU'_{(\alpha+\beta)}(s).$

**Proposition 4.4** If  $s > 0$  and  $0 < \alpha \leq 1$  then

$$\int_0^\infty e^{-s\frac{t^\alpha}{\alpha}} \left( \frac{{}_tT_\alpha f(t)}{t} \right) t^{2\alpha} dt = -\alpha F_\alpha(s) - \alpha s F'_\alpha(s).$$

**Proof** By Theorem 3.4 and Theorem 3.5,

$$\int_0^\infty e^{-s\frac{t^\alpha}{\alpha}} \left( \frac{{}_tT_\alpha f(t)}{t} \right) t^{2\alpha} dt = \int_0^\infty e^{-s\frac{t^\alpha}{\alpha}} \left( t^\alpha {}_tT_\alpha f(t) \right) t^{\alpha-1} dt = -\alpha \frac{d}{ds} \left( sF_\alpha(s) - f(0) \right) = -\alpha F_\alpha(s) - \alpha s F'_\alpha(s).$$

□

**Proposition 4.5** Let an exponential order function  $u(t)$  be twice differentiable on  $(0, \infty)$  and  $\alpha, \beta > 0$  such that  $\alpha + \beta \leq 1$ , then, the solution of the CFDE

$${}_tT_\beta \left( {}_tT_\alpha u(t) \right) + \left( t^\alpha + t^{-\beta} \right) \left( {}_tT_\alpha u(t) \right) = q(t) \tag{4.2}$$

is given by

$$u(t) = (L^0_{\alpha+\beta})^{-1} \left\{ \left( \frac{(s+1)^{\frac{1-\alpha}{\alpha+\beta}}}{s} \right) \left( \int \frac{(\alpha-1)u(0) - Q_{(\alpha+\beta)}(s)}{(\alpha+\beta)(s+1)^{\frac{\beta+1}{\alpha+\beta}}} ds + C \right) \right\}.$$

**Proof** Applying  $L^0_{\alpha+\beta}$  to the both sides of (4.2), and using Theorem 4.2 and Theorem 4.3, we obtain

$$\begin{aligned} \alpha u(0) - (2\alpha + \beta)sU_{(\alpha+\beta)}(s) - (\alpha + \beta)s^2U'_{(\alpha+\beta)}(s) - (\alpha + \beta)U_{(\alpha+\beta)}(s) - (\alpha + \beta)sU'_{(\alpha+\beta)}(s) - u(0) + sU_{(\alpha+\beta)}(s) \\ = Q_{(\alpha+\beta)}(s). \end{aligned}$$

Therefore, we can write

$$U'_{(\alpha+\beta)}(s) + \left( \frac{\alpha - 1}{(\alpha + \beta)(s + 1)} + \frac{1}{s} \right) U_{(\alpha+\beta)}(s) = \frac{(\alpha - 1)u(0) - Q_{(\alpha+\beta)}(s)}{s(s + 1)(\alpha + \beta)}.$$

By solving this first-order differential equation we obtain

$$U_{(\alpha+\beta)}(s) = \left( e^{-\int \left( \frac{\alpha-1}{(\alpha+\beta)(s+1)} + \frac{1}{s} \right) ds} \right) \left( \int \frac{(\alpha - 1)u(0) - Q_{(\alpha+\beta)}(s)}{s(s + 1)(\alpha + \beta)} e^{\int \left( \frac{\alpha-1}{(\alpha+\beta)(s+1)} + \frac{1}{s} \right) ds} ds + C \right).$$

Hence,

$$U_{(\alpha+\beta)}(s) = \left( \frac{(s + 1)^{\frac{1-\alpha}{\alpha+\beta}}}{s} \right) \left( \int \frac{(\alpha - 1)u(0) - Q_{(\alpha+\beta)}(s)}{(\alpha + \beta)(s + 1)^{\frac{\beta+1}{\alpha+\beta}}} ds + C \right).$$

Now, the solution  $u(t)$  can be found by applying the CF inverse transform. □

*In particular, when  $\alpha = \beta = \frac{1}{2}$  and  $u(0)=0$ , the solution of the CFDE*

$${}_tT_{\frac{1}{2}} \left( {}_tT_{\frac{1}{2}} u(t) \right) + \left( \sqrt{t} + \frac{1}{\sqrt{t}} \right) \left( {}_tT_{\frac{1}{2}} u(t) \right) = 5t + 2t^2$$

is given by

$$u(t) = (L_{(\frac{1}{2}+\frac{1}{2})}^0)^{-1} \{ U_{(\frac{1}{2}+\frac{1}{2})}(s) \} = L^{-1} \left\{ \frac{2}{s^3} + C \left( \frac{\sqrt{s+1}}{s} \right) \right\} = t^2 + C \left( \frac{e^{-t}}{\sqrt{\pi t}} + erf(\sqrt{t}) \right).$$

Since  $u(0) = 0$ ,  $u(t) = t^2$ .

**Theorem 4.6** *Let  $s > 0$ ,  $0 < \alpha \leq 1$  and an exponential order function  $g(t)$  be twice differentiable on  $(0, \infty)$  such that  $L_{\alpha}^0 \{g(t)\} = G_{\alpha}(s)$ , then*

$$L_{\alpha}^0 \left\{ {}_tT_{\alpha} g(t) + t^{2-\alpha} g''(t) \right\} = -\alpha s G_{\alpha}(s) - \alpha s^2 G'_{\alpha}(s). \tag{4.3}$$

**Proof** By (3.5),

$$L_{\alpha}^0 \left\{ f(t) + \frac{t}{\alpha} f'(t) \right\} = L_{\alpha}^0 \{f(t)\} + \frac{1}{\alpha} L_{\alpha}^0 \left\{ t^{\alpha} \left( {}_tT_{\alpha} f(t) \right) \right\} = F_{\alpha}(s) + \frac{1}{\alpha} (-\alpha) \left( s F_{\alpha}(s) - f(0) \right)' = -s F'_{\alpha}(s).$$

Now, letting  $f(t) = {}_tT_{\alpha} g(t)$  and using Theorem 3.4 we can write

$$L_{\alpha}^0 \left\{ {}_tT_{\alpha} g(t) + \frac{t}{\alpha} \left( {}_tT_{\alpha} g(t) \right)' \right\} = -s \left( s G_{\alpha}(s) - g(0) \right)' = -s G_{\alpha}(s) - s^2 G'_{\alpha}(s).$$

Since  ${}_tT_{\alpha} g(t) + \frac{t}{\alpha} \left( {}_tT_{\alpha} g(t) \right)' = \frac{1}{\alpha} t^{1-\alpha} g'(t) + \frac{1}{\alpha} t^{2-\alpha} g''(t)$ ,

$$L_{\alpha}^0 \left\{ {}_tT_{\alpha} g(t) + t^{2-\alpha} g''(t) \right\} = L_{\alpha}^0 \left\{ t^{1-\alpha} g'(t) + t^{2-\alpha} g''(t) \right\} = -\alpha s G_{\alpha}(s) - \alpha s^2 G'_{\alpha}(s).$$

□

Therefore, if  $\alpha, \beta > 0$  and  $\alpha + \beta \leq 1$ , then

$$\begin{aligned} L_{\alpha+\beta}^0 \left\{ \left( {}_tT_{(\alpha+\beta)}u(t) \right) + t^{2-(\alpha+\beta)}u''(t) \right\} &= L_{\alpha+\beta}^0 \left\{ t^{1-(\alpha+\beta)}u'(t) + t^{2-(\alpha+\beta)}u''(t) \right\} \\ &= -(\alpha + \beta)sU_{(\alpha+\beta)}(s) - (\alpha + \beta)s^2U'_{(\alpha+\beta)}(s). \end{aligned}$$

**Proposition 4.7** Suppose that  $u(t)$  is twice differentiable on  $(0, \infty)$ ,  $\alpha, \beta > 0$  and  $\alpha + \beta \leq 1$ . Then, the solution of the CFDE

$${}_tT_\beta \left( {}_tT_\alpha u(t) \right) + \left( {}_tT_{\alpha+\beta}u(t) \right) + t^{2-(\alpha+\beta)}u''(t) = q(t) \tag{4.4}$$

is given by

$$u(t) = \left( L_{\alpha+\beta}^0 \right)^{-1} \left\{ \left( \frac{1}{s^{\frac{3\alpha+2\beta}{2\alpha+2\beta}}} \right) \left( \int \frac{\alpha u(0) - Q_{\alpha+\beta}(s)}{2(\alpha + \beta)s^{\frac{\alpha+2\beta}{2\alpha+2\beta}}} ds + C \right) \right\}.$$

**Proof** Applying  $L_{\alpha+\beta}^0$  to the both sides of (4.4), and using Theorem 4.2 and Theorem 4.6, we find that

$$\alpha u(0) - (2\alpha + \beta)sU_{(\alpha+\beta)}(s) - s^2(\alpha + \beta)U'_{(\alpha+\beta)}(s) - (\alpha + \beta)sU_{(\alpha+\beta)}(s) - (\alpha + \beta)s^2U'_{(\alpha+\beta)}(s) = Q_{(\alpha+\beta)}(s).$$

This differential equation can be written as

$$2(\alpha + \beta)s^2U'_{(\alpha+\beta)}(s) + (3\alpha + 2\beta)sU_{(\alpha+\beta)}(s) = \alpha u(0) - Q_{(\alpha+\beta)}(s).$$

Therefore, we can write

$$U'_{(\alpha+\beta)}(s) + \left( \frac{\alpha}{(2\alpha + 2\beta)s} + \frac{1}{s} \right) U_{(\alpha+\beta)}(s) = \frac{\alpha u(0) - Q_{(\alpha+\beta)}(s)}{(2\alpha + 2\beta)s^2}.$$

By solving this first-order ODE we obtain

$$\begin{aligned} U_{\alpha+\beta}(s) &= \left( e^{-\int \left( \frac{\alpha}{(2\alpha+2\beta)s} + \frac{1}{s} \right) ds} \right) \left( \int \frac{\alpha u(0) - Q_{\alpha+\beta}(s)}{(2\alpha + 2\beta)s^2} e^{\int \left( \frac{\alpha}{(2\alpha+2\beta)s} + \frac{1}{s} \right) ds} ds + C \right) \\ &= \frac{1}{s^{\frac{3\alpha+2\beta}{2\alpha+2\beta}}} \left( \int \frac{\alpha u(0) - Q_{\alpha+\beta}(s)}{2(\alpha + \beta)s^{\frac{\alpha+2\beta}{2\alpha+2\beta}}} ds + C \right). \end{aligned}$$

Now, the solution  $u(t)$  can be found by applying the CF inverse transform. □

In particular, when  $\alpha = \beta = \frac{1}{2}$  and  $u(0)=1$ , the solution of the CFDE

$${}_tT_{\frac{1}{2}} \left( {}_tT_{\frac{1}{2}}u(t) \right) + {}_tT_{\frac{1}{2}+\frac{1}{2}}u(t) + t^{2-(\frac{1}{2}+\frac{1}{2})}u''(t) = \left( \frac{3}{2} + 2t \right) e^t$$

is given by

$$u(t) = L^{-1} \left\{ \frac{1}{s^{1+\frac{1}{4}}} \left( \frac{s^{\frac{5}{4}}}{s-1} + C \right) \right\} = L^{-1} \left\{ \frac{1}{s-1} \right\} + CL^{-1} \left\{ \frac{1}{s^{1+\frac{1}{4}}} \right\} = e^t + C \left( \frac{2\sqrt{2}\Gamma(\frac{3}{4})t^{\frac{1}{4}}}{\pi} \right).$$



**Theorem 4.8** *If  $s, \alpha > 0$  and  $\alpha \leq 1$ , then*

$$L_{\alpha}^0\{t^{\alpha+1}u'(t)\} = 2\alpha^2U'_{\alpha}(s) + \alpha^2sU''_{\alpha}(s). \tag{4.5}$$

**Proof** By Theorem 4.3,

$$L_{\alpha}^0\{tu'(t)\} = -\alpha U_{\alpha}(s) - \alpha sU'_{\alpha}(s).$$

Now, Theorem 3.5 allows us to write

$$L_{\alpha}^0\{t^{\alpha+1}u'(t)\} = L_{\alpha}^0\{t^{\alpha}(tu'(t))\} = -\alpha \frac{d}{ds} \left( -\alpha U_{\alpha}(s) - \alpha sU'_{\alpha}(s) \right) = 2\alpha^2U'_{\alpha}(s) + \alpha^2sU''_{\alpha}(s).$$

□

**Proposition 4.9** *Consider the CFDE*

$${}_tT_{\beta} \left( {}_tT_{\alpha}u(t) \right) - {}_tT_{\alpha} \left( {}_tT_{\beta}u(t) \right) = 0. \tag{4.6}$$

*If  $\alpha, \beta > 0$  and  $\alpha + \beta \leq 1$ , then  $\alpha = \beta$  or  $u(t)$  is constant.*

**Proof** Applying  $L_{\alpha+\beta}^0$  and using Theorem 4.2 we obtain

$$\alpha u(0) - (2\alpha + \beta)sU_{(\alpha+\beta)}(s) - (\alpha + \beta)s^2U'_{(\alpha+\beta)}(s) - \beta u(0) + (2\beta + \alpha)sU_{(\alpha+\beta)}(s) + (\alpha + \beta)s^2U'_{(\alpha+\beta)}(s) = 0.$$

Therefore,

$$(\alpha - \beta) \left( u(0) - sU_{(\alpha+\beta)}(s) \right) = 0.$$

Therefore,

$$(\alpha = \beta) \text{ or } U_{(\alpha+\beta)}(s) = \frac{u(0)}{s}.$$

Finally, we apply  $(L_{\alpha+\beta}^0)^{-1}$  to obtain

$$u(t) = \left( L_{\alpha+\beta}^0 \right)^{-1} \left\{ U_{(\alpha+\beta)}(s) \right\} = \left( L_{\alpha+\beta}^0 \right)^{-1} \left\{ \frac{u(0)}{s} \right\} = u(0).$$

□

**Proposition 4.10** *Assume that  $u(t)$  is twice differentiable on  $(0, \infty)$ ,  $k, m \in \mathbb{R}$ ,  $\alpha, \beta > 0$  and  $\alpha + \beta \leq 1$ . Then, the solution of the CFDE*

$${}_tT_{\beta}[{}_tT_{\alpha}u(t)] + {}_tT_{\alpha}[{}_tT_{\beta}u(t)] + ktu'(t) + mu(t) = q(t) \tag{4.7}$$

*is given by*

$$u(t) = \left( L_{\alpha+\beta}^0 \right)^{-1} \left\{ \left( \frac{s^{\frac{m}{k(\alpha+\beta)} - 1}}{(2s+k)^{\frac{m}{k(\alpha+\beta)} + \frac{1}{2}}} \right) \left( \int \frac{(\alpha + \beta)u(0) - Q_{(\alpha+\beta)}(s)}{(\alpha + \beta)s^{\frac{m}{k(\alpha+\beta)}}} (2s+k)^{\frac{2m-k(\alpha+\beta)}{2k(\alpha+\beta)}} ds + C \right) \right\}.$$

**Proof** Applying the CF Laplace transform to the both sides of (4.7), and using Theorem 4.2 and Theorem 4.3, we obtain

$$(\alpha + \beta)u(0) - (3\alpha + 3\beta)sU_{\alpha+\beta}(s) - (2\alpha + 2\beta)s^2U'_{\alpha+\beta}(s) - k(\alpha + \beta)\left(U_{(\alpha+\beta)}(s) + sU'_{(\alpha+\beta)}(s)\right) + mU_{(\alpha+\beta)}(s) = Q_{(\alpha+\beta)}(s).$$

Equivalently,

$$U'_{(\alpha+\beta)}(s) + \left(\frac{3(\alpha + \beta)s + k(\alpha + \beta) - m}{(\alpha + \beta)s(2s + k)}\right)U_{(\alpha+\beta)}(s) = \frac{(\alpha + \beta)u(0) - Q_{(\alpha+\beta)}(s)}{(\alpha + \beta)s(2s + k)}.$$

By solving this first-order ODE we can write

$$\begin{aligned} U_{(\alpha+\beta)}(s) &= \left(e^{-\int \left(\frac{3(\alpha+\beta)s+k(\alpha+\beta)-m}{(\alpha+\beta)s(2s+k)}\right)ds}\right) \left(\int \frac{(\alpha + \beta)u(0) - Q_{(\alpha+\beta)}(s)}{(\alpha + \beta)s(2s + k)} e^{\int \left(\frac{3(\alpha+\beta)s+k(\alpha+\beta)-m}{(\alpha+\beta)s(2s+k)}\right)ds} ds + C\right) \\ &= \left(\frac{s^{\frac{m}{k(\alpha+\beta)}-1}}{(2s + k)^{\frac{m}{k(\alpha+\beta)}+\frac{1}{2}}}\right) \left(\int \frac{(\alpha + \beta)u(0) - Q_{(\alpha+\beta)}(s)}{(\alpha + \beta)s^{\frac{m}{k(\alpha+\beta)}}} (2s + k)^{\frac{2m-k(\alpha+\beta)}{2k(\alpha+\beta)}} ds + C\right). \end{aligned}$$

Now, the solution  $u(t)$  can be found by applying the CF inverse transform. □

Therefore the solution may not be unique. For example, if  $\alpha = \frac{1}{3}$ ,  $\beta = \frac{2}{3}$  and  $k = m = 1$ , then the conformable-type problem

$${}_tT_{\frac{2}{3}}\left({}_tT_{\frac{1}{3}}u(t)\right) + {}_tT_{\frac{1}{3}}\left({}_tT_{\frac{2}{3}}u(t)\right) + tu'(t) + u(t) = 4t^3 + 15t^2, \quad u(0) = 0$$

has the solution given by

$$\begin{aligned} u(t) &= (L^0_{\frac{1}{3}+\frac{2}{3}})^{-1} \left\{ \left(\frac{s^{\frac{1}{1(\frac{1}{3}+\frac{2}{3})}-1}}{(2s + 1)^{\frac{1}{1(\frac{1}{3}+\frac{2}{3})}+\frac{1}{2}}}\right) \left(\int \frac{(\frac{1}{3} + \frac{2}{3})(0) - \frac{24}{s^4} - \frac{30}{s^3}}{(\frac{1}{3} + \frac{2}{3})s^{\frac{1}{1(\frac{1}{3}+\frac{2}{3})}}} (2s + 1)^{\frac{2(1)-1(\frac{1}{3}+\frac{2}{3})}{2(1)(\frac{1}{3}+\frac{2}{3})}} ds + C\right) \right\} \\ &= (L_1)^{-1} \left\{ \frac{1}{(2s + 1)^{\frac{3}{2}}} \left(\int \left(\frac{-24}{s^5} - \frac{30}{s^4}\right)(2s + 1)^{\frac{1}{2}} ds + C\right) \right\} \\ &= L^{-1} \left\{ \frac{1}{(2s + 1)^{\frac{3}{2}}} \left(\frac{6(2s + 1)^{\frac{3}{2}}}{s^4} + C\right) \right\} \\ &= L^{-1} \left\{ \frac{6}{s^4} \right\} + CL^{-1} \left\{ \frac{1}{(2s + 1)^{\frac{3}{2}}} \right\} \\ &= t^3 + C \left(\frac{\sqrt{2}}{2} \sqrt{\frac{t}{\pi}} e^{-\frac{1}{2}t}\right). \end{aligned}$$

### 5. Solving a class of second-order ODEs via the fractional Laplace transform

In this section, we use the fractional Laplace transform to solve some second-order ODEs.

**Proposition 5.1** *The solution of the second-order ODE*

$$2tu''(t) + (t + 1)u'(t) + u(t) = q(t) \tag{5.1}$$

is given by

$$u(t) = L^{-1} \left\{ \frac{1}{(2s + 1)^{\frac{3}{2}}} \left( \int \frac{u(0) - Q(s)}{s} (2s + 1)^{\frac{1}{2}} ds + C \right) \right\}.$$

**Proof** Equation (5.1) can be rewritten as

$$\frac{2}{3}u'(t) + tu''(t) + \frac{1}{3}u'(t) + tu''(t) + tu'(t) + u(t) = q(t),$$

or equivalently,

$$\left(1 - \frac{1}{3}\right)t^{1 - (\frac{1}{3} + \frac{2}{3})}u'(t) + t^{2 - (\frac{1}{3} + \frac{2}{3})}u''(t) + \left(1 - \frac{2}{3}\right)t^{1 - (\frac{2}{3} + \frac{1}{3})}u'(t) + t^{2 - (\frac{2}{3} + \frac{1}{3})}u''(t) + tu'(t) + u(t) = q(t).$$

Using (4.1) we can write

$${}_tT_{\frac{2}{3}} \left( {}_tT_{\frac{1}{3}}u(t) \right) + {}_tT_{\frac{1}{3}} \left( {}_tT_{\frac{2}{3}}u(t) \right) + 1tu'(t) + 1u(t) = q(t).$$

Now, (4.7) gives us

$$\begin{aligned} u(t) &= (L_1^0)^{-1} \left\{ \frac{1}{(2s + 1)^{\frac{3}{2}}} \left( \int \frac{u(0) - Q_1(s)}{s} (2s + 1)^{\frac{1}{2}} ds + C \right) \right\} \\ &= L^{-1} \left\{ \frac{1}{(2s + 1)^{\frac{3}{2}}} \left( \int \frac{u(0) - Q(s)}{s} (2s + 1)^{\frac{1}{2}} ds + C \right) \right\}. \end{aligned}$$

□

*For example, the solution of the differential equation*

$$2tu''(t) + (t + 1)u'(t) + u(t) = te^{-t}, \quad u(0) = 1$$

is given by

$$\begin{aligned} u(t) &= (L_1^0)^{-1} \left\{ \frac{1}{(2s + 1)^{\frac{3}{2}}} \left( \int \frac{1 - \frac{1}{(s+1)^2}}{s} (2s + 1)^{\frac{1}{2}} ds + C \right) \right\} = L^{-1} \left\{ \frac{1}{(2s + 1)^{\frac{3}{2}}} \left( \frac{(2s + 1)^{\frac{3}{2}}}{s + 1} + C \right) \right\} \\ &= L^{-1} \left\{ \frac{1}{s + 1} \right\} + L^{-1} \left\{ \frac{C}{(2s + 1)^{\frac{3}{2}}} \right\} \\ &= e^{-t} + C \left( \frac{\sqrt{2}}{2} \sqrt{\frac{t}{\pi}} e^{-\frac{1}{2}t} \right). \end{aligned}$$

**Proposition 5.2** *If  $k, m \in \mathbb{R}$ , then the solution of the second-order ODE*

$$2tu''(t) + (kt + 1)u'(t) + mu(t) = q(t) \tag{5.2}$$

is given by

$$u(t) = L^{-1} \left\{ \left( \frac{s^{\frac{m-k}{k}}}{(2s+k)^{\frac{2m+k}{2k}}} \right) \left( \int \frac{u(0) - Q(s)}{s^{\frac{m}{k}}} (2s+k)^{\frac{2m-k}{2k}} ds + C \right) \right\}.$$

**Proof** Equation (5.2) can be rewritten as

$$\frac{2}{3} u'(t) + tu''(t) + \frac{1}{3} u'(t) + tu''(t) + ktu'(t) + mu(t) = q(t),$$

or equivalently,

$$\left(1 - \frac{1}{3}\right)t^{1-(\frac{1}{3}+\frac{2}{3})}u'(t) + t^{2-(\frac{1}{3}+\frac{2}{3})}u''(t) + \left(1 - \frac{2}{3}\right)t^{1-(\frac{2}{3}+\frac{1}{3})}u'(t) + t^{2-(\frac{2}{3}+\frac{1}{3})}u''(t) + ktu'(t) + mu(t) = q(t).$$

Using (4.1) we can write

$${}_tT_{\frac{2}{3}} \left( {}_tT_{\frac{1}{3}} u(t) \right) + {}_tT_{\frac{1}{3}} \left( {}_tT_{\frac{2}{3}} u(t) \right) + ktu'(t) + mu(t) = q(t).$$

Finally, (4.7) gives us

$$\begin{aligned} u(t) &= (L^0_{(\frac{1}{3}+\frac{2}{3})})^{-1} \left\{ U_{(\frac{1}{3}+\frac{2}{3})}(s) \right\} \\ &= \left( L^0_{(\frac{1}{3}+\frac{2}{3})} \right)^{-1} \left\{ \left( \frac{s^{\frac{m}{k(\frac{1}{3}+\frac{2}{3})}-1}}{(2s+k)^{\frac{m}{k(\frac{1}{3}+\frac{2}{3})}+\frac{1}{2}}} \right) \left( \int \frac{(\frac{1}{3}+\frac{2}{3})u(0) - Q_{\frac{1}{3}+\frac{2}{3}}(s)}{(\frac{1}{3}+\frac{2}{3})s^{\frac{m}{k(\frac{1}{3}+\frac{2}{3})}}} (2s+k)^{\frac{2m-k(\frac{1}{3}+\frac{2}{3})}{2k(\frac{1}{3}+\frac{2}{3})}} ds + C \right) \right\} \\ &= L^{-1} \left\{ \left( \frac{s^{\frac{m-k}{k}}}{(2s+k)^{\frac{2m+k}{2k}}} \right) \left( \int \frac{u(0) - Q(s)}{s^{\frac{m}{k}}} (2s+k)^{\frac{2m-k}{2k}} ds + C \right) \right\}. \end{aligned}$$

□

**Proposition 5.3** Let  $a, b, p \in \mathbb{R}$ , and  $a, b \neq 0$ . Then, the solution of the second-order ODE

$$2atu''(t) + (bt+a)u'(t) + pu(t) = q(t) \tag{5.3}$$

is given by

$$u(t) = L^{-1} \left\{ \left( \frac{s^{\frac{p}{b}-1}}{(2s+\frac{b}{a})^{\frac{p}{b}+\frac{1}{2}}} \right) \left( \int \frac{u(0) - \frac{1}{a}Q(s)}{s^{\frac{p}{b}}} (2s+\frac{b}{a})^{\frac{p}{b}-\frac{1}{2}} ds + C \right) \right\}.$$

**Proof** Equation (5.3) can be rewritten as

$$\frac{2}{3} u'(t) + tu''(t) + \frac{1}{3} u'(t) + tu''(t) + \frac{b}{a} tu'(t) + \frac{p}{a} u(t) = \frac{1}{a} q(t),$$

or equivalently

$$\left(1 - \frac{1}{3}\right)t^{1-(\frac{1}{3}+\frac{2}{3})}u'(t) + t^{2-(\frac{1}{3}+\frac{2}{3})}u''(t) + \left(1 - \frac{2}{3}\right)t^{1-(\frac{2}{3}+\frac{1}{3})}u'(t) + t^{2-(\frac{2}{3}+\frac{1}{3})}u''(t) + \frac{b}{a}tu'(t) + \frac{p}{a}u(t) = \frac{1}{a}q(t).$$

Using (4.1) we can write

$${}_tT_{\frac{2}{3}}\left({}_tT_{\frac{1}{3}}u(t)\right) + {}_tT_{\frac{1}{3}}\left({}_tT_{\frac{2}{3}}u(t)\right) + \frac{b}{a}tu'(t) + \frac{p}{a}u(t) = \frac{1}{a}q(t).$$

Letting  $k = \frac{b}{a}$ ,  $m = \frac{p}{a}$  and  $q(t) = \frac{1}{a}q(t)$  in (4.7) we get

$$u(t) = L^{-1}\left\{\left(\frac{s^{\frac{p}{a}-1}}{(2s + \frac{b}{a})^{\frac{p}{a}+\frac{1}{2}}}\right)\left(\int \frac{u(0) - \frac{1}{a}Q(s)}{s^{\frac{p}{a}}}(2s + \frac{b}{a})^{\frac{p}{a}-\frac{1}{2}}ds + C\right)\right\}.$$

□

In particular, if  $a = b = 2$ ,  $p = 1$  and  $u(0) = 0$ , then the solution of the ODE

$$4tu''(t) + (2t + 2)u'(t) + u(t) = (2 - 17t + 12t^2)e^{-2t}$$

is given by

$$u(t) = te^{-2t} + C\left(\frac{-\sqrt{2}}{2}Ie^{-\frac{1}{2}t}erf\left(\frac{\sqrt{2}}{2}I\sqrt{t}\right)\right).$$

Since  $u(0) = 0$ ,  $C=0$ . So, we conclude that

$$u(t) = te^{-2t}.$$

### 6. Solving some conformable fractional differential equations via the fractional Laplace transform

In this section, we use the fractional Laplace transform to solve the Laguerre fractional differential equation. Also, we obtain some new results on some second-order ODEs and some other conformable fractional differential equations.

**Theorem 6.1** *If  $s$ ,  $\alpha > 0$ ,  $\alpha \leq 1$  and  $L_{\alpha}^0\{u(t)\} = U_{\alpha}(s)$ , then the following equalities hold.*

$$1) L_{\alpha}^0\{t^{2-\alpha}u''(t)\} = (-1 - \alpha)sU_{\alpha}(s) + u(0) - \alpha s^2U'_{\alpha}(s).$$

$$2) L_{\alpha}^0\{t^2u''(t)\} = (\alpha + \alpha^2)U_{\alpha}(s) + \alpha s(1 + 3\alpha)U'_{\alpha}(s) + \alpha^2s^2U''_{\alpha}(s).$$

**Proof** By Theorem 4.6,

$$L_{\alpha}^0\left\{{}_tT_{\alpha}u(t) + t^{2-\alpha}u''(t)\right\} = -\alpha sU_{\alpha}(s) - \alpha s^2U'_{\alpha}(s).$$

Therefore, Theorem 3.4 gives us the first formula:

$$L_{\alpha}^0\{t^{2-\alpha}u''(t)\} = (-1 - \alpha)sU_{\alpha}(s) + u(0) - \alpha s^2U'_{\alpha}(s).$$

Now, using Theorem 3.5 we can write

$$L_{\alpha}^0\left\{t^{\alpha}(t^{2-\alpha}u''(t))\right\} = -\alpha \frac{d}{ds}\left((-1 - \alpha)sU_{\alpha}(s) + u(0) - \alpha s^2U'_{\alpha}(s)\right).$$

So, we obtain

$$L_{\alpha}^0 \{t^2 u''(t)\} = (\alpha + \alpha^2)U_{\alpha}(s) + \alpha s(1 + 3\alpha)U'_{\alpha}(s) + \alpha^2 s^2 U''_{\alpha}(s),$$

which completes the proof. □

Therefore, if  $\alpha, \beta > 0, \alpha + \beta \leq 1$  and  $m \in \mathbb{R}$ , then we obtain the following equalities.

- 1)  $L_{\alpha+\beta}^0 \{t^{1-(\alpha+\beta)} u'(t)\} = L_{\alpha+\beta}^0 \{T_{\alpha+\beta} u(t)\} = -u(0) + sU_{\alpha+\beta}(s).$
- 2)  $L_{\alpha+\beta}^0 \{t^{2-(\alpha+\beta)} u''(t)\} = (-1 - (\alpha + \beta))sU_{\alpha+\beta}(s) + u(0) - (\alpha + \beta)s^2 U'_{\alpha+\beta}(s).$
- 3)  $L_{\alpha+\beta}^0 \{(m + 1)t^{1-(\alpha+\beta)} u'(t) + t^{2-(\alpha+\beta)} u''(t)\} = -mu(0) - (\alpha + \beta - m)sU_{\alpha+\beta}(s) - (\alpha + \beta)s^2 U'_{\alpha+\beta}(s).$

**Theorem 6.2** *If  $s, \alpha, \beta, \gamma > 0$  and  $\alpha + \beta + \gamma \leq 1$ , then the following equalities hold.*

$$I) \int_0^{\infty} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} (t^2 u'''(t)) dt = -2 \int_0^{\infty} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} (tu''(t)) dt + s \int_0^{\infty} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} (t^{\alpha+\beta+\gamma+1} u''(t)) dt.$$

$$\begin{aligned} II) & \int_0^{\infty} \left( (1 - 2\alpha - \beta)t + st^{\alpha+\beta+\gamma+1} \right) e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u''(t) dt \\ &= - \int_0^{\infty} \left( (1 - 2\alpha - \beta) + s(\alpha + \beta + \gamma + 1)t^{\alpha+\beta+\gamma} \right) \left( e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u'(t) \right) dt \\ & \quad + s(1 - 2\alpha - \beta) \int_0^{\infty} t^{\alpha+\beta+\gamma} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u'(t) dt + s^2 \int_0^{\infty} t^{2\alpha+2\beta+2\gamma} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u'(t) dt. \end{aligned}$$

**Proof** Using integration by parts we can write

$$\begin{aligned} & \int_0^{\infty} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} (t^2 u'''(t)) dt \\ &= t^2 e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u''(t) \Big|_0^{\infty} - \int_0^{\infty} u''(t) (2t - st^{\alpha+\beta+\gamma+1}) e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} dt \\ &= 0 - 2 \int_0^{\infty} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} (tu''(t)) dt + s \int_0^{\infty} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} (t^{\alpha+\beta+\gamma+1} u''(t)) dt. \end{aligned}$$

Also,

$$\begin{aligned} & \int_0^{\infty} \left( (1 - 2\alpha - \beta)t + st^{\alpha+\beta+\gamma+1} \right) e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u''(t) dt \\ &= \left( (1 - 2\alpha - \beta)t + st^{\alpha+\beta+\gamma} \right) e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u'(t) \Big|_0^{\infty} \\ & \quad - \int_0^{\infty} \left\{ (1 - 2\alpha - \beta) + s(\alpha + \beta + \gamma + 1)t^{\alpha+\beta+\gamma} \right\} \left( e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u'(t) \right) dt \end{aligned}$$

$$\begin{aligned}
 & +s(1-2\alpha-\beta) \int_0^\infty t^{\alpha+\beta+\gamma} \left( e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} \right) u'(t) dt + s^2 \int_0^\infty t^{2\alpha+2\beta+2\gamma} \left( e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} \right) u'(t) dt \\
 = & 0 - \int_0^\infty \left\{ (1-2\alpha-\beta) + s(\alpha+\beta+\gamma+1)t^{\alpha+\beta+\gamma} \right\} \left( e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u'(t) \right) dt \\
 & +s(1-2\alpha-\beta) \int_0^\infty t^{\alpha+\beta+\gamma} \left( e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} \right) u'(t) dt + s^2 \int_0^\infty t^{2\alpha+2\beta+2\gamma} \left( e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} \right) u'(t) dt.
 \end{aligned}$$

□

**Theorem 6.3** *If  $s, \alpha, \beta, \gamma > 0$  and  $\alpha + \beta + \gamma \leq 1$ , then*

$$\begin{aligned}
 & \int_0^\infty \left\{ (\alpha^2 + \alpha\beta) - (3\alpha + 2\beta + \gamma)st^{\alpha+\beta+\gamma} + s^2 t^{2\alpha+2\beta+2\gamma} \right\} \left\{ e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u'(t) \right\} dt \\
 & = -(\alpha^2 + \alpha\beta)u(0) + (3\alpha + 2\beta + \gamma)(\alpha + \beta + \gamma)sU_{\alpha+\beta+\gamma}(s) \\
 & \quad + 2(\alpha + \beta + \gamma)^2 s^2 U'_{\alpha+\beta+\gamma}(s) + (\alpha^2 + \alpha\beta)sU_{\alpha+\beta+\gamma}(s) \\
 & \quad + (3\alpha + 2\beta + \gamma)(\alpha + \beta + \gamma)s^2 U'_{\alpha+\beta+\gamma}(s) + (\alpha + \beta + \gamma)^2 s^3 U''_{\alpha+\beta+\gamma}(s).
 \end{aligned}$$

**Proof** Using integration by parts and Theorem 3.5 we obtain

$$\begin{aligned}
 & \int_0^\infty \left\{ (\alpha^2 + \alpha\beta) - (3\alpha + 2\beta + \gamma)st^{\alpha+\beta+\gamma} + s^2 t^{2\alpha+2\beta+2\gamma} \right\} \left\{ e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u'(t) \right\} dt \\
 = & \left\{ (\alpha^2 + \alpha\beta) - (3\alpha + 2\beta + \gamma)st^{\alpha+\beta+\gamma} + s^2 t^{2\alpha+2\beta+2\gamma} \right\} \left( e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u(t) \right) \Big|_0^\infty \\
 & - \int_0^\infty \left\{ (-3\alpha - 2\beta - \gamma)s(\alpha + \beta + \gamma)t^{\alpha+\beta+\gamma-1} + s^2(2\alpha + 2\beta + 2\gamma)t^{\alpha+\beta+\gamma-1}(t^{\alpha+\beta+\gamma}) \right\} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u(t) dt \\
 & + \int_0^\infty \left\{ st^{\alpha+\beta+\gamma-1} (\alpha^2 + \alpha\beta - s(3\alpha + 2\beta + \gamma)) (t^{\alpha+\beta+\gamma}) + s^3 t^{\alpha+\beta+\gamma-1} (t^{2\alpha+2\beta+2\gamma}) \right\} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u(t) dt \\
 = & (-\alpha^2 - \alpha\beta)u(0) \\
 & + \int_0^\infty (3\alpha + 2\beta + \gamma)s(\alpha + \beta + \gamma)t^{\alpha+\beta+\gamma-1} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u(t) dt \\
 & - s^2 \int_0^\infty (2\alpha + 2\beta + 2\gamma)(t^{\alpha+\beta+\gamma} u(t)) t^{\alpha+\beta+\gamma-1} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} dt \\
 & + s(\alpha^2 + \alpha\beta) \int_0^\infty t^{\alpha+\beta+\gamma-1} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u(t) dt \\
 & - s^2(3\alpha + 2\beta + \gamma) \int_0^\infty (t^{\alpha+\beta+\gamma} u(t)) t^{\alpha+\beta+\gamma-1} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} dt \\
 & + s^3 \int_0^\infty (t^{2\alpha+2\beta+2\gamma} u(t)) t^{\alpha+\beta+\gamma-1} e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} dt
 \end{aligned}$$

$$\begin{aligned}
 &= -(\alpha^2 + \alpha\beta)u(0) \\
 &\quad + (3\alpha + 2\beta + \gamma)(\alpha + \beta + \gamma)sU_{\alpha+\beta+\gamma}(s) \\
 &\quad + 2s^2(\alpha + \beta + \gamma)^2U'_{\alpha+\beta+\gamma}(s) + (\alpha^2 + \alpha\beta)sU_{\alpha+\beta+\gamma}(s) \\
 &\quad + (3\alpha + 2\beta + \gamma)(\alpha + \beta + \gamma)s^2U'_{\alpha+\beta+\gamma}(s) \\
 &\quad + s^3(\alpha + \beta + \gamma)^2U''_{\alpha+\beta+\gamma}(s).
 \end{aligned}$$

□

**Theorem 6.4** *If  $\alpha, \beta, \gamma > 0$  and  $\alpha + \beta + \gamma \leq 1$ , then*

$${}_tT_\gamma \left( {}_tT_\beta ( {}_tT_\alpha u(t) ) \right) = (1 - \alpha)(1 - \alpha - \beta)t^{1-(\alpha+\beta+\gamma)}u'(t) + (3 - (2\alpha + \beta))t^{2-(\alpha+\beta+\gamma)}u''(t) + t^{3-(\alpha+\beta+\gamma)}u'''(t).$$

**Proof** By Theorem 2.2 and Theorem 4.1,

$$\begin{aligned}
 {}_tT_\gamma \left( {}_tT_\beta ( {}_tT_\alpha u(t) ) \right) &= {}_tT_\gamma \left( (1 - \alpha)t^{1-(\alpha+\beta)}u'(t) + t^{2-(\alpha+\beta)}u''(t) \right) \\
 &= t^{1-\gamma} \frac{d}{dt} \left\{ (1 - \alpha)t^{1-(\alpha+\beta)}u'(t) + t^{2-(\alpha+\beta)}u''(t) \right\} \\
 &= (1 - \alpha)(1 - \alpha - \beta)t^{1-(\alpha+\beta+\gamma)}u'(t) + (3 - (2\alpha + \beta))t^{2-(\alpha+\beta+\gamma)}u''(t) + t^{3-(\alpha+\beta+\gamma)}u'''(t).
 \end{aligned}$$

□

**Theorem 6.5** *If  $s, \alpha, \beta, \gamma > 0$  and  $\alpha + \beta + \gamma \leq 1$ , then*

$$\begin{aligned}
 &L^0_{\alpha+\beta+\gamma} \left\{ {}_tT_\gamma \left( {}_tT_\beta ( {}_tT_\alpha u(t) ) \right) \right\} \\
 &= -(\alpha^2 + \alpha\beta)u(0) \\
 &\quad + \left\{ (3\alpha + 2\beta + \gamma)(\alpha + \beta + \gamma) + \alpha^2 + \alpha\beta \right\} sU_{\alpha+\beta+\gamma}(s) \\
 &\quad + (5\alpha + 4\beta + 3\gamma)(\alpha + \beta + \gamma)s^2U'_{\alpha+\beta+\gamma}(s) \\
 &\quad + (\alpha + \beta + \gamma)^2 s^3U''_{\alpha+\beta+\gamma}(s).
 \end{aligned}$$

**Proof** By Theorem 6.4,

$$\begin{aligned}
 &L^0_{\alpha+\beta+\gamma} \left\{ {}_tT_\gamma \left( {}_tT_\beta ( {}_tT_\alpha u(t) ) \right) \right\} \\
 &= L^0_{\alpha+\beta+\gamma} \left\{ (1 - \alpha)(1 - (\alpha + \beta))t^{1-(\alpha+\beta+\gamma)}u'(t) \right\} \\
 &\quad + L^0_{\alpha+\beta+\gamma} \left\{ (3 - (2\alpha + \beta))t^{2-(\alpha+\beta+\gamma)}u''(t) \right\}
 \end{aligned}$$



$$\begin{aligned}
 & +L_{\alpha+\beta+\gamma}^0 \left\{ t^{3-(\alpha+\beta+\gamma)} u'''(t) \right\} \\
 & = \left( (1-\alpha)^2 + \beta(\alpha-1) \right) \int_0^\infty e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} u'(t) dt \\
 & + (3-2\alpha-\beta) \int_0^\infty e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} (tu''(t)) dt \\
 & + \int_0^\infty e^{-s \frac{t(\alpha+\beta+\gamma)}{(\alpha+\beta+\gamma)}} (t^2 u'''(t)) dt.
 \end{aligned}$$

Now, the proof is straightforward using Theorem 6.2 and Theorem 6.3. □

**Proposition 6.6** *If  $\alpha, \beta > 0, \alpha + \beta \leq 1, k, m \in \mathbb{R}$  and  $k \neq 0$ , then the solution of the second-order ODE*

$$t^{2-(\alpha+\beta)} u''(t) + ktu'(t) + mu(t) = q(t) \tag{6.1}$$

is given by

$$u(t) = \left( L_{\alpha+\beta}^0 \right)^{-1} \left\{ \left( \frac{s^{\frac{m}{k(\alpha+\beta)}-1}}{(s+k)^{\frac{m+k}{k(\alpha+\beta)}}} \right) \left( \int \frac{u(0) - Q_{\alpha+\beta}(s)}{(\alpha+\beta)s^{\frac{m}{k(\alpha+\beta)}}} (s+k)^{\frac{m+k}{k(\alpha+\beta)}-1} ds + C \right) \right\}.$$

**Proof** Applying the CF Laplace transform  $L_{\alpha+\beta}^0$  to the both sides of (6.1), and using Theorem 4.3 and Theorem 6.1, we obtain

$$(-1-(\alpha+\beta))sU_{\alpha+\beta}(s) + u(0) - (\alpha+\beta)s^2U'_{\alpha+\beta}(s) + k(-(\alpha+\beta)U_{\alpha+\beta}(s) - (\alpha+\beta)sU'_{\alpha+\beta}(s)) + mU_{\alpha+\beta}(s) = Q_{\alpha+\beta}(s).$$

Equivalently,

$$U'_{\alpha+\beta}(s) + \left( \frac{s + (\alpha + \beta)(s + k) - m}{(\alpha + \beta)s(s + k)} \right) U_{\alpha+\beta}(s) = \frac{u(0) - Q_{\alpha+\beta}(s)}{(\alpha + \beta)s(s + k)}.$$

By solving this first-order ODE we can write

$$\begin{aligned}
 U_{\alpha+\beta}(s) & = \left( e^{-\int \frac{s+(\alpha+\beta)(s+k)-m}{(\alpha+\beta)s(s+k)} ds} \right) \left( \int \frac{u(0) - Q_{\alpha+\beta}(s)}{(\alpha + \beta)s(s + k)} e^{\int \frac{s+(\alpha+\beta)(s+k)-m}{(\alpha+\beta)s(s+k)} ds} ds + C \right) \\
 & = \left( \frac{s^{\frac{m}{k(\alpha+\beta)}-1}}{(s+k)^{\frac{m+k}{k(\alpha+\beta)}}} \right) \left( \int \frac{u(0) - Q_{\alpha+\beta}(s)}{(\alpha + \beta)s^{\frac{m}{k(\alpha+\beta)}}} (s+k)^{\frac{m+k}{k(\alpha+\beta)}-1} ds + C \right).
 \end{aligned}$$

Now, the solution  $u(t)$  can be found by applying the CF inverse transform. □

*For example, we use the fractional Laplace transform to find the solution of the second-order ODE*

$$tu''(t) + ktu'(t) + mu(t) = q(t),$$

where  $k, m \in \mathbb{R}$  and  $k \neq 0$ . To do so, we let  $\alpha = \beta = \frac{1}{2}$  in (6.1) to obtain

$$t^{2-(\frac{1}{2}+\frac{1}{2})} u''(t) + ktu'(t) + mu(t) = q(t).$$

Therefore,

$$u(t) = (L_1^0)^{-1}\{U_1(s)\} = L^{-1}\left\{\left(\frac{s^{\frac{m}{k}-1}}{(s+k)^{\frac{m}{k}+1}}\right)\left(\int(u(0)-Q(s))\left(1+\frac{k}{s}\right)^{\frac{m}{k}}ds+C\right)\right\}.$$

In particular, if  $k = m = 1$  and  $u(0) = 1$ , then the solution of the differential equation

$$tu''(t) + 1tu'(t) + 1u(t) = 2te^t + e^t$$

is given by

$$\begin{aligned} u(t) &= L^{-1}\left\{\left(\frac{1}{(s+1)^2}\right)\left(\int\left(1-\frac{2}{(s-1)^2}-\frac{1}{s-1}\right)\left(1+\frac{1}{s}\right)ds+C\right)\right\} \\ &= e^t + (C-3)te^{-t}. \end{aligned}$$

**Proposition 6.7** If  $\alpha, \beta > 0, \alpha + \beta \leq 1, a, b, m \in \mathbb{R}$  and  $a \neq 0$ , then the solution of the second-order ODE

$$at^{2-(\alpha+\beta)}u''(t) + bt^{1-(\alpha+\beta)}u'(t) + mu(t) = q(t) \tag{6.2}$$

is given by

$$u(t) = (L_{\alpha+\beta}^0)^{-1}\left\{\left(\frac{e^{-\frac{m}{a(\alpha+\beta)}s}}{s^{1-\frac{b-a}{a(\alpha+\beta)}}}\right)\left(\int\frac{(a-b)u(0)-Q_{\alpha+\beta}(s)}{a(\alpha+\beta)s^{1+\frac{b-a}{a(\alpha+\beta)}}}e^{\frac{m}{a(\alpha+\beta)}s}ds+C\right)\right\}.$$

**Proof** Applying the CF Laplace transform  $L_{\alpha+\beta}^0$  to the both sides of (6.2) and using Theorem 6.1 we find that

$$a\left\{\left(-1-(\alpha+\beta)\right)sU_{\alpha+\beta}(s)+u(0)-(\alpha+\beta)s^2U'_{\alpha+\beta}(s)\right\}+b\left\{-u(0)+sU_{\alpha+\beta}(s)\right\}+mU_{\alpha+\beta}(s)=Q_{\alpha+\beta}(s).$$

Therefore,

$$U'_{\alpha+\beta}(s)+\left(\frac{a(\alpha+\beta)s-bs+as-m}{a(\alpha+\beta)s^2}\right)U_{\alpha+\beta}(s)=\frac{(a-b)u(0)-Q_{\alpha+\beta}(s)}{a(\alpha+\beta)s^2}.$$

By solving this first-order ODE we obtain

$$U_{\alpha+\beta}(s)=\left(\frac{e^{-\frac{m}{a(\alpha+\beta)}s}}{s^{1-\frac{b-a}{a(\alpha+\beta)}}}\right)\left(\int\frac{(a-b)u(0)-Q_{\alpha+\beta}(s)}{a(\alpha+\beta)s^{1+\frac{b-a}{a(\alpha+\beta)}}}e^{\frac{m}{a(\alpha+\beta)}s}ds+C\right).$$

Now, the solution  $u(t)$  can be found by applying the CF inverse transform. □

In particular, letting  $\alpha = \beta = \frac{1}{2}$  in (6.2) we observe that the solution of the second-order ODE

$$atu''(t) + bu'(t) + mu(t) = q(t)$$

is given by

$$u(t) = L^{-1}\left\{\left(\frac{e^{-\frac{m}{a(\frac{1}{2}+\frac{1}{2})}s}}{s^{1-\frac{b-a}{a(\frac{1}{2}+\frac{1}{2})}}}\right)\left(\int\frac{(a-b)u(0)-Q_{\frac{1}{2}+\frac{1}{2}}(s)}{a(\frac{1}{2}+\frac{1}{2})s^{1+\frac{b-a}{a(\frac{1}{2}+\frac{1}{2})}}e^{\frac{m}{a(\frac{1}{2}+\frac{1}{2})}s}ds+C\right)\right\}$$

$$= L^{-1} \left\{ \frac{e^{-\frac{m}{as}}}{s^{2-\frac{b}{a}}} \left( \int \frac{(a-b)u(0) - Q(s)}{as^{\frac{b}{a}}} e^{\frac{m}{as}} ds + C \right) \right\}.$$

For example, if  $a = b = m = 1$ , then the solution of the differential equation

$$tu''(t) + u'(t) + u(t) = te^{-t}$$

is given by

$$\begin{aligned} u(t) &= L^{-1} \left\{ \frac{e^{-\frac{1}{s}}}{s} \left( \int -\frac{1}{(s+1)^2 s} e^{\frac{1}{s}} ds + C \right) \right\} = L^{-1} \left\{ \frac{e^{-\frac{1}{s}}}{s} \left( \frac{se^{\frac{1}{s}}}{s+1} + C \right) \right\} = L^{-1} \left\{ \frac{1}{s+1} \right\} + L^{-1} \left\{ C \frac{e^{-\frac{1}{s}}}{s} \right\} \\ &= e^{-t} + CBesselJ(0, 2\sqrt{t}). \end{aligned}$$

**Proposition 6.8** (The Laguerre conformable fractional differential equation) Suppose that  $u(t)$  is twice differentiable on  $(0, \infty)$ ,  $\alpha, \beta > 0$ ,  $\alpha + \beta \leq 1$ ,  $k, m, n \in \mathbb{R}$  and  $k \neq 0$ . Then, the solution of the CFDE

$${}_t T_\beta \left( {}_t T_\alpha u(t) \right) + (\alpha + m) \left( {}_t T_{\alpha+\beta} u(t) \right) + ktu'(t) + (n - m)u(t) = q(t) \tag{6.3}$$

is given by

$$u(t) = \left( L_{\alpha+\beta}^0 \right)^{-1} \left\{ \left( (s+k)^{\frac{(k+1)m-n}{k(\alpha+\beta)}} s^{\frac{n-m}{k(\alpha+\beta)}-1} \right) \left( \int \frac{-mu(0) - Q_{\alpha+\beta}(s)}{(\alpha+\beta)s^{\frac{n-m}{k(\alpha+\beta)}}} (s+k)^{\frac{n-m(k+1)}{k(\alpha+\beta)}-1} ds + C \right) \right\}.$$

**Proof** By Theorem 2.2 and Theorem 4.1, equation (6.3) can be rewritten as

$$t^{2-(\alpha+\beta)} u''(t) + (m+1)t^{1-(\alpha+\beta)} u'(t) + ktu'(t) + (n-m)u(t) = q(t). \tag{6.4}$$

Now, by applying  $L_{\alpha+\beta}^0$  to the both sides of (6.4), and using Theorem 4.3 and Theorem 6.1, we obtain

$$\begin{aligned} -mu(0) - (\alpha + \beta - m)sU_{\alpha+\beta}(s) - (\alpha + \beta)s^2U'_{\alpha+\beta}(s) + k \left\{ -(\alpha + \beta)U_{\alpha+\beta}(s) - (\alpha + \beta)sU'_{\alpha+\beta}(s) \right\} \\ + (n - m)U_{\alpha+\beta}(s) = Q_{\alpha+\beta}(s). \end{aligned}$$

Therefore,

$$U'_{\alpha+\beta}(s) + \left( \frac{(\alpha + \beta - m)s + (\alpha + \beta)k + m - n}{(\alpha + \beta)s(s+k)} \right) U_{\alpha+\beta}(s) = \frac{-mu(0) - Q_{\alpha+\beta}(s)}{(\alpha + \beta)(s)(s+k)}.$$

By solving this first-order ODE we obtain

$$\begin{aligned} U_{\alpha+\beta}(s) &= \left( e^{-\int \left( \frac{(\alpha+\beta-m)s+(\alpha+\beta)k+m-n}{(\alpha+\beta)s(s+k)} \right) ds} \right) \left( \int \frac{-mu(0) - Q_{\alpha+\beta}(s)}{(\alpha + \beta)s(s+k)} e^{\int \left( \frac{(\alpha+\beta-m)s+(\alpha+\beta)k+m-n}{(\alpha+\beta)s(s+k)} \right) ds} ds + C \right) \\ &= \left( (s+k)^{\frac{(k+1)m-n}{k(\alpha+\beta)}} s^{\frac{n-m}{k(\alpha+\beta)}-1} \right) \left( \int \frac{-mu(0) - Q_{\alpha+\beta}(s)}{(\alpha + \beta)s^{\frac{n-m}{k(\alpha+\beta)}}} (s+k)^{\frac{n-m(k+1)}{k(\alpha+\beta)}-1} ds + C \right). \end{aligned}$$

Now, the solution  $u(t)$  can be found by applying the CF inverse transform. □

In particular, if  $\alpha + \beta = 1$ ,  $k = -1$  and  $m, n \in \mathbb{N} \cup \{0\}$  in (6.4), then we obtain the solution of the differential equation

$$tu''(t) + (m + 1 - t)u'(t) + (n - m)u(t) = 0 \quad (6.5)$$

as

$$u(t) = L^{-1} \left\{ (s - 1)^n s^{m-n-1} \left( \int \frac{-mu(0)}{s^{m-n}} (s - 1)^{-n-1} ds + C \right) \right\}.$$

On the other hand, we can find the Laguerre dependent function using equation (6.5):

$$\psi_n^m(t) = \frac{d^m}{dt^m} \psi_n(t) = \frac{d^m}{dt^m} \left( e^t \frac{d^n}{dt^n} (t^n e^{-t}) \right) = L^{-1} \left\{ (s - 1)^n s^{m-n-1} \left( \int \frac{-mu(0)}{s^{m-n}} (s - 1)^{-n-1} ds + C \right) \right\}.$$

For example, letting  $m=0$  in (6.5) we find that the solution of the Laguerre differential equation

$$tu''(t) + (1 - t)u'(t) + nu(t) = 0$$

is given by

$$u(t) = L^{-1} \left\{ C \frac{(s - 1)^n}{s^{n+1}} \right\}.$$

So, we obtain the Laguerre polynomial of degree  $n$ , that is,

$$\psi_n(t) = L^{-1} \left\{ C \frac{(s - 1)^n}{s^{n+1}} \right\}.$$

## 7. Conclusion

The conformable fractional derivative is a new kind of fractional derivative that still needs to be further investigated. We discussed the fractional Laplace transform, as a transform compatible with this type of fractional derivatives. The conformable fractional derivative behaves well in the product rule and the chain rule, while complicated formulas appear in case of the usual fractional calculus. Some new results were reported, which were shown to be useful in the theory of conformable fractional differential equations. The method of fractional Laplace transform, as a powerful approach for extracting the exact solutions of differential equations, was developed for the conformable time fractional differential equations. Using the fractional Laplace transform, we were able to convert some ordinary differential equations and conformable fractional differential equations into first-order ordinary differential equations.

## References

- [1] Abdon A, Dumitru B, Ahmed AL. New properties of conformable derivative. Journal of Open Mathematics 2015; 13 (1): doi: 10.1515/math-2015-0081
- [2] Abdon A, Ilknur K. Chaos in a simple nonlinear system with Atangana–Baleanu derivatives with fractional order. Journal of Chaos, Solitons & Fractals 2016; 89: 447-454. doi: 10.1016/j.chaos.2016.02.012
- [3] Angelo B M. On generalized and fractional derivatives and their applications to classical mechanics. Journal of Physics A: Mathematical and Theoretical 2018; 51 (36): 365204.
- [4] Dumitru B, Jihad H A, Amin J. The fractional model of spring pendulum: New features within different kernels. Proceedings of the Romanian Academy Series A - Mathematics Physics Technical Sciences Information Science 2018; 19 (3): 447-454.

- [5] Dumitru B, Samaneh Sadat S, Amin J, Jihad H A. New features of the fractional Euler-Lagrange equations for a physical system within non-singular derivative operator. *The European Physical Journal Plus* 2019; 134 (4): 181. doi: 10.1140/epjp/i2019-12561-x
- [6] Hamid M, Ali Asghar JA. Two classes of conformable fractional Sturm-Liouville problems: Theory and applications. *Mathematical Methods in the Applied Sciences* 2021; 44 (1): 166-195. doi: 10.1002/mma.6719
- [7] Igor P. *Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications*. Elsevier 1998;
- [8] Keith BO, Jerome S. *The fractional calculus theory and applications of differentiation and integration to arbitrary order*. *Mathematics in Science and Engineering* 1974; 111.
- [9] Manuel D O, JA Tenreiro M. What is a fractional derivative? *Journal of computational Physics* 2015; 293: 4-13. doi: 10.1016/j.jcp.2014.07.019
- [10] Mehmet E, Mohammad M, Mostafa E, Qin Z, Seithuti P M, Anjan B, Milivoj B. Optical soliton perturbation with fractional-temporal evolution by first integral method with conformable fractional derivatives. *Journal of Optik* 2016; 127 (22): 10659-10669. doi: 10.1016/j.ijleo.2016.08.076
- [11] Michele C, Mauro F. A new definition of fractional derivative without singular kernel. *Journal of Progr. Fract. Differ. Appl* 2015; 1 (2): 1-13.
- [12] Mir Sajjad H. Invariant subspaces admitted by fractional differential equations with conformable derivatives. *Journal of Chaos, Solitons & Fractals* 2018; 107: 161-169. doi: 10.1016/j.chaos.2018.01.002
- [13] Mohammad M, Farhad DS, Mohammad J, Yaghoub M. New analytical methods for solving a class of conformable fractional differential equations by fractional Laplace transform. *Computational Methods for Differential Equations* 2022; 10 (2): 396-407. doi: 10.22034/cmde.2021.40834.1775
- [14] Ozan O, Ali K. The analytical solutions for conformable integral equations and integro-differential equations by conformable Laplace transform. *Journal of Opt. Quant. Electron* 2018; 50: 81.
- [15] Roshdi KH, Mohammed AL, Abdelrahman Y, Mohammad S. A new definition of fractional derivative. *Journal of Computational and Applied Mathematics* 2014; 264: 65-70. doi: 10.1016/j.cam.2014.01.002
- [16] Shaobo H, Kehui S, Xiaoyong M, Bo Y, Siwei XU. Numerical analysis of a fractional-order chaotic system based on conformable fractional-order derivative. *The European Physical Journal Plus* 2017; 132 (1): 1-11. doi: 10.1140/epjp/i2017-11306-3
- [17] Thabet A. On conformable fractional calculus. *Journal of Computational and Applied Mathematics* 2015; 279: 57-66. doi: 10.1016/j.cam.2014.10.016
- [18] YÜCEL C, Ali K. The solution of time fractional heat equation with new fractional derivative definition. *8th International Conference on Applied Mathematics, Simulation and Modelling* 2014; 195.
- [19] Yücel C, Dumitru B, Ali K, Orkun T. New exact solutions of Burgers' type equations with conformable derivative. *Waves in Random and complex Media* 2017; 27 (1): 103-116. doi: 10.1080/17455030.2016.1205237
- [20] Yousef ET, Mehrdad L, Mohsen R. Study of B-spline collocation method for solving fractional optimal control problems. *Transactions of the Institute of Measurement and Control* 2021; 0142331220987537. doi: 10.1177/0142331220987537
- [21] Zeyad AL, Nouf AL, Fatimah AL, Raed AL. New theoretical results and applications on conformable fractional Natural transform. *Ain Shams Engineering Journal* 2021; 12 (1): 927-933. doi: 10.1016/j.asej.2020.07.006