

1-1-2023

Novel correlation coefficients for interval-valued Fermatean hesitant fuzzy sets with pattern recognition application

İBRAHİM DEMİR

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

DEMİR, İBRAHİM (2023) "Novel correlation coefficients for interval-valued Fermatean hesitant fuzzy sets with pattern recognition application," *Turkish Journal of Mathematics*: Vol. 47: No. 1, Article 14.

<https://doi.org/10.55730/1300-0098.3355>

Available at: <https://journals.tubitak.gov.tr/math/vol47/iss1/14>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Novel correlation coefficients for interval-valued Fermatean hesitant fuzzy sets with pattern recognition application

İbrahim DEMİR* 

Turkish Statistical Institute (TURKSTAT), Ankara, Turkey

Received: 28.02.2022

Accepted/Published Online: 18.11.2022

Final Version: 13.01.2023

Abstract: A combination of interval-valued Fermatean fuzzy sets with Fermatean hesitant fuzzy elements in the form of interval values is known as an interval-valued Fermatean hesitant fuzzy set. Since Fermatean hesitant fuzzy sets are effective instruments for representing more complex, ambiguous, and hazy information, interval-valued Fermatean hesitant fuzzy sets are expansions of these sets. This investigation will concentrate on four different types of correlation coefficients for Fermatean hesitant fuzzy sets and expand them to include correlation coefficients and weighted correlation coefficients for interval-valued Fermatean hesitant fuzzy sets. Finally, the numerical examples demonstrate the viability and usefulness of the suggested methodologies in decision-making under many criteria.

Key words: Correlation, correlation coefficient, Fermatean hesitant fuzzy set, informational energy, interval-valued Fermatean hesitant fuzzy set

1. Introduction

1.1. Correlation coefficients

The correlation coefficient (KK) is a precise metric used in correlation studies to express the strength of the linear relationship between two variables. By measuring the distance of each data point from the variable mean, the formula indicates how well the link between the variables can be fit to an example line drawn across the data for two variables. A correlation is a sign of cochange. Using this statistical method, we may measure the magnitude and direction of the link between two or more variables. The KK is denoted by r . The KK can have a value between -1 and $+1$, according to mathematics. The sign of this coefficient indicates the direction of the link between the two variables, and its numerical value indicates the strength of the correlation. The reciprocal contact is taken into consideration while interpreting the connection. Correlation is a reciprocal relationship rather than a cause-and-effect relationship that can be used to explain the difference or resemblance between two variables. Due to correlation, the values of the related variables may be maximized if the affecting factors can be controlled or the relationship between the variables can be estimated by looking at the value of one variable.

A KK is a bivariate statistic if it depicts the connection between just two variables and a multivariate statistic if there are more than two variables. There are, thus, many diverse fields of study, from engineering to physics, from medicine to economics. In statistics, finding a KK between any two parameters or variables is quite common. Pearson's KK has been used in statistics research on data analysis and classification, pattern

*Correspondence: idemir@gmail.com

2010 *AMS Mathematics Subject Classification:* 03E72, 91A06, 62C86

recognition, clustering, medical diagnosis, and decision-making. It is shown that conventional correlation cannot handle data with ambiguous problems. The fuzzy logic approach attempts to quantify human perception and cognitive uncertainties. Scientists are used to employing binary logic to analyze data. Human logic is ambiguous and complicated; therefore, using binary logic to study human mental processes causes some distortion. Fuzzy logic is built on the base of human cognitive processes. Fuzzy logic, for instance, is defined as the "modeling of thinking and decision systems that enable people to make consistent and correct judgments in the presence of extensive and inaccurate information." The fuzzy type KK 's have been expanded based on mathematical statistics, statistical KK , and the fuzzy logic method. The KK generated for fuzzy data shows both the strength of the relationship between fuzzy sets as well as whether fuzzy sets are positively or negatively related.

1.2. Uncertainty

Numerous academic fields, including psychology, philosophy, cognitive science, and artificial intelligence, are interested in how humans reason and make decisions in the face of everyday situations. Usually, numerous mathematical and statistical models are used to try to characterize these processes. The issue of decision-making surfaces in this process. Decision-making (DM) is the process of choosing one or more of the available options for behavior when a person or institution is trying to accomplish a certain objective. According to research, while many daily judgments may be made instinctively, complicated and important choices require more than this. Multicriteria decision making (MCDM) is a group of analytical techniques that assesses the benefits and drawbacks of options based on a variety of criteria. To pick one or more alternatives from a group of alternatives with varying qualities by competing criteria or to rank these alternatives, MCDM approaches are employed to help the DM process. In other words, decision-makers use MCDM approaches to rate options with various features by comparing them to a variety of criteria. MCDM is a collection of techniques that are regularly applied at all levels and in all spheres of life.

A key idea in decision-making (DM) difficulties is uncertainty. Unpredictable events characterize uncertainty. Routine choices cannot be discussed under ambiguous circumstances. It is important to consider both the advantages and disadvantages of potential outcomes under unclear circumstances. It is crucial to do a thorough analysis of the environmental influences at this stage. Benefiting from prior experiences and decisions is not always successful when there is ambiguity, even while ultimate judgments are not in doubt. Thanks to Zadeh's notion of "fuzzy sets" (FS) [76], linguistic terms that we unintentionally employ regularly have become "computable". Fuzzy logic enabled the grading system to expand the realm of classical mathematics, which was previously restricted to certainty. This notion resulted in a paradigm shift that spread throughout the world as a result of its successful implementations in everyday situations. An element with a distinctive function is either an element of a set in the traditional sense of the word or it is not. The membership function (MF), which assigns each item a degree of membership in the range $[0, 1]$, in the FS notion, determines whether or not an apple is a member of a set.

The degree to which an element belongs to a set in the FS A is $\rho(A)$, and the degree to which it does not belong is $1 - \rho(A)$. As a result, one is equal to the total of the degrees of belonging and nonbelonging. This circumstance, however, falls short of adequately explaining the ambiguity in several issues. The intuitionistic fuzzy set (IFS) theory, which is an extension of the FS theory, was developed by Atanassov [4] as a result. In IFS theory, the nonmembership degree (ND) is specified in addition to the membership degree (MD), whereas FS theory is modeled to only reveal the membership degree (MD) defined in the range $[0, 1]$. According to IFS theory, MD and ND are both in the $[0, 1]$ region. Yager [71] proposed Pythagorean fuzzy sets (PFS)

and in certain circumstances developed them as an extension of IFSs because IFSs cannot adequately convey uncertainty. PFSs employ the notion that the sum of the squares of MD and ND is less than or equal to 1 for circumstances when decision-making is impossible when MD and ND are added together. In the literature, there is a lot of study on FS and its many expansions ([1, 15, 16, 20, 21, 28, 29, 36, 50, 74]).

1.3. Motivation

Tools like aggregation operators and information measurements are frequently used in decision-making difficulties. Another method for determining the best option is to utilize the KK s measure of the level of dependence between two sets. Using KK s, one may assess how strongly two variables are associated. Since the information is frequently vague, ambiguous, and incomplete in many situations, several academics have developed the KK s in fuzzy contexts. In addition to supplying the correlation for fuzzy information in accordance with conventional statistics, Chiang and Lin [10] provided a technique for KK of FSs. According to the conventional understanding of "KKs," the KK of fuzzy information has been investigated in [44] using a mathematical programming approximation. According to [4], intuitionistic fuzzy set (IFS) results were more complete and exact based on the findings from the FS theory. Both membership degree (MD) and nonmembership degree (ND) are taken into consideration by the IFS theory, and it requires that their sum be one or less than one. There are several applications for the IFS-derived KK s, including DM, cluster analysis, image processing, and pattern recognition ([42, 64–66, 69]). Thanks to Pythagorean fuzzy set (PFS) ([29, 34, 49, 50, 70, 71, 73, 77]), which were developed to address an IFS issue, several DM problems involving Pythagorean fuzzy information have been published in the literature.

Senapati and Yager [53] were the first to propose the Fermat fuzzy set (FFS). The MD and ND in the FFS achieve the property $0 \leq m_A^3 + n_A^3 \leq 1$. When identifying uncertainties, the FFS, a novel idea in the literature, performs better than the IFS and PFS. As an illustration, consider $0.9 + 0.6 > 1$, $0.92 + 0.62 > 1$, and $0.93 + 0.63 > 1$. Some FFS characteristics, score, and accuracy functions are provided in [53]. Additionally, the TOPSIS approach, which is widely used to solve MCDM issues, has been employed to solve FFS. Additionally, Senapati and Yager [53] used the TOPSIS method, which is frequently used in MCDM issues, to solve FFS difficulties. Senapati and Yager [54] continued this work by investigating a number of additional operations including arithmetic mean operations over FFSs in addition to using the FF weighted product model to address MCDM issues. New aggregation operations that are FFS-related are described and their associated attributes are studied in [55]. Shahzadi and Akram [56] created the new aggregated operators and provided a new decision support algorithm for the FFSS. Garg et al. [26] defined new FFS type aggregated operators defined by t-norm and t-conorm. Donghai et al. [13] suggested the concept of FF linguistic term sets. Operations, score, and accuracy functions belonging to these sets are given. In [14], a new similarity measure related to FF linguistic term sets is constructed. The new measurement is a combination of Euclidean distance measure and cosine similarity measure. Kirisci [30] defined FF soft sets and gave the measure of entropy based on FF soft sets. In [31], a new hesitant fuzzy set called the fermatean hesitant fuzzy set is given and some of its properties are investigated. Kirisci and Simsek [33] offer aggregation operations to extend FFHSs to interval-valued Fermatean hesitant fuzzy sets (IVFHFS) and to improve MCGDM methods in IVFHF environments. In [32], the ELECTRE I method is defined with Fermatean fuzzy sets according to the group DM process in which more than one individual interacts at the same time. In [?], various FF reference relations (consistent, incomplete, consistent incomplete, acceptable incomplete) are defined. An additive consistency based on a priority vector is given. In addition, a model is presented to obtain missing decisions in incomplete FF preference relations.

Torra and Narukawa [61] and Torra [62] provide HFSs, which authorize an object’s MD to a set of potential values. Particularly useful in GDM difficulties, HFSs have generated strong solutions for these issues. Because of these qualities, HFSs have gained prominence in the literature. Following that, several interval-valued hesitant fuzzy sets were investigated [2, 3, 5, 7, 9, 12, 27, 40, 41, 48, 51]. *KK*s for HFSs have been studied in [68] and [8]. In these investigations, the hesitant fuzzy element (HFE), which has fewer element cardinalities, has had its possible MDs enlarged by the addition of some extreme values. If the extreme value is distant from the other possible membership values, the extended HFE will be very different from the original HFE. Additionally, it is impossible to determine the *KK*s between two HFS if one HFS’s hesitant MDs are always zero. With respect to the mean of an HFE, Liao et al. [43] established a unique *KK* formula and expanded the *KK*s’ range to the interval $[-1; 1]$. The *KK* between an HFS and other HFSs will still not be calculated when an HFS is characterized by a constant function. The improvement of the counterintuition of the current *KK*s of HFSs in [8, 43, 68] was the main goal of Sun et al.’s [59] study. Its key contribution is to better the weighted *KK*s; however, the general case is not improved. Meng and Chen [45] suggested the *KK*s of HFSs based on IVF measures, and they [46] extended the technique to study the situation. The term is too complicated, though. The *KK*s of dual HFSs were also explored in [63] and [75]. The *KK*s of HF soft sets were discussed by Das et al. [11]. Although there are many *KK*s for HFSs, these notions might lead to a lot of irrational claims.

The originality of this work can be stated as follows. There have been various extensions of the classical *KK*s such as fuzzy *KK*, IF *KK*, and PF *KK*. These extensions have improved the performance of the *KK*s. FFSs can handle problems with ambiguity and incomplete information more efficiently than that of IFSs and PFSs Figure. In this study, the interval-valued Fermatean fuzzy *KK*s were developed considering the hesitant *KK*s, intuitionistic fuzzy *KK*s, Pythagorean fuzzy *KK*s, and Fermatean fuzzy *KK*s studies. Since the $MD^3 + ND^3 \leq 1$ requirement is satisfied for an object in the use of FFSs, there will be the possibility to cover more elements than IFSs and PFSs. Examples of pattern recognition and medical decision-making regarding the new *KK*s are given. *KK*s based on different fuzzy sets given in previous studies were compared with newly proposed *KK*s.




Figure 1. Comparison of space of FMGs, PMGs. and IFGs [53].

In this study, for IVFHFSSs, *KK*s and weighted *KK*s were defined, and some characteristics were examined. The correlation coefficient for PFSs is a particular instance of the correlation coefficient for FFSs. As a result,

IVFHFS's discovery of correlation coefficients provides more alternatives for handling uncertainty. The suggested correlation coefficient is consequently more general than the current ones, making it more appropriate for solving problems in the real world. Illustrative examples such as pattern recognition and medical decision-making to show the practicality and efficiency of the suggested KFs are given.

The organization of this work's structure is as follows. Fundamental ideas and findings from HFSs, FHFSs, and IVFHFSs are given in Section 2. The informational energies, correlations, and KK s and weighted KK s of IVFHFSs are defined in Section 3. Using the weighted KK s recommended in IVFHF environments, numerical examples are given in Section 4 demonstrating the effectiveness of the proposed method. A comparison of the new method with previous methods is made in Section 5 and this section is concluded with a subsection explaining the advantages of the new method.

2. Preliminaries

It will be regarded as the $\mathfrak{X} = \{x_1, x_2, \dots, x_n\}$ initial universe throughout the work.

The intuitionistic fuzzy set (IFS) in \mathfrak{X} is defined:

$$N = \{(x, \zeta_N(x), \eta_N(x)) | x \in \mathfrak{X}\}. \quad (2.1)$$

In this definition, $\zeta_N(x), \eta_N(x) : \mathfrak{X} \rightarrow [0, 1]$ is said to be MD and ND, with $\zeta_N(x) + \eta_N(x) \leq 1$.

The Pythagorean fuzzy set (PFS) is characterized as:

$$N = \{(x, \zeta_N(x), \eta_N(x)) | x \in \mathfrak{X}\}, \quad (2.2)$$

if $\zeta_N(x), \eta_N(x) : \mathfrak{X} \rightarrow [0, 1]$ are MD and ND of element of the $x \in \mathfrak{X}$, with $\zeta_N^2(x) + \eta_N^2(x) \leq 1$.

Fermatean fuzzy set (FFS) is given as:

$$N = \{(x, \zeta_N(x), \eta_N(x)) | x \in \mathfrak{X}\}, \quad (2.3)$$

if $\zeta_N(x), \eta_N(x) : \mathfrak{X} \rightarrow [0, 1]$ are MD and ND of element of the $x \in \mathfrak{X}$, with $\zeta_N^3(x) + \eta_N^3(x) \leq 1$.

The set

$$\Gamma = \{(u, \tau_\Gamma(u)) : u \in U\} \quad (2.4)$$

is called hesitant fuzzy set, where $\tau_\Gamma(u)$ indicates the set of some values in unit interval, that is probable MD of $u \in U$ to Γ [67].

The set

$$F_T = \{(u, \zeta_{F_T}(u), \eta_{F_T}(u)) : u \in U\} \quad (2.5)$$

is called a Fermatean hesitant fuzzy set (FHFS) [31], where

(i.) For each element $u \in U$, $\zeta_{F_T}(u)$, $\eta_{F_T}(u)$ are functions from U to $[0, 1]$, demonstrating a likely MD and ND of element $u \in U$ in F_T respectively,

(ii.) $\forall t_{F_T}(u) \in \zeta_{F_T}(u)$, $\exists t'_{F_T}(u) \in \eta_{F_T}(u)$, such that $0 \leq t_{F_T}^3(u) + t'_{F_T}{}^3(u) \leq 1$,

(iii.) $\forall t'_{F_T}(u) \in \eta_{F_T}(u), \exists t_{F_T}(u) \in \zeta_{F_T}(u)$, such that $0 \leq t_{F_T}^3(u) + t'_{F_T}{}^3(u) \leq 1$.

From this stage on, the set of all elements belonging to FHFSSs will be denoted by $\mathfrak{F}\mathfrak{H}(U)$. If U has only $(u, \zeta_{F_T}(u), \eta_{F_T}(u))$, then it is said to be a fermatean hesitant fuzzy number (FHFN) and is represented by $\tilde{t} = \{\zeta_{\tilde{t}}, \eta_{\tilde{t}}\}$.

$$\mathcal{F} = \{(u, h_{\mathcal{F}}(u) : u \in U)\}, \tag{2.6}$$

is called an interval-valued Fermatean hesitant (IVFHH) \mathcal{F} on U , where

$$h_{\mathcal{F}}(u) = \left\{ (\widehat{\zeta}_{\mathcal{F}}(u), \widehat{\eta}_{\mathcal{F}}(u)) : \widehat{\zeta}_{\mathcal{F}}(u) = [\zeta_{\mathcal{F}}^-, \zeta_{\mathcal{F}}^+] \in D[0, 1], \widehat{\eta}_{\mathcal{F}}(u) = [\eta_{\mathcal{F}}^-, \eta_{\mathcal{F}}^+] \in D[0, 1], (\zeta_{\mathcal{F}}^+)^3 + (\eta_{\mathcal{F}}^+)^3 \leq 1 \right\},$$

where $\widehat{\zeta}_{\mathcal{F}}(u)$ is the possible Fermatean membership interval and $\widehat{\eta}_{\mathcal{F}}(u)$ is the possible Fermatean nonmembership intervals of \mathcal{F} . Throughout this article, Υ will show the set of all IVFHHs.

Principle of recognition is defined as: In discourse universe \mathfrak{X} , let it be assumed that there are m patterns defined by FFS \mathfrak{N}_k ($k = 1, 2, \dots, m$). Again, let us suppose that there is a model to be identified with FFS \mathfrak{P} in \mathfrak{X} .

The relationship index degree between FFSs \mathfrak{N}_k and \mathfrak{P} is described as:

$$R(\mathfrak{N}_{k0}, \mathfrak{P}) = \max_{1 \leq k \leq m} \{R(\mathfrak{N}_k, \mathfrak{P})\}.$$

In this case, it is decided that sample \mathfrak{P} belongs to \mathfrak{N}_{k0} .

The set

$$IE_F(N) = \sum_{i=1}^n [\zeta_N^6(x_i) + \eta_N^6(x_i) + \theta_N^6(x_i)] \tag{2.7}$$

is called informational energies of Fermatean fuzzy set N [35].

For FFSs N and M , correlation and KK are defined as [35]:

$$C_F(N, M) = \sum_{i=1}^n [\zeta_N^3(x_i) \cdot \zeta_M^3(x_i) + \eta_N^3(x_i) \cdot \eta_M^3(x_i) + \theta_N^3(x_i) \cdot \theta_M^3(x_i)] \tag{2.8}$$

$$\mathfrak{C}_F(N, M) = \frac{C_F(N, M)}{\sqrt{IE_F(N) \cdot IE_F(M)}} \tag{2.9}$$

$$= \frac{\sum_{i=1}^n [\zeta_N^3(x_i) \cdot \zeta_M^3(x_i) + \eta_N^3(x_i) \cdot \eta_M^3(x_i) + \theta_N^3(x_i) \cdot \theta_M^3(x_i)]}{\sqrt{\sum_{i=1}^n [\zeta_N^6(x_i) + \eta_N^6(x_i) + \theta_N^6(x_i)]} \cdot \sqrt{\sum_{i=1}^n [\zeta_M^6(x_i) + \eta_M^6(x_i) + \theta_M^6(x_i)]}} \tag{2.10}$$

$$\mathfrak{D}_F(N, M) = \frac{C_I(N, M)}{\max\{IE(N), IE(M)\}} \tag{2.11}$$

$$= \frac{\sum_{i=1}^n [\zeta_N(x_i) \cdot \zeta_M(x_i) + \eta_N(x_i) \cdot \eta_M(x_i) + \theta_N^3(x_i) \cdot \theta_M^3(x_i)]}{\max[\sum_{i=1}^n [\zeta_N^6(x_i) + \eta_N^6(x_i) + \theta_N^6(x_i)], \sum_{i=1}^n [\zeta_M^6(x_i) + \eta_M^6(x_i) + \theta_M^6(x_i)]]}. \tag{2.12}$$

3. Correlation coefficients

The set

$$\mathcal{F} = \{(u, h_{\mathcal{F}}(u)) : u \in U\}, \tag{3.1}$$

is called an IVFHI \mathcal{F} on U , where

$$h_{\mathcal{F}}(u) = \left\{ (\widehat{\zeta}_{\mathcal{F}}(u), \widehat{\eta}_{\mathcal{F}}(u)) : \widehat{\zeta}_{\mathcal{F}}(u) = [\zeta_{\mathcal{F}}^-, \zeta_{\mathcal{F}}^+] \in D[0, 1], \widehat{\eta}_{\mathcal{F}}(u) = [\eta_{\mathcal{F}}^-, \eta_{\mathcal{F}}^+] \in D[0, 1], (\zeta_{\mathcal{F}}^+)^3 + (\eta_{\mathcal{F}}^+)^3 \leq 1 \right\},$$

where $\widehat{\zeta}_{\mathcal{F}}(u)$ is the possible Fermatean membership interval and $\widehat{\eta}_{\mathcal{F}}(u)$ is the possible Fermatean nonmembership intervals of \mathcal{F} . Throughout this article, Υ will show the set of all IVFHs.

The indeterminacy degree of u to \mathcal{F} is described as $\theta_{\mathcal{F}}(u) = \sqrt[3]{1 - (\zeta_{\mathcal{F}}^3(u) + \eta_{\mathcal{F}}^3(u))}$, for any FFS \mathcal{F} and $u \in U$.

Let $\mathcal{F} = \{u_i, \zeta_{\mathcal{F}}(u_i), \eta_{\mathcal{F}}(u_i) | u_i \in U\}$ be a IVFHFS, where $\zeta_{\mathcal{F}}(u_i), \eta_{\mathcal{F}}(u_i) \in [0, 1]$ and $\zeta_{\mathcal{F}}^3(u_i) + \eta_{\mathcal{F}}^3(u_i) | u_i \leq 1$ for each $u_i \in U$, we define the informational energy of the FFS \mathcal{F} as

$$IE(\mathcal{F}) = \sum_{i=1}^n \left[(\zeta_{\mathcal{F}}^-(u_i))^6 + (\zeta_{\mathcal{F}}^+(u_i))^6 + (\eta_{\mathcal{F}}^-(u_i))^6 + (\eta_{\mathcal{F}}^+(u_i))^6 + (\theta_{\mathcal{F}}^-(u_i))^6 + (\theta_{\mathcal{F}}^+(u_i))^6 \right]. \tag{3.2}$$

Suppose that two IVFHFS's $\mathcal{F} = \{u_i, \zeta_{\mathcal{F}}(u_i), \eta_{\mathcal{F}}(u_i) | u_i \in U\}$ and $\mathcal{G} = \{u_i, \zeta_{\mathcal{G}}(u_i), \eta_{\mathcal{G}}(u_i) | u_i \in U\}$, where $\zeta_{\mathcal{F}}(u_i), \eta_{\mathcal{F}}(u_i), \zeta_{\mathcal{G}}(u_i), \eta_{\mathcal{G}}(u_i) \in [0, 1]$ for each $u_i \in U$. Hence, the correlation of the FFVs \mathcal{F}, \mathcal{G} is defined:

$$C(\mathcal{F}, \mathcal{G}) = \sum_{i=1}^n \left[(\zeta_{\mathcal{F}}^-(u_i))^3 (\zeta_{\mathcal{G}}^-(u_i))^3 + (\eta_{\mathcal{F}}^-(u_i))^3 (\eta_{\mathcal{G}}^-(u_i))^3 + (\theta_{\mathcal{F}}^-(u_i))^3 (\theta_{\mathcal{G}}^-(u_i))^3 + \right. \tag{3.3}$$

$$\left. (\zeta_{\mathcal{F}}^+(u_i))^3 (\zeta_{\mathcal{G}}^+(u_i))^3 + (\eta_{\mathcal{F}}^+(u_i))^3 (\eta_{\mathcal{G}}^+(u_i))^3 + (\theta_{\mathcal{F}}^+(u_i))^3 (\theta_{\mathcal{G}}^+(u_i))^3 \right].$$

For the correlation of FFSs, the conditions

- (1) $C(\mathcal{F}, \mathcal{F}) = IE(\mathcal{F})$
- (2) $C(\mathcal{F}, \mathcal{G}) = C(\mathcal{G}, \mathcal{F})$

are hold.

Definition 3.1 Choose two FFSs \mathcal{F} and \mathcal{G} on X . Then the KK between \mathcal{F}, \mathcal{G} is defined by

$$\begin{aligned} \mathfrak{c}(\mathcal{F}, \mathcal{G}) &= \frac{C(\mathcal{F}, \mathcal{G})}{[IE(\mathcal{F}).IE(\mathcal{G})]^{1/2}} \tag{3.4} \\ &= \frac{\sum_{i=1}^n (\zeta_{\mathcal{F}}^-(u_i))^3 (\zeta_{\mathcal{G}}^-(u_i))^3 + (\eta_{\mathcal{F}}^-(u_i))^3 (\eta_{\mathcal{G}}^-(u_i))^3 + (\theta_{\mathcal{F}}^-(u_i))^3 (\theta_{\mathcal{G}}^-(u_i))^3}{\sqrt{\sum_{i=1}^n \left[(\zeta_{\mathcal{F}}^-(u_i))^6 + (\zeta_{\mathcal{F}}^+(u_i))^6 + (\eta_{\mathcal{F}}^-(u_i))^6 + (\eta_{\mathcal{F}}^+(u_i))^6 + (\theta_{\mathcal{F}}^-(u_i))^6 + (\theta_{\mathcal{F}}^+(u_i))^6 \right]}} \\ &\quad \frac{+ (\zeta_{\mathcal{F}}^+(u_i))^3 (\zeta_{\mathcal{G}}^+(u_i))^3 + (\eta_{\mathcal{F}}^+(u_i))^3 (\eta_{\mathcal{G}}^+(u_i))^3 + (\theta_{\mathcal{F}}^+(u_i))^3 (\theta_{\mathcal{G}}^+(u_i))^3}{\sqrt{\sum_{i=1}^n \left[(\zeta_{\mathcal{G}}^-(u_i))^6 + (\zeta_{\mathcal{G}}^+(u_i))^6 + (\eta_{\mathcal{G}}^-(u_i))^6 + (\eta_{\mathcal{G}}^+(u_i))^6 + (\theta_{\mathcal{G}}^-(u_i))^6 + (\theta_{\mathcal{G}}^+(u_i))^6 \right]}} \\ &\times \sqrt{\sum_{i=1}^n \left[(\zeta_{\mathcal{G}}^-(u_i))^6 + (\zeta_{\mathcal{G}}^+(u_i))^6 + (\eta_{\mathcal{G}}^-(u_i))^6 + (\eta_{\mathcal{G}}^+(u_i))^6 + (\theta_{\mathcal{G}}^-(u_i))^6 + (\theta_{\mathcal{G}}^+(u_i))^6 \right]} \end{aligned}$$

Theorem 3.2 For any two FFSs \mathcal{F}, \mathcal{G} in X , the KK of FFSs satisfies the following conditions:

$$(P1) \quad \mathfrak{C}(\mathcal{F}, \mathcal{G}) = \mathfrak{C}(\mathcal{G}, \mathcal{F}),$$

$$(P2) \quad \text{If } \mathcal{F} = \mathcal{G}, \text{ then } \mathfrak{C}(\mathcal{F}, \mathcal{G}) = 1,$$

$$(P3) \quad 0 \leq \mathfrak{C}(\mathcal{F}, \mathcal{G}) \leq 1.$$

Proof We only proved condition (P2). Obviously, $\mathfrak{C}(\mathcal{F}, \mathcal{G}) \geq 0$.

$$\begin{aligned} C(\mathcal{F}, \mathcal{G}) &= \sum_{i=1}^n \left[(\zeta_{\mathcal{F}}^-(u_i))^3 (\zeta_{\mathcal{G}}^-(u_i))^3 + (\eta_{\mathcal{F}}^-(u_i))^3 (\eta_{\mathcal{G}}^-(u_i))^3 + (\theta_{\mathcal{F}}^-(u_i))^3 (\theta_{\mathcal{G}}^-(u_i))^3 \right. \\ &\quad \left. + (\zeta_{\mathcal{F}}^+(u_i))^3 (\zeta_{\mathcal{G}}^+(u_i))^3 + (\eta_{\mathcal{F}}^+(u_i))^3 (\eta_{\mathcal{G}}^+(u_i))^3 + (\theta_{\mathcal{F}}^+(u_i))^3 (\theta_{\mathcal{G}}^+(u_i))^3 \right] \\ &= \left[(\zeta_{\mathcal{F}}^-(u_1))^3 (\zeta_{\mathcal{G}}^-(u_1))^3 + (\eta_{\mathcal{F}}^-(u_1))^3 (\eta_{\mathcal{G}}^-(u_1))^3 + (\theta_{\mathcal{F}}^-(u_1))^3 (\theta_{\mathcal{G}}^-(u_1))^3 \right. \\ &\quad \left. + (\zeta_{\mathcal{F}}^+(u_1))^3 (\zeta_{\mathcal{G}}^+(u_1))^3 + (\eta_{\mathcal{F}}^+(u_1))^3 (\eta_{\mathcal{G}}^+(u_1))^3 + (\theta_{\mathcal{F}}^+(u_1))^3 (\theta_{\mathcal{G}}^+(u_1))^3 \right] \\ &= \left[(\zeta_{\mathcal{F}}^-(u_2))^3 (\zeta_{\mathcal{G}}^-(u_2))^3 + (\eta_{\mathcal{F}}^-(u_2))^3 (\eta_{\mathcal{G}}^-(u_2))^3 + (\theta_{\mathcal{F}}^-(u_2))^3 (\theta_{\mathcal{G}}^-(u_2))^3 \right. \\ &\quad \left. + (\zeta_{\mathcal{F}}^+(u_2))^3 (\zeta_{\mathcal{G}}^+(u_2))^3 + (\eta_{\mathcal{F}}^+(u_2))^3 (\eta_{\mathcal{G}}^+(u_2))^3 + (\theta_{\mathcal{F}}^+(u_2))^3 (\theta_{\mathcal{G}}^+(u_2))^3 \right] \\ &\quad + \cdots + \left[(\zeta_{\mathcal{F}}^-(u_n))^3 (\zeta_{\mathcal{G}}^-(u_n))^3 + (\eta_{\mathcal{F}}^-(u_n))^3 (\eta_{\mathcal{G}}^-(u_n))^3 + (\theta_{\mathcal{F}}^-(u_n))^3 (\theta_{\mathcal{G}}^-(u_n))^3 \right. \\ &\quad \left. + (\zeta_{\mathcal{F}}^+(u_n))^3 (\zeta_{\mathcal{G}}^+(u_n))^3 + (\eta_{\mathcal{F}}^+(u_n))^3 (\eta_{\mathcal{G}}^+(u_n))^3 + (\theta_{\mathcal{F}}^+(u_n))^3 (\theta_{\mathcal{G}}^+(u_n))^3 \right]. \end{aligned}$$

Using the Cauchy–Schwarz inequality, for $(\zeta_1 + \cdots + \zeta_n) \in R^n$ and $(\eta_1 + \cdots + \eta_n) \in R^n$,

$$(\zeta_1 \eta_1 + \zeta_2 \eta_2 + \cdots + \zeta_n \eta_n)^2 \leq (\zeta_1^2 + \cdots + \zeta_n^2) \cdot (\eta_1^2 + \cdots + \eta_n^2).$$

Then,

$$\begin{aligned}
 [C(\mathcal{F}, \mathcal{G})]^2 &\leq \left[((\zeta_{\mathcal{F}}^-(u_1))^6 + (\eta_{\mathcal{F}}^-(u_1))^6 + (\theta_{\mathcal{F}}^-(u_1))^6) + ((\zeta_{\mathcal{F}}^+(u_1))^6 + (\eta_{\mathcal{F}}^+(u_1))^6 + (\theta_{\mathcal{F}}^+(u_1))^6) \right. \\
 &+ ((\zeta_{\mathcal{F}}^-(u_2))^6 + (\eta_{\mathcal{F}}^-(u_2))^6 + (\theta_{\mathcal{F}}^-(u_2))^6) + ((\zeta_{\mathcal{F}}^+(u_2))^6 + (\eta_{\mathcal{F}}^+(u_2))^6 + (\theta_{\mathcal{F}}^+(u_2))^6) + \dots \\
 &+ \left. ((\zeta_{\mathcal{F}}^-(u_n))^6 + (\eta_{\mathcal{F}}^-(u_n))^6 + (\theta_{\mathcal{F}}^-(u_n))^6) + ((\zeta_{\mathcal{F}}^+(u_n))^6 + (\eta_{\mathcal{F}}^+(u_n))^6 + (\theta_{\mathcal{F}}^+(u_n))^6) \right] \\
 &\times \left[((\zeta_{\mathcal{G}}^-(u_1))^6 + (\eta_{\mathcal{G}}^-(u_1))^6 + (\theta_{\mathcal{G}}^-(u_1))^6) + ((\zeta_{\mathcal{G}}^+(u_1))^6 + (\eta_{\mathcal{G}}^+(u_1))^6 + (\theta_{\mathcal{G}}^+(u_1))^6) \right. \\
 &+ ((\zeta_{\mathcal{G}}^-(u_2))^6 + (\eta_{\mathcal{G}}^-(u_2))^6 + (\theta_{\mathcal{G}}^-(u_2))^6) + ((\zeta_{\mathcal{G}}^+(u_2))^6 + (\eta_{\mathcal{G}}^+(u_2))^6 + (\theta_{\mathcal{G}}^+(u_2))^6) + \dots \\
 &+ \left. ((\zeta_{\mathcal{G}}^-(u_n))^6 + (\eta_{\mathcal{G}}^-(u_n))^6 + (\theta_{\mathcal{G}}^-(u_n))^6) + ((\zeta_{\mathcal{G}}^+(u_n))^6 + (\eta_{\mathcal{G}}^+(u_n))^6 + (\theta_{\mathcal{G}}^+(u_n))^6) \right] \\
 &= \sum_{i=1}^n [((\zeta_{\mathcal{F}}^-(u_i))^6 + (\eta_{\mathcal{F}}^-(u_i))^6 + (\theta_{\mathcal{F}}^-(u_i))^6) + ((\zeta_{\mathcal{F}}^+(u_i))^6 + (\eta_{\mathcal{F}}^+(u_i))^6 + (\theta_{\mathcal{F}}^+(u_i))^6)] \\
 &\times \sum_{i=1}^n [((\zeta_{\mathcal{G}}^-(u_i))^6 + (\eta_{\mathcal{G}}^-(u_i))^6 + (\theta_{\mathcal{G}}^-(u_i))^6) + ((\zeta_{\mathcal{G}}^+(u_i))^6 + (\eta_{\mathcal{G}}^+(u_i))^6 + (\theta_{\mathcal{G}}^+(u_i))^6)] \\
 &= IE(\mathcal{F}).IE(\mathcal{G}).
 \end{aligned}$$

Therefore, $[C(\mathcal{F}, \mathcal{G})]^2 \leq IE(\mathcal{F}).IE(\mathcal{G})$ and $\mathfrak{C}(\mathcal{F}, \mathcal{G}) \leq 1$. □

Example 3.3 Take three FFSs $\mathcal{F} = \{([0.6, 0.8], [0.5, 0.7])\}$, $\mathcal{G} = \{([0.3, 0.7], [0.5, 0.6]), ([0.4, 0.7], [0.5, 0.7])\}$ in \mathfrak{X} . By Equation 3.2, the informational energies of \mathcal{F} , \mathcal{G} :

$$IE(\mathcal{F}) = (0.6^6 + 0.8^6 + 0.87^6) + (0.5^6 + 0.7^6 + 0.53^6) = 0.9$$

$$IE(\mathcal{G}) = (0.3^6 + 0.7^6 + 0.95^6) + (0.5^6 + 0.6^6 + 0.76^6) + (0.4^6 + 0.7^6 + 0.93^6) + (0.5^6 + 0.7^6 + 0.68^6) = 2.74$$

By using Equation 3.4, the correlation between the FFSs \mathcal{F}, \mathcal{G} is written as:

$$\begin{aligned}
 C(\mathcal{F}, \mathcal{G}) &= \left[(\zeta_{\mathcal{F}}^-)^3(\zeta_{\mathcal{G}_1}^-)^3 + (\eta_{\mathcal{F}}^-)^3(\eta_{\mathcal{G}_1}^-)^3 + (\theta_{\mathcal{F}}^-)^3(\theta_{\mathcal{G}_1}^-)^3 + (\zeta_{\mathcal{F}}^+)^3(\zeta_{\mathcal{G}_1}^+)^3 + (\eta_{\mathcal{F}}^+)^3(\eta_{\mathcal{G}_1}^+)^3 + (\theta_{\mathcal{F}}^+)^3(\theta_{\mathcal{G}_1}^+)^3 + \right. \\
 &(\zeta_{\mathcal{F}}^-)^3(\zeta_{\mathcal{G}_2}^-)^3 + (\eta_{\mathcal{F}}^-)^3(\eta_{\mathcal{G}_2}^-)^3 + (\theta_{\mathcal{F}}^-)^3(\theta_{\mathcal{G}_2}^-)^3 + (\zeta_{\mathcal{F}}^+)^3(\zeta_{\mathcal{G}_2}^+)^3 + (\eta_{\mathcal{F}}^+)^3(\eta_{\mathcal{G}_2}^+)^3 + (\theta_{\mathcal{F}}^+)^3(\theta_{\mathcal{G}_2}^+)^3 \left. \right] \\
 &= 0.6^3 0.3^3 + 0.8^3 0.7^3 + 0.87^3 0.95^3 + 0.5^3 0.5^3 + 0.7^3 0.6^3 + 0.53^3 0.76^3 \\
 &+ 0.6^3 0.4^3 + 0.8^3 0.7^3 + 0.87^3 0.93^3 + 0.5^3 0.5^3 + 0.7^3 0.7^3 + 0.53^3 0.68^3 = 1.81
 \end{aligned}$$

Hence, the KK between the FFSs \mathcal{F}, \mathcal{G} is given by:

$$\mathfrak{C}(\mathcal{F}, \mathcal{G}) = \frac{C(\mathcal{F}, \mathcal{G})}{[IE(\mathcal{F}).IE(\mathcal{G})]^{1/2}} = \frac{1.81}{[(0.9).(2.74)]^{1/2}} = 0.734$$

Definition 3.4 For \mathcal{F} and \mathcal{G} , the definition of KK as:

$$\begin{aligned} \mathfrak{D}(\mathcal{F}, \mathcal{G}) &= \frac{C(\mathcal{F}, \mathcal{G})}{\max [IE(\mathcal{F}).IE(\mathcal{G})]} \tag{3.5} \\ &= \frac{\sum_{i=1}^n (\zeta_{\mathcal{F}}^-(u_i))^3 (\zeta_{\mathcal{G}}^-(u_i))^3 + (\eta_{\mathcal{F}}^-(u_i))^3 (\eta_{\mathcal{G}}^-(u_i))^3 + (\theta_{\mathcal{F}}^-(u_i))^3 (\theta_{\mathcal{G}}^-(u_i))^3}{\max \left\{ \sum_{i=1}^n \left[(\zeta_{\mathcal{F}}^-(u_i))^6 + (\zeta_{\mathcal{F}}^+(u_i))^6 + (\eta_{\mathcal{F}}^-(u_i))^6 + (\zeta_{\mathcal{F}}^+(u_i))^6 + (\theta_{\mathcal{F}}^-(u_i))^6 + (\theta_{\mathcal{F}}^+(u_i))^6 \right], \right. \\ &\quad \left. \frac{\sum_{i=1}^n \left[(\zeta_{\mathcal{F}}^+(u_i))^3 (\zeta_{\mathcal{G}}^+(u_i))^3 + (\eta_{\mathcal{F}}^+(u_i))^3 (\eta_{\mathcal{G}}^+(u_i))^3 + (\theta_{\mathcal{F}}^+(u_i))^3 (\theta_{\mathcal{G}}^+(u_i))^3 \right]}{\sum_{i=1}^n \left[(\zeta_{\mathcal{G}}^-(u_i))^6 + (\zeta_{\mathcal{G}}^+(u_i))^6 + (\eta_{\mathcal{G}}^-(u_i))^6 + (\zeta_{\mathcal{G}}^+(u_i))^6 + (\theta_{\mathcal{G}}^-(u_i))^6 + (\theta_{\mathcal{G}}^+(u_i))^6 \right]} \right\}} \end{aligned}$$

Theorem 3.5 For any two FFSs \mathcal{F}, \mathcal{G} $\mathfrak{D}(\mathcal{F}, \mathcal{G})$ satisfies the following conditions:

(P1) $\mathfrak{D}(\mathcal{F}, \mathcal{G}) = \mathfrak{D}(\mathcal{G}, \mathcal{F})$,

(P2) $\mathcal{F} = \mathcal{G}$ iff $\mathfrak{D}(\mathcal{F}, \mathcal{G}) = 1$,

(P3) $0 \leq \mathfrak{D}(\mathcal{F}, \mathcal{G}) \leq 1$.

Proof We only proved condition (P2). It is clear that $\mathfrak{D}(\mathcal{F}, \mathcal{G}) \geq 0$. Since from Theorem 3.2, $\left[C(\mathcal{F}, \mathcal{G}) \right]^2 \leq IE(\mathcal{F}).IE(\mathcal{G})$. Therefore, $C(\mathcal{F}, \mathcal{G}) \leq \max [IE(\mathcal{F}).IE(\mathcal{G})]$; thus, $\mathfrak{D}(\mathcal{F}, \mathcal{G}) \leq 1$. \square

It is possible to define these KK s in a different way. Weights will be used for these new definitions because, in numerous real-life practices, the distinct sets can have diverse weights. Therefore, weight ρ_i of every element $x_i \in \mathfrak{X}$ must be considered in new definitions. The KK s to be defined by weights will be called weighted KK s. For these definitions, choose the weight vector as ρ with the condition $\sum_{i=1}^n \rho_i = 1$ for $\rho_i \geq 1$. Therefore, $C_{\rho}(\mathcal{F}, \mathcal{G})$, $\mathfrak{C}_{\rho}(\mathcal{F}, \mathcal{G})$ are defined as follows:

$$\begin{aligned} \mathfrak{C}_{\rho}(\mathcal{F}, \mathcal{G}) &= \frac{C_{\rho}(\mathcal{F}, \mathcal{G})}{[IE_{\rho}(\mathcal{F}).IE_{\rho}(\mathcal{G})]^{1/2}} \tag{3.6} \\ &= \frac{\sum_{i=1}^n \rho_i (\zeta_{\mathcal{F}}^-(u_i))^3 (\zeta_{\mathcal{G}}^-(u_i))^3 + (\eta_{\mathcal{F}}^-(u_i))^3 (\eta_{\mathcal{G}}^-(u_i))^3 + (\theta_{\mathcal{F}}^-(u_i))^3 (\theta_{\mathcal{G}}^-(u_i))^3}{\sqrt{\sum_{i=1}^n \rho_i \left[(\zeta_{\mathcal{F}}^-(u_i))^6 + (\zeta_{\mathcal{F}}^+(u_i))^6 + (\eta_{\mathcal{F}}^-(u_i))^6 + (\zeta_{\mathcal{F}}^+(u_i))^6 + (\theta_{\mathcal{F}}^-(u_i))^6 + (\theta_{\mathcal{F}}^+(u_i))^6 \right]}} \\ &\quad \frac{\sum_{i=1}^n \left[(\zeta_{\mathcal{F}}^+(u_i))^3 (\zeta_{\mathcal{G}}^+(u_i))^3 + (\eta_{\mathcal{F}}^+(u_i))^3 (\eta_{\mathcal{G}}^+(u_i))^3 + (\theta_{\mathcal{F}}^+(u_i))^3 (\theta_{\mathcal{G}}^+(u_i))^3 \right]}{\sqrt{\sum_{i=1}^n \left[(\zeta_{\mathcal{G}}^-(u_i))^6 + (\zeta_{\mathcal{G}}^+(u_i))^6 + (\eta_{\mathcal{G}}^-(u_i))^6 + (\zeta_{\mathcal{G}}^+(u_i))^6 + (\theta_{\mathcal{G}}^-(u_i))^6 + (\theta_{\mathcal{G}}^+(u_i))^6 \right]}} \end{aligned}$$

$$\begin{aligned}
\mathfrak{D}_\rho(\mathcal{F}, \mathcal{G}) &= \frac{C_\rho(\mathcal{F}, \mathcal{G})}{\max[IE_\rho(\mathcal{F}), IE_\rho(\mathcal{G})]} \\
&= \frac{\sum_{i=1}^n \rho_i (\zeta_{\mathcal{F}}^-(u_i))^3 (\zeta_{\mathcal{G}}^-(u_i))^3 + (\eta_{\mathcal{F}}^-(u_i))^3 (\eta_{\mathcal{G}}^-(u_i))^3 + (\theta_{\mathcal{F}}^-(u_i))^3 (\theta_{\mathcal{G}}^-(u_i))^3}{\max \sum_{i=1}^n \rho_i \left[(\zeta_{\mathcal{F}}^-(u_i))^6 + (\zeta_{\mathcal{F}}^+(u_i))^6 + (\eta_{\mathcal{F}}^-(u_i))^6 + (\eta_{\mathcal{F}}^+(u_i))^6 + (\theta_{\mathcal{F}}^-(u_i))^6 + (\theta_{\mathcal{F}}^+(u_i))^6 \right.} \\
&\quad \left. + (\zeta_{\mathcal{F}}^+(u_i))^3 (\zeta_{\mathcal{G}}^+(u_i))^3 + (\eta_{\mathcal{F}}^+(u_i))^3 (\eta_{\mathcal{G}}^+(u_i))^3 + (\theta_{\mathcal{F}}^+(u_i))^3 (\theta_{\mathcal{G}}^+(u_i))^3 \right]}{\sum_{i=1}^n \rho_i \left[(\zeta_{\mathcal{G}}^-(u_i))^6 + (\zeta_{\mathcal{G}}^+(u_i))^6 + (\eta_{\mathcal{G}}^-(u_i))^6 + (\eta_{\mathcal{G}}^+(u_i))^6 + (\theta_{\mathcal{G}}^-(u_i))^6 + (\theta_{\mathcal{G}}^+(u_i))^6 \right]}
\end{aligned} \tag{3.7}$$

The following theorems are proved as similar to Theorems 3.2 and 3.5:

Theorem 3.6 Take a weight vector of x_i as ρ and it is considered to satisfy the $\sum_{i=1}^n \rho_i = 1$ condition for $\rho_i \geq 1$. The weighted KK between the FFSs $\mathfrak{R}, \mathfrak{M}$ defined by Equation 3.6 satisfies:

- (P1) $\mathfrak{C}_\rho(\mathcal{F}, \mathcal{G}) = \mathfrak{C}_\rho(\mathcal{F}, \mathcal{G})$,
- (P2) $0 \leq \mathfrak{C}_\rho(\mathcal{F}, \mathcal{G}) \leq 1$;
- (P3) $\mathfrak{C}_\rho(\mathcal{F}, \mathcal{G}) = 1$ iff $\mathcal{F} = \mathcal{G}$

Theorem 3.7 The weighted KK between the FFSs \mathcal{F}, \mathcal{G} defined by Equation 3.7 satisfies:

- (P1) $\mathfrak{D}_\rho(\mathcal{F}, \mathcal{G}) = \mathfrak{D}_\rho(\mathcal{F}, \mathcal{G})$,
- (P2) $0 \leq \mathfrak{D}_\rho(\mathcal{F}, \mathcal{G}) \leq 1$;
- (P3) $\mathfrak{D}_\rho(\mathcal{F}, \mathcal{G}) = 1$ iff $\mathcal{F} = \mathcal{G}$

4. Numerical examples

4.1. Pattern recognition application

In this subsection, an example for the multicriteria decision-making problem of alternatives, from the field of pattern recognition is used as the demonstration of the application of the proposed decision-making method, as well as the effectiveness of the proposed method.

Choose three known patterns PR_1, PR_2, PR_3 . For the finite universe $U = \{u_1, u_2, u_3\}$ as

$$\begin{aligned}
PR_1 &= \{(u_1, [1.0, 0.0], [0.9, 0.2]), (u_2, [0.8, 0.0], [0.7, 0.1]), (u_3, [0.7, 0.4], [0.6, 0.5])\} \\
PR_2 &= \{(u_1, [0.8, 0.1], [0.9, 0.1]), (u_2, [1.0, 0.0], [0.9, 0.1]), (u_3, [0.8, 0.3], [0.7, 0.4])\} \\
PR_3 &= \{(u_1, [0.6, 0.2], [0.7, 0.3]), (u_2, [0.8, 0.2], [0.7, 0.2]), (u_3, [1.0, 0.0], [0.9, 0.1])\}.
\end{aligned}$$

Consider an unknown pattern $UP \in FFS(U)$ that will be recognized, where

$$UP = \{(u_1, [0.5, 0.3], [0.4, 0.3]), (u_2, [0.6, 0.2], [0.6, 0.3]), (u_3, [0.8, 0.1], [0.8, 0.2])\}.$$

The target of this problem is to classify the pattern UP in one of the classes PR_1, PR_2, PR_3 . For it, proposed correlation coefficient index, $\mathfrak{C}, \mathfrak{D}$, have been computed from UP to PR_i ($i = 1, 2, 3$) and are given as follows:

$$\begin{aligned}
\mathfrak{C}(PR_1, UP) &= 0.098, & \mathfrak{C}(PR_2, UP) &= 0.084, & \mathfrak{C}(PR_3, UP) &= 0.1167, \\
\mathfrak{D}(PR_1, UP) &= 0.2652, & \mathfrak{D}(PR_2, UP) &= 0.1195, & \mathfrak{D}(PR_3, UP) &= 0.4582.
\end{aligned}$$

Thus, from these two proposed correlation coefficient indexes, we conclude that the pattern UP belongs to the pattern PR_3 .

On the other hand, if we assume that weights of u_1, u_2, u_3 are 0.5, 0.3, 0.2, respectively, then we utilize the correlation coefficient $\mathfrak{C}_\rho, \mathfrak{D}_\rho$ for obtaining the most suitable pattern as:

$$\begin{aligned}\mathfrak{C}_\rho(PR_1, UP) &= 0.175, & \mathfrak{C}_\rho(PR_2, UP) &= 0.162, & \mathfrak{C}_\rho(PR_3, UP) &= 0.29, \\ \mathfrak{D}_\rho(PR_1, UP) &= 0.2, & \mathfrak{D}_\rho(PR_2, UP) &= 0.056, & \mathfrak{D}_\rho(PR_3, UP) &= 0, 327.\end{aligned}$$

Thus, ranking order of the three patterns PR_1, PR_2, PR_3 and hence PR_3 is the most desirable pattern to be classified with UP .

4.2. Multicriteria decision-making

This subsection will apply the correlation coefficients for interval-valued Pythagorean hesitant fuzzy sets (IVPHFSs) to MCDM problems and clustering analysis.

Let $L = \{L_i : i = 1, 2, \dots, m\}$ be a finite set of alternatives and $K = \{K_j : j = 1, 2, \dots, n\}$ be a set of criteria, ω be a weight vector of the criteria ($\omega_j \in [0, 1], \sum_{j=1}^n \omega_j = 1$). Suppose $M = (h_{\mathcal{F}_j})_{m \times n}$ is an interval-valued Fermatean hesitant fuzzy decision matrix (IVFHFDM) where $h_{\mathcal{F}_j} = \{([\zeta_{ij}^-, \zeta_{ij}^+], [\eta_{ij}^-, \eta_{ij}^+]) : (\zeta_{ij}^+)^3 + (\eta_{ij}^+)^3 \leq 1\}$ ($i = 1, 2, \dots, m; j = 1, 2, \dots, n$) is an INFHFE given by decision makers to evaluate the alternative L_i with respect to the criteria K_j .

The concrete algorithm is listed as follows:

Step 1: Input $M = (h_{\mathcal{F}_j})_{m \times n}$, $\omega = (\omega_1, \dots, \omega_n)^T$ and $\lambda \in [0, 1]$.

Step 2: Make M the revised IVFHFDM $M' = (h'_{\mathcal{F}_j})_{m \times n}$.

Step 3: Compute the weighted correlation coefficients between each \mathcal{F}_i and \mathcal{F}_i^* .

Step 4: Get the priority of the alternatives L_i by ranking the above correlation coefficients.

Example 4.1 A scientific committee should evaluate four possible R&D projects L_i ($i = 1, 2, 3, 4$) according to the three criteria K_j ($j = 1, 2, 3$). Here C_1, C_2 are both benefit criteria and C_3 is a cost criterion. Take the weight vector of criteria as $\omega = (0.3, 0.5, 0.3)^T$. The decision matrix is given by the experts as Table 1.

Considering that $|h_{\mathcal{F}_\infty}| \neq |h_{\mathcal{F}_\infty}|$ for any $j \in \{1, 2, 3\}$, we revised the matrix M and M' in Table 2. Furthermore, compute the four correlation coefficients between these alternatives and the ideal alternatives in Table 3. The result shows the R&D project L_3 are always the most optimal, although the four ranking results (Table 4) are little different.

4.3. Medical decision-making

Example 4.2 A clinic wants to buy a panoramic X-ray machine. In the research conducted for this purchase process, a selection will be made from four different brands ($\widehat{\mathcal{F}} = \{\widehat{\mathcal{F}}_1, \widehat{\mathcal{F}}_2, \widehat{\mathcal{F}}_3, \widehat{\mathcal{F}}_4\}$). When making this choice, three different criteria are considered: C_1 : Robustness and permanence, C_2 : service support, C_3 : price

Table 1. IVFHFDM M .

	C_1	C_2	C_3
\mathcal{F}_1	$\{([0.6, 0.8], [0.4, 0.5]),$ $([0.5, 0.8], [0.1, 0.3]),$	$\{([0.5, 0.7], [0.1, 0.3]),$ $([0.4, 0.6], [0.2, 0.5])\}$	$\{([0.5, 0.7], [0.3, 0.4])\}$
\mathcal{F}_2	$\{([0.4, 0.7], [0.2, 0.3])\}$	$\{([0.5, 0.6], [0.3, 0.4])\}$ $\{([0.3, 0.5], [0.2, 0.3])\}$ $\{([0.2, 0.7], [0.2, 0.4])\}$ $\{([0.3, 0.5], [0.2, 0.7])\}$	$\{([0.3, 0.7], [0.2, 0.4]),$ $([0.2, 0.5], [0.3, 0.4])\}$ $([0.2, 0.6], [0.2, 0.3])\}$
\mathcal{F}_3	$\{([0.3, 0.5], [0.1, 0.2])\}$ $\{([0.5, 0.7], [0.3, 0.4])\}$	$\{([0.6, 0.8], [0.2, 0.3]),$ $([0.6, 0.9], [0.1, 0.2])\}$	$\{([0.3, 0.4], [0.5, 0.6]),$ $\{([0.3, 0.4], [0.6, 0.7])\}$ $\{([0.4, 0.5], [0.4, 0.6])\}$
\mathcal{F}_4	$\{([0.5, 0.6], [0.2, 0.4])\}$	$\{([0.4, 0.5], [0.3, 0.4]),$ $([0.3, 0.5], [0.3, 0.4])\}$	$\{([0.5, 0.8], [0.3, 0.4]),$

Table 2. Revised IVFHFDM M' .

	C_1	C_2	C_3
\mathcal{F}_1	$\{([0.5, 0.8], [0.1, 0.3]),$ $([0.6, 0.8], [0.4, 0.5]),$	$\{([0.5, 0.7], [0.1, 0.3]),$ $([0.4, 0.6], [0.2, 0.5])\}$ $([0.4, 0.6], [0.2, 0.5])\}$ $([0.4, 0.6], [0.2, 0.5])\}$	$\{([0.5, 0.7], [0.3, 0.4])\}$ $\{([0.5, 0.7], [0.3, 0.4])\}$ $\{([0.5, 0.7], [0.3, 0.4])\}$
\mathcal{F}_2	$\{([0.4, 0.7], [0.2, 0.3])\}$ $\{([0.4, 0.7], [0.2, 0.3])\}$	$\{([0.5, 0.6], [0.3, 0.4])\}$ $\{([0.2, 0.7], [0.2, 0.4])\}$ $\{([0.3, 0.5], [0.2, 0.3])\}$ $\{([0.3, 0.5], [0.2, 0.7])\}$	$\{([0.3, 0.7], [0.2, 0.4]),$ $\{([0.2, 0.6], [0.2, 0.3])\}$ $\{([0.2, 0.5], [0.3, 0.4])\}$
\mathcal{F}_3	$\{([0.5, 0.7], [0.3, 0.4])\}$ $\{([0.3, 0.5], [0.1, 0.2])\}$	$\{([0.6, 0.8], [0.2, 0.3]),$ $\{([0.6, 0.8], [0.2, 0.3]),$ $\{([0.6, 0.9], [0.1, 0.2])\}$ $\{([0.6, 0.9], [0.1, 0.2])\}$	$\{([0.4, 0.5], [0.4, 0.6]),$ $\{([0.3, 0.4], [0.5, 0.6])\}$ $([0.3, 0.4], [0.6, 0.7])\}$
\mathcal{F}_4	$\{([0.5, 0.6], [0.2, 0.4])\}$ $\{([0.5, 0.6], [0.2, 0.4])\}$	$\{([0.4, 0.5], [0.3, 0.4]),$ $([0.4, 0.5], [0.3, 0.4])\}$ $([0.3, 0.5], [0.3, 0.4])\}$ $([0.3, 0.5], [0.3, 0.4])\}$	$\{([0.5, 0.8], [0.3, 0.4]),$ $\{([0.5, 0.8], [0.3, 0.4])\}$ $\{([0.5, 0.8], [0.3, 0.4])\}$

Table 3. Correlations Coefficients of IVFHFSSs.

	$(\mathcal{F}_1, \mathcal{F}^*)$	$(\mathcal{F}_2, \mathcal{F}^*)$	$(\mathcal{F}_3, \mathcal{F}^*)$	$(\mathcal{F}_4, \mathcal{F}^*)$
\mathfrak{C}	0.3144	0.207	0.4737	0.2116
\mathfrak{D}	0.2225	0.1864	0.3974	0.1721
\mathfrak{C}_ρ	0.5937	0.4851	0.6803	0.498
\mathfrak{D}_ρ	0.4765	0.4469	0.5154	0.4319

Table 4. Ranking.

$\mathfrak{C}(\mathcal{F}_3, \mathcal{F}^*) > \mathfrak{C}(\mathcal{F}_1, \mathcal{F}^*) > \mathfrak{C}(\mathcal{F}_4, \mathcal{F}^*) > \mathfrak{C}(\mathcal{F}_2, \mathcal{F}^*)$
$\mathfrak{D}(\mathcal{F}_3, \mathcal{F}^*) > \mathfrak{D}(\mathcal{F}_1, \mathcal{F}^*) > \mathfrak{D}(\mathcal{F}_2, \mathcal{F}^*) > \mathfrak{D}(\mathcal{F}_4, \mathcal{F}^*)$
$\mathfrak{C}_\rho(\mathcal{F}_3, \mathcal{F}^*) > \mathfrak{C}_\rho(\mathcal{F}_1, \mathcal{F}^*) > \mathfrak{C}_\rho(\mathcal{F}_4, \mathcal{F}^*) > \mathfrak{C}_\rho(\mathcal{F}_2, \mathcal{F}^*)$
$\mathfrak{D}_\rho(\mathcal{F}_3, \mathcal{F}^*) > \mathfrak{D}_\rho(\mathcal{F}_1, \mathcal{F}^*) > \mathfrak{D}_\rho(\mathcal{F}_2, \mathcal{F}^*) > \mathfrak{D}_\rho(\mathcal{F}_4, \mathcal{F}^*)$

performance. C_1 and C_2 are benefit criteria, C_3 are cost criteria. Let the weight vector of criteria be $\omega = (0.28, 0.47, 0.25)^T$.

Table 5. IVFHF M .

	C_1	C_2	C_3
$\widehat{\mathcal{F}}_1$	$\{\langle [0.58, 0.68], [0.41, 0.52] \rangle, \langle [0.57, 0.76], [0.18, 0.33] \rangle\}$	$\{\langle [0.46, 0.59], [0.15, 0.35] \rangle, \langle [0.37, 0.58], [0.23, 0.54] \rangle\}$	$\{\langle [0.56, 0.72], [0.18, 0.37] \rangle\}$
$\widehat{\mathcal{F}}_2$	$\{\langle [0.46, 0.67], [0.27, 0.41] \rangle\}$	$\{\langle [0.54, 0.62], [0.35, 0.42] \rangle, \langle [0.38, 0.51], [0.20, 0.33] \rangle, \langle [0.25, 0.65], [0.24, 0.40] \rangle, \langle [0.35, 0.48], [0.21, 0.68] \rangle\}$	$\{\langle [0.39, 0.51], [0.22, 0.35] \rangle, \langle [0.21, 0.53], [0.37, 0.42] \rangle, \langle [0.25, 0.58], [0.14, 0.36] \rangle\}$
$\widehat{\mathcal{F}}_3$	$\{\langle [0.33, 0.45], [0.16, 0.22] \rangle, \langle [0.58, 0.68], [0.26, 0.37] \rangle\}$	$\{\langle [0.60, 0.80], [0.14, 0.22] \rangle, \langle [0.66, 0.72], [0.24, 0.29] \rangle\}$	$\{\langle [0.33, 0.44], [0.37, 0.59] \rangle, \langle [0.38, 0.44], [0.55, 0.75] \rangle, \langle [0.43, 0.58], [0.34, 0.67] \rangle\}$
$\widehat{\mathcal{F}}_4$	$\{\langle [0.55, 0.65], [0.22, 0.44] \rangle\}$	$\{\langle [0.46, 0.51], [0.37, 0.42] \rangle, \langle [0.28, 0.48], [0.31, 0.42] \rangle\}$	$\{\langle [0.48, 0.77], [0.23, 0.37] \rangle\}$

IVFHF values are given in Table 5. Table 6 shows the revised IVFHF decision matrix, since $|h_{\widehat{\mathcal{F}}_{1j}}| \neq |h_{\widehat{\mathcal{F}}_{2j}}|$ ($j = 1, 2, 3$). Table 7 gives the correlations and informational energies of the four IVFHFSSs. The values in Table 8 are the calculated four KKs between these alternatives and the ideal alternatives.

As can be seen from these results, there is not much difference between the ranking results, and it is seen that the $\widehat{\mathcal{F}}_3$ panoramic X-ray machine is suitable for all criteria.

5. Discussion

5.1. Comparison

The KK formulas given below were defined by Chen et al. [8] and Garg [78], respectively. For the values given below, we will calculate the correlation coefficients using the formulas 5.1 and 5.2.

In order to comprehend the examples, it is helpful to keep in mind the following details: The correlation coefficient is a measure of how strongly and in what direction the independent variables are related. The range of values for this coefficient is -1 to $+1$. A direct linear relationship is shown by a positive number, and an inverse linear relationship is indicated by a negative value. If the correlation coefficient is 0, the aforementioned variables do not have a linear connection. The closer the coefficient is to $+1$ or -1 , the stronger the linearity of the relationship.

Table 6. Revised IVFHF M' .

	C_1	C_2	C_3
$\widehat{\mathcal{F}}_1$	$\{\langle [0.57, 0.76], [0.18, 0.33] \rangle, \langle [0.58, 0.68], [0.41, 0.52] \rangle\}$	$\{\langle [0.46, 0.59], [0.15, 0.35] \rangle, \langle [0.46, 0.59], [0.15, 0.35] \rangle\}$ $\{\langle [0.37, 0.58], [0.23, 0.54] \rangle, \langle [0.37, 0.58], [0.23, 0.54] \rangle\}$	$\{\langle [0.56, 0.72], [0.18, 0.37] \rangle, \langle [0.56, 0.72], [0.18, 0.37] \rangle\}$ $\{\langle [0.56, 0.72], [0.18, 0.37] \rangle\}$
$\widehat{\mathcal{F}}_2$	$\{\langle [0.46, 0.67], [0.27, 0.41] \rangle, \langle [0.46, 0.67], [0.27, 0.41] \rangle\}$	$\{\langle [0.54, 0.62], [0.35, 0.42] \rangle, \langle [0.38, 0.51], [0.20, 0.33] \rangle\}$ $\langle [0.2, 0.7], [0.2, 0.4] \rangle,$ $\langle [0.35, 0.48], [0.21, 0.68] \rangle\}$	$\{\langle [0.39, 0.51], [0.22, 0.35] \rangle, \langle [0.25, 0.58], [0.14, 0.36] \rangle\}$ $\langle [0.21, 0.53], [0.37, 0.42] \rangle\}$
$\widehat{\mathcal{F}}_3$	$\{\langle [0.58, 0.68], [0.26, 0.37] \rangle, \langle [0.33, 0.45], [0.16, 0.22] \rangle\}$	$\{\langle [0.60, 0.80], [0.14, 0.22] \rangle, \langle [0.60, 0.80], [0.14, 0.22] \rangle\}$ $\langle [0.66, 0.72], [0.24, 0.29] \rangle,$ $\langle [0.38, 0.44], [0.55, 0.75] \rangle\}$	$\{\langle [0.43, 0.58], [0.34, 0.67] \rangle, \langle [0.3, 0.4], [0.4, 0.6] \rangle\}$ $\langle [0.38, 0.44], [0.55, 0.75] \rangle\}$
$\widehat{\mathcal{F}}_4$	$\{\langle [0.55, 0.65], [0.22, 0.44] \rangle, \langle [0.55, 0.65], [0.22, 0.44] \rangle\}$	$\{\langle [0.46, 0.51], [0.37, 0.42] \rangle, \langle [0.46, 0.51], [0.37, 0.42] \rangle\}$ $\{\langle [0.28, 0.48], [0.31, 0.42] \rangle, \langle [0.28, 0.48], [0.31, 0.42] \rangle\}$	$\{\langle [0.48, 0.77], [0.23, 0.37] \rangle, \langle [0.48, 0.77], [0.23, 0.37] \rangle\}$ $\{\langle [0.48, 0.77], [0.23, 0.37] \rangle\}$

Table 7. Correlations of M' .

	C	D	C_ρ	D_ρ
$\widehat{\mathcal{F}}_1$	0.108	0.2623	0.3941	0.2616
$\widehat{\mathcal{F}}_2$	0.0963	0.2037	0.3628	0.1962
$\widehat{\mathcal{F}}_3$	0.1485	0.2809	0.4039	0.2508
$\widehat{\mathcal{F}}_4$	0.0842	0.1875	0.3612	0.2084

Table 8. Correlation coefficients of IVFHFSSs.

	$(\widehat{\mathcal{F}}_1, \widehat{\mathcal{F}}^*)$	$(\widehat{\mathcal{F}}_2, \widehat{\mathcal{F}}^*)$	$(\widehat{\mathcal{F}}_3, \widehat{\mathcal{F}}^*)$	$(\widehat{\mathcal{F}}_4, \widehat{\mathcal{F}}^*)$
\mathfrak{C}	0.4033	0.2816	0.5413	0.2896
\mathfrak{D}_ρ	0.2861	0.2440	0.3837	0.2121
\mathfrak{C}_ρ	0.6514	0.5702	0.7302	0.5728
\mathfrak{D}_ρ	0.5029	0.4576	0.5734	0.4562

$$\mathfrak{C}(F, G)_{Chen_1} = \frac{\sum_{i=1}^n \left[\frac{1}{l_i} \sum_{j=1}^{l_i} \rho_{F\sigma(j)}(u_i) \rho_{G\sigma(j)}(u_i) \right]}{\left[\sum_{i=1}^n \left(\frac{1}{l_{F_i}} \sum_{j=1}^{l_{F_i}} \rho_{F\sigma(j)}^2(u_i) \right) \right]^{1/2} \left[\sum_{i=1}^n \left(\frac{1}{l_{G_i}} \sum_{j=1}^{l_{G_i}} \rho_{G\sigma(j)}^2(u_i) \right) \right]^{1/2}} \quad (5.1)$$

Consider the two HFSs $F_1 = \{(u, [0.0])\}$, $F_2 = \{(u, [0.2, 0.3])\}$ for $U = \{u\}$. Using the formula 5.1, since the informational energy of F is equal to 0, then the correlation coefficient between F_1 and F_2 cannot compute.

For $h_{\mathcal{F}}(u) = \{\rho_{\mathcal{F}}(u)\}$, take the HFS $\mathcal{F} = \{(u, h_{\mathcal{F}}(u)) : u \in U\}$. Now, we can generalize $h_{\mathcal{F}}(u)$ to $\mathcal{F} = \{(\rho_{\mathcal{F}}(u), \sqrt{1 - \rho_{\mathcal{F}}^2(u)})\}$. Therefore, \mathcal{F} is extended to Pythagorean hesitant fuzzy set (PHFS). Zheng et al. [78] developed the LCME method to solve the problems caused by the different cardinalities of the PHFEs. According to this method, the KK of the PHFS is as follows:

$$\mathfrak{C}(F, G)_{Chen_2} = \frac{\frac{1}{n} \sum_{i=1}^n \left(\frac{1}{3S_i} \sum_{j=1}^{S_i} \left[\rho_{F\sigma(j)}^2 \rho_{G\sigma(j)}^2 + \eta_{F\sigma(j)}^2 \eta_{G\sigma(j)}^2 + \theta_{F\sigma(j)}^2 \theta_{G\sigma(j)}^2 \right] \right)}{\sqrt{\frac{1}{n} \sum_{i=1}^n \left(\rho_{F_j}^4(u_i) + \eta_{F_j}^4(u_i) + \theta_{F_j}^4(u_i) \right)} \sqrt{\frac{1}{n} \sum_{i=1}^n \left(\rho_{G_j}^4(u_i) + \eta_{G_j}^4(u_i) + \theta_{G_j}^4(u_i) \right)}} \quad (5.2)$$

Consider the two PHFSs $F_1 = \{(u, [0.0])\}$, $F_2 = \{(u, [0.2, 0.3])\}$ for $U = \{u\}$. Then, we can obtain $h_{F_1}(u) = \{(0, 1)\}$, $h_{F_2}(u) = \{(0.2, \sqrt{0.96}), (0.3, \sqrt{0.91})\}$. Using the formula 5.2, $\mathfrak{C}(F, G)_{Chen_2} = 0.935$.

If we use similar thinking for FHS and calculate with the same values, then $h_{F_1}(u) = \{(0, 1)\}$, $h_{F_2}(u) = \{(0.2, \sqrt{0.997}), (0.3, \sqrt{0.991})\}$, where $\mathcal{F} = \{(\rho_{\mathcal{F}}(u), \sqrt[3]{1 - \rho_{\mathcal{F}}^3(u)})\}$. Hence, we can obtain $\mathfrak{C}(F, G) = 0.9969$.

5.2. Advantages of the new approach

The following merits have been taken into account from the proposed correlation coefficient:

As mentioned above, the FFS is one of the generalizations of the classic set, FS, IFS, and PFS. As PFS is one of the most successful extensions, which is characterized by the degrees of membership and nonmembership satisfaction of the particular alternative concerning the criteria, such that their sum of the square is equal to or less than 1. However, there may be a situation where the decision-maker may provide the degree of membership and nonmembership of a particular attribute in such a way that their sum of squares is greater than 1. Therefore, this situation is not properly handled in the PFS. To overcome this shortcoming, FFS theory is one of the more general and can handle not only incomplete information but also indeterminate information and inconsistent information, which exist commonly in real situations. Therefore, the Fermatean fuzzy information decision-making is more suitable for real scientific and engineering applications.

Furthermore, it has been observed from the existing studies that various researchers proposed an algorithm by using a correlation coefficient for PFSs. As mentioned above, some situations cannot be represented by PFSs, so their corresponding algorithm may not give appropriate results.

The correlation coefficient for PFSs is a special case of the correlation coefficient of FFSs. Accordingly, obtaining correlation coefficients according to IVFHS gives wider opportunities to deal with uncertainty. Therefore, the proposed correlation coefficient is more generalized and suitable to solve real-life problems more accurately than the existing ones.

6. Conclusion

In this paper, a correlation coefficient for IVFHSs has been proposed. The short-coming of the existing operators have also been highlighted in this paper. Based on that, the present paper have extended the theory of correlation coefficient from PFS to the FFSs in which the constraint condition of sum of membership and nonmembership degrees be less than one has been relaxed. Numerical example have been given that demonstrate that the proposed correlation coefficient can easily handle the situation where the existing correlation coefficient in PFS environment fails. The proposed correlation coefficient in FFS has been developed by taking the degree

of membership, nonmembership, and their degree of hesitation between them. Also to deal with the situations where the elements in a set are correlative, a weighted correlation coefficient has been defined. To demonstrate the efficiency of the proposed coefficients, numerical examples of pattern recognition and multicriteria decision-making have been taken. From the experimental studies, it has been concluded that the proposed correlation coefficient in the FFS environment can suitably handle the real-life decision-making problem with their targets. In future studies, it will be possible to create new concepts with the methods proposed in this article and ideas from studies such as [37–39, 47, 57].

Acknowledgments

The author would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

References

- [1] Abdullah S, Qiyas M, Naeem MM, Liu Y. Pythagorean Cubic fuzzy Hamacher aggregation operators and their application in green supply selection problem[J]. *AIMS Mathematics* 2022; 7 (3): 4735-4766. <https://doi.org/10.3934/math.2022263>
- [2] Akram M, Adeel A, Alcantud JCR. Hesitant fuzzy N-soft sets: A new model with applications in decision-making. *Journal of Intelligent & Fuzzy Systems* 2019; 36 (6): 6113-6127.
- [3] Akram M, Adeel A, Alcantud JCR. Group decision-making methods based on hesitant N-soft sets. *Expert Systems with Applications* 2019; 115; 95-105.
- [4] Atanassov KD. Intuitionistic fuzzy sets. *Fuzzy Sets and Systems* 1986; 20 (1): 87-96. [https://doi.org/10.1016/S0165-0114\(86\)80034-3](https://doi.org/10.1016/S0165-0114(86)80034-3)
- [5] Bustince H, Burillo P. Correlation of interval-valued intuitionistic fuzzy sets. *Fuzzy Sets Systems* 1995; 74 (2): 237-244.
- [6] Chen Y, Peng X, Guan G, Jiang H. Approaches to multiple attribute decision making based on the correlation coefficient with dual hesitant fuzzy information. *Journal of Intelligent & Fuzzy Systems* 2014; 26 (5): 2547-2556.
- [7] Chen N, Xu Z, Xia M. Interval-valued hesitant preference relations and their applications to group decision making. *Knowledge-Based Systems* 2013; 37: 528-540.
- [8] Chen N, Xu Z, Xia M. Correlation coefficients of hesitant fuzzy sets and their applications to clustering analysis, *Applied Mathematical Modelling* 2013; 37 (4), 2197-2211.
- [9] Chen N, Xu Z. Properties of interval-valued hesitant fuzzy sets. *Journal of Intelligent & Fuzzy Systems* 2014; 27 (1): 143-158.
- [10] Chiang DA, Lin NP. Correlation of fuzzy sets. *Fuzzy Sets and Systems* 1999; 202 (2): 221-226. [https://doi.org/10.1016/S0165-0114\(97\)00127-9](https://doi.org/10.1016/S0165-0114(97)00127-9)
- [11] Das S, Malakar D, Kar S, Pal T. Correlation measure of hesitant fuzzy soft sets and their application in decision making. *Neural Computing and Applications* 2019; 31 (4): 1023-1039.
- [12] De A, Das S, Kar S. Multiple attribute decision making based on probabilistic interval-valued intuitionistic hesitant fuzzy set and extended TOPSIS method. *Journal of Intelligent & Fuzzy Systems* 2019; 37 (4); 5229-5248.
- [13] Donghai L, Yuanyuan L, Xiaohong C. Fermatean fuzzy linguistic set and its application in multicriteria decision making. *International Journal of Intelligent Systems* 2019; 34 (3): 878–894. <https://doi.org/10.1002/int.22079>
- [14] Donghai L, Yuanyuan L, Lizhen W. Distance measure for Fermatean fuzzy linguistic term sets based on linguistic scale function: An illustration of the TODIM and TOPSIS methods. *International Journal of Intelligent Systems* 2019; 34 (1): 2807–2834. <https://doi.org/10.1002/int.22162>

- [15] Ejegwa PA. Distance and similarity measures for Pythagorean fuzzy sets. *Granular Computing* 2018; 5: 225-238. <https://doi.org/10.1007/s41066-018-00149-z>
- [16] Ejegwa PA, Onasanya BO. Improved intuitionistic fuzzy composite relation and its application to medical diagnostic process. *Notes on Intuitionistic Fuzzy Sets* 2018; 25: 43-58. <https://doi.org/10.7546/nifs.2019.25.1.43-58>
- [17] Ejegwa PA. Novel Correlation Coefficient for Intuitionistic Fuzzy Sets and its Application to Multi-criteria Decision-making Problems. *International Journal of Fuzzy System Applications* 2021; 10 (2): 39-58.
- [18] Ejegwa PA, Jana C. Some New Weighted Correlation Coefficients between Pythagorean Fuzzy Sets and their Applications. *Pythagorean fuzzy sets* 2021; 39-64. https://doi.org/10.1007/978-981-16-1989-2_2
- [19] Farhadinia B. Correlation for dual hesitant fuzzy sets and dual interval-valued hesitant fuzzy set. *International Journal of Intelligent Systems* 2014; 29 (2): 184-205.
- [20] Garg H. A New Generalized Pythagorean Fuzzy Information Aggregation Using Einstein Operations and Its Application to Decision Making. *International Journal of Intelligent Systems* 2016; 31: 886-920.
- [21] Garg H. Some series of intuitionistic averaging aggregation operators. *SpringerPlus* 2016; 5 (1): 999. <https://doi.org/10.1186/s40064-016-2591-9>
- [22] Garg H. Novel correlation coefficients under the intuitionistic multiplicative environment and their applications to decision-making process. *Journal of Industrial & Management Optimization* 2018; 14 (4): 1501-1519. <https://doi.org/10.3934/jimo.2018018>
- [23] Garg H. A Novel Correlation Coefficients between Pythagorean Fuzzy Sets and Its Applications to Decision-Making. *International Journal of Intelligent Systems* 2016; 31 (12): 1234-1252. <https://doi.org/10.1002/int.21827>
- [24] Garg H, Kumar K. A novel correlation coefficient of intuitionistic fuzzy sets based on the connection number of set pair analysis and its application. *Scientia Iranica* 2018; 25 (4): 2373-2388.
- [25] Garg H, Rani D. A robust correlation coefficient measure of complex intuitionistic fuzzy sets and their applications in decision-making. *Applied Intelligence* 2019; 49 (2): 496-512.
- [26] Garg H, Shahzadi G, Akram M. Decision-Making Analysis Based on Fermatean Fuzzy Yager Aggregation Operators with Application in COVID-19 Testing Facility. *Mathematical Problems in Engineering* 2020; Article ID 7279027. <https://doi.org/10.1155/2020/7279027>
- [27] Ju Y, Liu X, Yang S. Interval-valued dual hesitant fuzzy aggregation operators and their applications to multiple attribute decision making. *Journal of Intelligent & Fuzzy Systems* 2014; 27 (3): 1203-1218.
- [28] Khan AA, Ashraf S, Abdullah S, Qiyas M, Luo J et al. Pythagorean Fuzzy Dombi Aggregation Operators and Their Application in Decision Support. *System Symmetry* 2019; 11: 383. <https://doi.org/10.3390/sym11030383>
- [29] Kirişçi M. Ω - Soft Sets and medical decision-making application. *International Journal of Computer Mathematics* 2021; 98(4): 690-704. <https://doi.org/10.1080/00207160.2020.1777404>
- [30] Kirişçi M. New Entropy and Distance Measures for Fermatean Fuzzy Soft Sets with Medical Decision-Making and Pattern Recognition Applications. <https://doi.org/10.21203/rs.3.rs-1796355/v1>
- [31] Kirişçi M. Fermatean Hesitant Fuzzy Sets with Medical Decision Making Application. <https://doi.org/10.21203/rs.3.rs-1151389/v2>
- [32] Kirişçi M., Demir I., Simşek N. Fermatean Fuzzy ELECTRE Multi-Criteria Group Decision-Making and Most Suitable Biomedical Material Selection. *Artificial Intelligence in Medicine* 2022. <https://doi.org/10.1016/j.artmed.2022.102278>
- [33] Kirişçi M., Simşek N. Interval-Valued Fermatean Hesitant Fuzzy Sets and Infectious Diseases Application. <https://doi.org/10.21203/rs.3.rs-1273874/v1>
- [34] Kirişçi M. A Case Study for medical decision making with the fuzzy soft sets. *Afrika Matematika* 2020; 31: 557-564. <https://doi.org/10.1007/s13370-019-00741-9>

- [35] Kirişci M. Correlation Coefficients of Fermatean Fuzzy Sets with a Medical Application. *Journal of Mathematical Sciences and Modelling* 2022; 5 (1): 16-23. <https://doi.org/10.33187/jmsm.1039613>
- [36] Kirişci M, Simşek N. Decision making method related to Pythagorean Fuzzy Soft Sets with infectious diseases application. *Journal of King Saud University - Computer and Information Sciences* 2021; <https://doi.org/10.1016/j.jksuci.2021.08.010>
- [37] Kiyas M, Naeem M, Abdullah S, Khan N, Ali A. Similarity Measures Based on q-Rung Linear Diophantine Fuzzy Sets and Their Application in Logistics and Supply Chain Management. *Journal of Mathematics* 2022; <https://doi.org/10.1155/2022/4912964>
- [38] Kiyas M, Saleem A. Sine trigonometric spherical fuzzy aggregation operators and their application in decision support system, TOPSIS, VIKOR. *The Korean Journal of Mathematics* 2021; 29: 137-167.
- [39] Kiyas M, Saleem A, Khan F, Naeem M. Banzhaf-Choquet-Copula-based aggregation operators for managing fractional orthotriple fuzzy information. *Alexandria Engineering Journal* 2022; 61 (6): 4659-4677.
- [40] Krishankumar R, Ravichandran KS, Kar S, Gupta P, Mehlatat MK. Interval-valued probabilistic hesitant fuzzy set for multi-criteria group decision-making. *Soft Computing* 2019; 23 (21): 10853-10879.
- [41] Krishankumar R, Saranya R, Nethra R, Ravichandran K, Kar S. A decision-making framework under probabilistic linguistic term set for multi-criteria group decision-making problem. *Journal of Intelligent & Fuzzy Systems* 2019; 36 (6): 5783-5795.
- [42] Liang Z, Shi P. Similarity measures on intuitionistic fuzzy sets. *Pattern Recognition Letters* 2003; 24 (15): 2687-2693. [https://doi.org/10.1016/S0167-8655\(03\)00111-9](https://doi.org/10.1016/S0167-8655(03)00111-9)
- [43] Liao H, Xu Z, Zeng XJ. Novel correlation coefficients between hesitant fuzzy sets and their application in decision making. *Knowledge-Based Systems* 2015; 82: 115-127. <https://doi.org/10.1016/j.knosys.2015.02.020>
- [44] Liu ST, Kao C. Fuzzy measures for correlation coefficient of fuzzy numbers. *Fuzzy Sets and Systems* 2002; 128 (2): 267-275. [https://doi.org/10.1016/S0165-0114\(01\)00199-3](https://doi.org/10.1016/S0165-0114(01)00199-3)
- [45] Meng F, Chen X. Correlation coefficients of hesitant fuzzy sets and their application based on fuzzy measures. *cognitive computing* 2015; 7 (4): 445-463. <https://doi.org/10.1007/s12559-014-9313-9>
- [46] Meng F, Wang C, Chen X, Zhang Q. Correlation coefficients of interval-valued hesitant fuzzy sets and their application based on the Shapley function. *International Journal of Intelligent Systems* 2016; 31 (1): 17-43.
- [47] Naeem M, Kiyas M, Al-Shomrani MM, Saleem A. Similarity Measures for Fractional Orthotriple Fuzzy Sets Using Cosine and Cotangent Functions and Their Application in Accident Emergency Response. *Mathematics* 2020; 8 (10): 1653. <https://doi.org/10.3390/math8101653>
- [48] Naeem M, Khan MA, Abdullah S, Qiyas M, Khan S. Extended TOPSIS method based on the entropy measure and probabilistic hesitant fuzzy information and their application in decision support system. *Journal of Intelligent & Fuzzy Systems* 2021; 40 (6): 11479-11490.
- [49] Peng X, Yang Y, Song J, Jiang Y. Pythagorean fuzzy soft set and its application. *Computer Engineering* 2015; 41 (7): 224-229. <https://doi.org/10.3969/j.issn.1000-3428.2015.07.043>
- [50] Peng X, Selvachandran G. Pythagorean fuzzy set: state of the art and future directions. *Artificial Intelligence Review* 2019; 52 (3): 1873-1927. <https://doi.org/10.1007/s10462-017-9596-9>
- [51] Roy J, Das S, Kar S, Pamučar D. An extension of the CODAS approach using the interval-valued intuitionistic fuzzy set for sustainable material selection in construction projects with incomplete weight information. *Symmetry* 2019; 11 (3): 393.
- [52] Saeed M, Saqlain M, Mehmood A, Naseer K, Yaqoob S. Multi-Polar Neutrosophic Soft Sets with Application in Medical Diagnosis and Decision-Making. *Neutrosophic Set Systems* 2020; 33 (3): 183-207.
- [53] Senapati T, Yager RR. Fermatean fuzzy sets. *Journal of Ambient Intelligence and Humanized Computing* 2020; 11 (2): 663-674.

- [54] Senapati T, Yager RR. Some new operations over Fermatean fuzzy numbers and application of Fermatean fuzzy WPM in multiple criteria decision making. *Informatica* 2019; 30 (1): 391-412.
- [55] Senapati T, Yager RR. Fermatean fuzzy weighted averaging/geometric operators and its application in multi-criteria decision-making methods. *Engineering Applications of Artificial Intelligence* 2019; 85: 112-121. <https://doi.org/10.1016/j.engappai.2019.05.012>
- [56] Shahzadi G, Akram M. Group decision-making for the selection of an antivirus mask under fermatean fuzzy soft information. *Journal of Intelligent & Fuzzy Systems* 2021; 40 (1): 1401-1416.
- [57] Shahzaib A, Saleem A, Aslam M, Kiyas M, Kutbi MA. Spherical fuzzy sets and its representation of spherical fuzzy t-norms and t-conorms. *Journal of Intelligent & Fuzzy Systems* 2019; 36 (6): 6089-6102.
- [58] Singh S, Ganie AH. On some correlation coefficients in Pythagorean fuzzy environment with applications. *International Journal of Intelligent Systems* 2020; 35 (4): 682-717.
- [59] Sun G, Guan X, Yi X, Zhou Z. Improvements on correlation coefficients of hesitant fuzzy sets and their applications. *cognitive computing* 2019; 11 (4): 529-544.
- [60] Thao NX. A new correlation coefficient of the intuitionistic fuzzy sets and its application. *Journal of Intelligent & Fuzzy Systems* 2018; 35 (2): 1959-1968. <https://doi.org/10.3233/JIFS-171589>
- [61] Torra V, Narukawa Y. On hesitant fuzzy sets and decision, in *Proc. IEEE International Conference on Fuzzy Systems*, Jeju Island, South Korea, 2009; 1378-1382.
- [62] Torra V. Hesitant fuzzy sets. *International Journal of Intelligent Systems* 2010; 25 (6): 529-539.
- [63] Tyagi SK. Correlation coefficient of dual hesitant fuzzy sets and its applications. *Applied Mathematical Modelling* 2015; 39 (22): 7082-7092.
- [64] Vander WD, Nachttegaal M, Kerre EE. A new similarity measure for image processing. *Journal of Computational Methods in Sciences and Engineering* 2003; 3 (2): 209-222.
- [65] Vander WD, Nachttegaal M, Kerre EE. Using similarity measures and homogeneity for the comparison of images. *Image and Vision Computing* 2004; 22 (9): 695-702. <https://doi.org/10.1016/j.imavis.2004.03.002>
- [66] Wei WG, Wang HJ, Lin R, Application of correlation coefficient to interval-valued intuitionistic fuzzy multiple attribute decision-making with incomplete weight information. *Knowledge and Information Systems* 2011; 26: 337-349.
- [67] Xia M, Xu Z. Hesitant fuzzy information aggregation in decision making. *International Journal of Approximate Reasoning* 2011; 52 (3): 395-407. <https://doi.org/10.1016/j.ijar.2010.09.002>
- [68] Xu Z, Xia M. On distance and correlation measures of hesitant fuzzy information. *International Journal of Intelligent Systems* 2011; 26 (5): 410-425.
- [69] Xu ZS, Chen J, Wu JJ. Cluster algorithm for intuitionistic fuzzy sets. *Information Sciences* 2008; 178 (19): 3775-3790. <https://doi.org/10.1016/j.ins.2008.06.008>
- [70] Yager RR. Pythagorean fuzzy subsets. *Proc Joint IFSA World Congress and NAFIPS. Annual Meeting*; Edmonton, Canada; 2013.
- [71] Yager RR. Pythagorean membership grade in multicriteria decision makng. *IEEE Transactions on Fuzzy Systems* 2014; 22 (4): 958-965. <https://doi.org/10.1109/TFUZZ.2013.2278989>
- [72] Yager RR. Multicriteria decision making with ordinal linguistic intuitionistic fuzzy sets for mobile apps. *Transactions on Fuzzy Systems* 2016; 24 (3): 590-599. <https://doi.org/10.1109/TFUZZ.2015.2463740>
- [73] Yager RR, Abbasov AM. Pythagorean membership grades, complex numbers and decision making. *International Journal of Intelligent Systems* 2013; 28 (5): 436-452.
- [74] Yahya M, Naeem M, Abdullah S, Qiyas M, Aamir M. A novel approach on the intuitionistic fuzzy rough frank aggregation operator-based EDAS method for multicriteria group decision-making. *Complexity* 2021; <https://doi.org/10.1155/2021/5534381>

- [75] Ye J. Correlation coefficient of dual hesitant fuzzy sets and its application to multiple attribute decision making. *Applied Mathematical Modelling* 20214; 38 (2); 659-666.
- [76] Zadeh LA. Fuzzy sets. *Inf Comp* 1965; 8: 338–353.
- [77] Zhang XL, Xu ZS. Extension of TOPSIS to multi-criteria decision making with Pythagorean fuzzy sets. *International Journal of Intelligent Systems* 2014; 29 (12): 1061-1078.
- [78] Zheng T, Zhang M, Li L, Wu Q, Zhou L. Correlation Coefficients of Interval-Valued Pythagorean Hesitant Fuzzy Sets and Their Applications. *IEEE Access* 2020; 8; 9271-9286. <https://doi.org/10.1109/ACCESS.2020.2964580>
- [79] Zhou Q, Mo H, Deng Y. A New Divergence Measure of Pythagorean Fuzzy Sets Based on Belief Function and Its Application in Medical Diagnosis. *Mathematics* 2020; 8 (1): 2227-7390. <https://doi.org/10.3390/math8010142>