

1-1-2023

## Metrical almost periodicity, metrical approximations of functions and applications

BELKACEM CHAOUCHI

MARKO KOSTIC

DANIEL VELINOV

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)




### Recommended Citation

CHAOUCHI, BELKACEM; KOSTIC, MARKO; and VELINOV, DANIEL (2023) "Metrical almost periodicity, metrical approximations of functions and applications," *Turkish Journal of Mathematics*: Vol. 47: No. 2, Article 25. <https://doi.org/10.55730/1300-0098.3393>

Available at: <https://journals.tubitak.gov.tr/math/vol47/iss2/25>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Metrical almost periodicity, metrical approximations of functions and applications

Belkacem CHAOUCHI<sup>1</sup> , Marko KOSTIĆ<sup>2</sup> , Daniel VELINOV<sup>3,\*</sup> 

<sup>1</sup>Laboratory of Energy and Intelligent Systems, Khemis Miliana University, Khemis Miliana, Algeria

<sup>2</sup>Department of Fundamental Sciences, Faculty of Technical Sciences,  
University of Novi Sad, Novi Sad, Serbia

<sup>3</sup>Department for Mathematics and Informatics, Faculty of Civil Engineering,  
Ss. Cyril and Methodius University in Skopje, Skopje, N. Macedonia

Received: 27.10.2022

Accepted/Published Online: 13.02.2023

Final Version: 09.03.2023

**Abstract:** In this paper, we analyze Levitan and Bebutov metrical approximations of functions  $F : \Lambda \times X \rightarrow Y$  by trigonometric polynomials and  $\rho$ -periodic type functions, where  $\emptyset \neq \Lambda \subseteq \mathbb{R}^n$ ,  $X$  and  $Y$  are complex Banach spaces, and  $\rho$  is a general binary relation on  $Y$ . We also analyze various classes of multidimensional Levitan almost periodic functions in general metric and multidimensional Bebutov uniformly recurrent functions in general metric. We provide several applications of our theoretical results to the abstract Volterra integro-differential equations and the partial differential equations.

**Key words:** Levitan metrical almost periodicity, Bebutov metrical almost periodicity, abstract Volterra integro-differential equations

### 1. Introduction and preliminaries

The notion of almost periodicity was introduced by the Danish mathematician H. Bohr around 1924-1926 and later generalized by many other authors (see the research monographs [10, 24, 26, 31, 33, 40, 41, 46, 47, 60] for more details about almost periodic functions and their applications). Suppose that  $(X, \|\cdot\|)$  is a complex Banach space, and  $F : \mathbb{R}^n \rightarrow X$  is a continuous function, where  $n \in \mathbb{N}$ . Then it is said that  $F(\cdot)$  is almost periodic if for each  $\epsilon > 0$  there exists a finite real number  $l > 0$  such that for each  $\mathbf{t}_0 \in \mathbb{R}^n$  there exists  $\tau \in B(\mathbf{t}_0, l) \equiv \{\mathbf{t} \in \mathbb{R}^n : |\mathbf{t} - \mathbf{t}_0| \leq l\}$  such that

$$\|F(\mathbf{t} + \tau) - F(\mathbf{t})\| \leq \epsilon, \quad \mathbf{t} \in \mathbb{R}^n;$$

here,  $|\cdot - \cdot|$  denotes the Euclidean distance in  $\mathbb{R}^n$ . Equivalently,  $F(\cdot)$  is almost periodic if for any sequence  $(\mathbf{b}_k)$  in  $\mathbb{R}^n$  there exists a subsequence  $(\mathbf{a}_k)$  of  $(\mathbf{b}_k)$  such that the sequence of translations  $(F(\cdot + \mathbf{a}_k))$  converges in  $C_b(\mathbb{R}^n : X)$ , the Banach space of all bounded continuous functions on  $\mathbb{R}^n$ , equipped with the sup-norm. Any trigonometric polynomial in  $\mathbb{R}^n$  is almost periodic, and a continuous function  $F(\cdot)$  is almost periodic if there exists a sequence of trigonometric polynomials in  $\mathbb{R}^n$  which converges uniformly to  $F(\cdot)$ .

A continuous function  $F : \mathbb{R}^n \rightarrow X$  is said to be uniformly recurrent (uniformly Poisson stable) if there exists a sequence  $(\tau_k)$  in  $\mathbb{R}^n$  such that  $\lim_{k \rightarrow +\infty} |\tau_k| = +\infty$  and  $\lim_{k \rightarrow +\infty} F(\mathbf{t} + \tau_k) = F(\mathbf{t})$ , uniformly in

\*Correspondence: velinovd@gf.ukim.edu.mk

2010 AMS Mathematics Subject Classification: 42A75, 43A60, 47D99

$\mathbf{t} \in \mathbb{R}^n$  (uniformly in  $\mathbf{t} \in K$ , for any compact subset  $K \subseteq \mathbb{R}^n$ ). Any almost periodic function is uniformly recurrent and any uniformly recurrent function is uniformly Poisson stable, while the converse statements are not true in general; cf. also [15, 25, 34] for some recent results about almost periodic functions and uniformly recurrent functions.

Further on, let  $F : \mathbb{R}^n \rightarrow X$  be a continuous function. Then it is said that the function  $F(\cdot)$  is almost automorphic if for every sequence  $(\mathbf{b}_k)$  in  $\mathbb{R}^n$  there exist a subsequence  $(\mathbf{a}_k)$  of  $(\mathbf{b}_k)$  and a mapping  $G : \mathbb{R}^n \rightarrow X$  such that

$$\lim_{k \rightarrow \infty} F(\mathbf{t} + \mathbf{a}_k) = G(\mathbf{t}) \quad \text{and} \quad \lim_{k \rightarrow \infty} G(\mathbf{t} - \mathbf{a}_k) = F(\mathbf{t}), \tag{1.1}$$

pointwisely for  $\mathbf{t} \in \mathbb{R}^n$ . The range of an almost automorphic function  $F(\cdot)$  is relatively compact in  $X$ , and the limit function  $G(\cdot)$  is bounded on  $\mathbb{R}^n$  but not necessarily continuous on  $\mathbb{R}^n$ . If the convergence of limits appearing in (1.1) is uniform on compact subsets of  $\mathbb{R}^n$ , then we say that the function  $F(\cdot)$  is compactly almost automorphic. It is well known that an almost automorphic function  $F(\cdot)$  is compactly almost automorphic if  $F(\cdot)$  is uniformly continuous (cf. [16] for further information about the multidimensional almost automorphic functions and their applications).

On the other hand, the class of almost periodic functions can be generalized following the approach of B. M. Levitan (see, e.g., [41] for the one-dimensional setting): Suppose that  $F : \mathbb{R}^n \rightarrow X$  is a continuous function,  $N > 0$  and  $\epsilon > 0$ . Then a point  $\tau \in \mathbb{R}^n$  is said to be an  $\epsilon, N$ -almost period of function  $F(\cdot)$  if

$$\|F(\mathbf{t} + \tau) - F(\mathbf{t})\| \leq \epsilon \text{ for all } \mathbf{t} \in \mathbb{R}^n \text{ with } |\mathbf{t}| \leq N;$$

denote by  $E(\epsilon, N)$  the set consisting of all  $\epsilon, N$ -almost periods of function  $F(\cdot)$ . Let us say, maybe for the first time in the existing literature, that a continuous function  $F(\cdot)$  is Levitan pre-almost periodic if for each  $N > 0$  and  $\epsilon > 0$ , there exists a finite real number  $l > 0$  such that for each  $\mathbf{t}_0 \in \mathbb{R}^n$  there exists  $\tau \in B(\mathbf{t}_0, l) \cap E(\epsilon, N)$ , i.e. the set  $E(\epsilon, N)$  is relatively dense in  $\mathbb{R}^n$  for each  $N > 0$  and  $\epsilon > 0$ . It is worth noting that B. Ya. Levin has shown, in [39], that the sum of two Levitan pre-almost periodic functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  need not be Levitan pre-almost periodic, in general. In the definition of a Levitan ( $N$ -)almost periodic function  $F : \mathbb{R}^n \rightarrow X$ , we additionally require that, for every real numbers  $N > 0$  and  $\epsilon > 0$ , there exist a finite real number  $\eta > 0$  and a relatively dense set  $E_{\eta, N}$  of  $(\eta, N)$ -almost periods of  $F(\cdot)$  such that  $E_{\eta, N} \pm E_{\eta, N} \subseteq E(\epsilon, N)$ ; cf. [41, condition (2), p. 54, l. 3] and the corresponding footnote for more details concerning this issue in the one-dimensional setting. Due to R. Yuan’s result [59, theorem 3.1], we know that a bounded continuous function  $f : \mathbb{R} \rightarrow X$  is compactly almost automorphic if  $f(\cdot)$  is Levitan  $N$ -almost periodic (cf. also A. Reich [49] and references cited therein for more details about the relationship between the almost automorphic functions and the Levitan  $N$ -almost periodic functions on topological groups).

The notion of a recurrent function in the continuous Bebutov system is based on the use of topology of uniform convergence on compact sets (cf. also Subsection 2.3.9 in the monograph [9] by G. Bertotti and I. D. Mayergoyz, the paper [22] by L. I. Danilov and references cited therein for further information in this direction). A uniformly recurrent function is also called pseudoperiodic by H. Bohr, which has been accepted by many other authors later on; a recurrent function in the continuous Bebutov system is also called (uniformly) Poisson-stable motion by M. V. Bebutov. The Levitan almost periodic solutions and the uniformly Poisson stable solutions for various classes of (abstract) differential equations have been sought in many research articles so far; see, e.g., [17, 18, 42, 43, 45, 54–56] and the references cited therein; it is also worth noting that M. Akhmet, M.

Tleubergenova, and A. Zhamanshin have recently analyzed the existence and uniqueness of modulo periodic Poisson stable solutions of quasilinear differential equations in [3].

Concerning the Poisson stability of motions of dynamical systems and solutions of differential equations, we would like to specifically mention the research monographs [51, 52] by B. A. Shcherbakov. It is also worth noticing that T. Caraballo and D. Cheban have analyzed the existence and uniqueness of Levitan/Bohr almost periodic (almost automorphic) solutions of the second-order monotone differential equations in [12]. The Poisson stability of motions for monotone nonautonomous dynamical systems and of solutions for certain classes of monotone nonautonomous differential equations has been analyzed by D. Cheban and Z. Liu in [20]; cf. also [19, 51–53, 56–58] and the references cited therein for related results obtained within the theory of dynamical systems. Finally, let us note that the interpolation by Levitan almost periodic functions was considered by S. Hartman in [28] (1974), while the difference property for perturbations of vector-valued Levitan almost periodic functions was considered by B. Basit and H. Günzler in [8]; see also M. G. Lyubarskii [43].

On the other hand, the first systematic study of metrical almost periodicity was conducted by the second named author in 2021 ([34]). The Stepanov, Weyl, and Besicovitch classes of metrical  $\rho$ -almost periodic type functions have been considered in [35], [36], and [37], respectively. In our joint research article [14] with B. Chaouchi and D. Velinov, we have recently studied the metrical approximations of functions  $F : \Lambda \times X \rightarrow Y$  by trigonometric polynomials and  $\rho$ -periodic type functions, where  $\emptyset \neq \Lambda \subseteq \mathbb{R}^n$ ,  $X$  and  $Y$  are complex Banach spaces, and  $\rho$  is a general binary relation on  $Y$ ; we have also introduced and analyzed various classes of almost periodic functions and uniformly recurrent functions in general metric therein.

The main aim of this research article is to continue the analysis raised in [14] by investigating the Levitan and Bebutov metrical approximations of functions  $F : \Lambda \times X \rightarrow Y$  by trigonometric polynomials and  $\rho$ -periodic type functions. As mentioned in the abstract, we analyze here various classes of multidimensional Levitan almost periodic functions in general metric and multidimensional Bebutov uniformly recurrent functions in general metric.

The organization and main ideas of this paper can be briefly described as follows. After explaining the notation and terminology used throughout the paper as well as the main concepts necessary for understanding anything that follows, we introduce the basic function spaces of metrically almost periodic functions in the sense of Levitan/Bebutov approach in the second section of paper; cf. Definition 2.1 and Definitions 2.4–2.5. The main structural results established in this section are Propositions 2.6, 2.9, and 2.11; cf. also Examples 2.7, 2.8, and 2.10. Levitan  $(N, c)$ -almost periodic functions and uniformly Poisson  $c$ -stable functions [multidimensional Levitan  $N$ -almost periodic functions] are specifically analyzed in Subsection 2.1 [Subsection 2.2]. In Subsection 2.2, we introduce the spaces of strongly Levitan  $N$ -almost periodic functions of type 1, the Levitan  $N$ -almost periodic functions of type 1, and the strongly Levitan  $N$ -almost periodic functions. We show that these spaces have the linear vector structure and propose an open problem whether the Levitan  $N$ -almost periodic functions form a vector space with the usual operations if  $n \geq 2$ . By a simple counterexample, we show that the Bogolybov theorem (see, e.g., [41, pp. 55–57]) cannot be straightforwardly extended to the higher-dimensional case  $n \geq 2$ ; cf. also Proposition 2.15 and Example 2.16.

In Section 3, we present several applications of our theoretical results to the abstract Volterra integro-differential equations. Subsection 3.1 is devoted to the study of invariance of Levitan  $N$ -almost like periodicity under the actions of the infinite convolution products and certain applications to the abstract Cauchy problems without initial conditions; Subsection 3.2 investigates the convolution invariance of certain kinds of multidimensional

mensional Levitan  $N$ -almost periodic type functions (the second application of this subsection, concerning the fractional diffusion-wave equations with Caputo-Dzhrbashyan fractional derivatives, and the third application of this subsection, concerning the biharmonic partial differential operator, are essentially new and not considered anywhere else in the existing literature). In this subsection, we also reconsider several results established recently by A. Nawrocki in [45]. Subsection 3.3 continues our analysis of the wave equation in  $\mathbb{R}^n$ ; the final section of the paper is reserved for some comments and final remarks about the introduced spaces of Levitan  $N$ -almost periodic type functions. We propose several open problems to our readers, providing also a great number of important references concerning the subjects under our consideration (we feel it is our duty to say that the article is maybe a little bit oversaturated with definitions as well as that we have not been able to properly shorten some notions; for example, we work with the class of Levitan  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \Lambda', \rho, \mathcal{P}_{\mathbf{K}})$ -almost periodic functions henceforth but it is not clear whether we can so simply drop any of the parameters used above from the definition of the class).

**Notation and terminology.** Suppose that  $X, Y, Z$  and  $T$  are given nonempty sets. Let us recall that a binary relation between  $X$  and  $Y$  is any subset  $\rho \subseteq X \times Y$ . If  $\rho \subseteq X \times Y$  and  $\sigma \subseteq Z \times T$  with  $Y \cap Z \neq \emptyset$ , then we define  $\sigma \cdot \rho = \sigma \circ \rho \subseteq X \times T$  by  $\sigma \circ \rho := \{(x, t) \in X \times T : \exists y \in Y \cap Z \text{ such that } (x, y) \in \rho \text{ and } (y, t) \in \sigma\}$ . As is well known, the domain and range of  $\rho$  are defined by  $D(\rho) := \{x \in X : \exists y \in Y \text{ such that } (x, y) \in \rho\}$  and  $R(\rho) := \{y \in Y : \exists x \in X \text{ such that } (x, y) \in \rho\}$ , respectively;  $\rho(x) := \{y \in Y : (x, y) \in \rho\}$  ( $x \in X$ ),  $x \rho y \Leftrightarrow (x, y) \in \rho$ . If  $\rho$  is a binary relation on  $X$  and  $n \in \mathbb{N}$ , then we define  $\rho^n$  inductively. Set  $\rho(X') := \{y : y \in \rho(x) \text{ for some } x \in X'\}$  ( $X' \subseteq X$ ).

We will always assume henceforth that  $(X, \|\cdot\|)$  and  $(Y, \|\cdot\|_Y)$  are complex Banach spaces,  $n \in \mathbb{N}$ ,  $\emptyset \neq \Lambda \subseteq \mathbb{R}^n$ , and  $\mathcal{B}$  is a nonempty collection of nonempty subsets of  $X$  satisfying that for each  $x \in X$  there exists  $B \in \mathcal{B}$  such that  $x \in B$ . For the sequel, we set

$$\Lambda'' := \{\tau \in \mathbb{R}^n : \tau + \Lambda \subseteq \Lambda\}.$$

By  $L(X, Y)$ , we denote the Banach space of all bounded linear operators from  $X$  into  $Y$ ,  $L(X, X) \equiv L(X)$ ;  $I$  denotes the identity operator on  $Y$ . Define  $\mathbb{N}_n := \{1, \dots, n\}$ . If  $A$  and  $B$  are nonempty sets, then we define  $B^A := \{f | f : A \rightarrow B\}$ . By  $\|\cdot\|_\infty$  we denote the sup-norm; the symbol  $f|_K(\cdot)$  denotes the restriction of a function  $f(\cdot)$  to a nonempty subset  $K$  of its domain.

Suppose now that  $\nu : \Lambda \rightarrow (0, \infty)$  and the function  $1/\nu(\cdot)$  is locally bounded. Then the vector space  $C_{b,\nu}(\Lambda : Y)$  consists of all continuous functions  $u : \Lambda \rightarrow Y$  such that  $\sup_{\mathbf{t} \in \Lambda} \|u(\mathbf{t})\|_Y \nu(\mathbf{t}) < +\infty$ . Equipped with the norm  $\|\cdot\| := \sup_{\mathbf{t} \in \Lambda} \|\nu(\mathbf{t}) \cdot (\mathbf{t})\|_Y$ ,  $C_{b,\nu}(\Lambda : Y)$  is a Banach space.

We need to recall the following notion ([33]):

**Definition 1.1** (i) Let  $\omega \in \mathbb{R}^n \setminus \{0\}$ ,  $\rho$  be a binary relation on  $Y$  and  $\omega \in \Lambda''$ . A continuous function  $F : \Lambda \times X \rightarrow Y$  is said to be  $(\omega, \rho)$ -periodic [ $\rho$ -periodic] if  $F(\mathbf{t} + \omega; x) \in \rho(F(\mathbf{t}; x))$ ,  $\mathbf{t} \in \Lambda$ ,  $x \in X$  [there exists  $\omega \in (\mathbb{R}^n \setminus \{0\}) \cap \Lambda''$  such that  $F(\cdot; \cdot)$  is  $(\omega, \rho)$ -periodic].

(ii) Let  $\omega_j \in \mathbb{R} \setminus \{0\}$ ,  $\rho_j \in \mathbb{C} \setminus \{0\}$  be a binary relation on  $Y$  and  $\omega_j e_j + \Lambda \subseteq \Lambda$  ( $1 \leq j \leq n$ ). A continuous function  $F : \Lambda \times X \rightarrow Y$  is said to be  $(\omega_j, \rho_j)_{j \in \mathbb{N}_n}$ -periodic [ $(\rho_j)_{j \in \mathbb{N}_n}$ -periodic] if  $F(\mathbf{t} + \omega_j e_j; x) \in \rho_j(F(\mathbf{t}; x))$ ,  $\mathbf{t} \in \Lambda$ ,  $x \in X$ ,  $j \in \mathbb{N}_n$  [there exist nonzero real numbers  $\omega_j$  such that  $\omega_j e_j \in \Lambda''$  for all  $j \in \mathbb{N}_n$  and  $F(\cdot; \cdot)$  is  $(\omega_j, \rho_j)_{j \in \mathbb{N}_n}$ -periodic].

(iii) Let  $\omega_j \in \mathbb{R} \setminus \{0\}$ , and  $\omega_j e_j + \Lambda \subseteq \Lambda$  ( $1 \leq j \leq n$ ). A continuous function  $F : \Lambda \times X \rightarrow Y$  is said to be periodic if  $F(\cdot; \cdot)$  is  $(\rho_j)_{j \in \mathbb{N}_n}$ -periodic with  $\rho_j = I$  for all  $j \in \mathbb{N}_n$ .

Finally, let us recall that a trigonometric polynomial  $P : \Lambda \times X \rightarrow Y$  is any linear combination of functions like

$$e^{i[\lambda_1 t_1 + \lambda_2 t_2 + \dots + \lambda_n t_n]} c(x),$$

where  $\lambda_i$  are real numbers ( $1 \leq i \leq n$ ) and  $c : X \rightarrow Y$  is a continuous mapping.

## 2. Metrical approximations: Levitan and Bebutov concepts

In this section, we assume that  $\emptyset \neq \Lambda \subseteq \mathbb{R}^n$  and  $\varphi : [0, \infty) \rightarrow [0, \infty)$ . If  $\emptyset \neq K \subseteq \mathbb{R}^n$  is a compact set and  $K \cap \Lambda \neq \emptyset$ , then we assume that the function  $\mathbb{F}_K : K \cap \Lambda \rightarrow [0, \infty)$  and the pseudometric space  $\mathcal{P}_K = (P_K, d_K)$ , where  $P_K \subseteq [0, \infty)^{K \cap \Lambda}$ , are given. Define  $\|g\|_{P_K} := d_K(g, 0)$  for any  $g \in P_K$ .

The following notion plays an important role in our analysis (cf. also [14, definition 2.1]):

**Definition 2.1** Suppose that  $\emptyset \neq \Lambda \subseteq \mathbb{R}^n$  and  $F : \Lambda \times X \rightarrow Y$ . Then we say that  $F(\cdot; \cdot)$  is: Levitan strongly  $(\varphi, \mathbb{F}_K, \mathcal{B}, \mathcal{P}_K)$ -almost periodic (Levitan semi- $(\varphi, \rho, \mathbb{F}_K, \mathcal{B}, \mathcal{P}_K)$ -periodic, Levitan semi- $(\varphi, \rho_j, \mathbb{F}_K, \mathcal{B}, \mathcal{P}_K)_{j \in \mathbb{N}_n}$ -periodic) if for each  $B \in \mathcal{B}$  and for each nonempty compact set  $K \subseteq \mathbb{R}^n$  such that  $K \cap \Lambda \neq \emptyset$  there exists a sequence  $(P_k^{B,K} : \Lambda \times X \rightarrow Y)$  of trigonometric polynomials ( $\rho$ -periodic functions  $(P_k^{B,K} : \Lambda \times X \rightarrow Y)$ ,  $(\rho_j)_{j \in \mathbb{N}_n}$ -periodic functions  $(P_k^{B,K} : \Lambda \times X \rightarrow Y)$ ) such that  $\mathbb{F}_K(\cdot)[\varphi(\|P_k^{B,K}(\cdot; x) - F(\cdot; x)\|_Y)]|_{K \cap \Lambda} \in P_K$  for all  $x \in X$ , and

$$\lim_{k \rightarrow +\infty} \sup_{x \in B} \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \|P_k^{B,K}(\cdot; x) - F(\cdot; x)\|_Y \right) \right] \Big|_{K \cap \Lambda} \right\|_{P_K} = 0.$$

As in our former research studies, we omit the term “ $\varphi$ ” if  $\varphi(x) \equiv x$ , “ $\rho$ ” if  $\rho = I$ , the term “ $\mathcal{B}$ ” if  $X = \{0\}$  and  $\mathcal{B} = \{X\}$ . We will also omit the term “ $\mathbb{F}_K$ ” if  $\mathbb{F}_K \equiv 1$  for each nonempty compact set  $K \subseteq \mathbb{R}^n$  such that  $K \cap \Lambda \neq \emptyset$  (in the case that  $\varphi(x) \equiv x$ ,  $\rho = I$ ,  $P_K = C_b(K \cap \Lambda)$  and  $\mathbb{F}_K \equiv 1$  for each nonempty compact set  $K \subseteq \mathbb{R}^n$  such that  $K \cap \Lambda \neq \emptyset$ , the notion of Levitan strong  $(\varphi, \mathbb{F}_K, \mathcal{B}, \mathcal{P}_K)$ -almost periodicity is completely regardless; for example, any continuous function  $F : \mathbb{R}^n \rightarrow Y$  is Levitan strongly  $(x, \mathbb{F}_K, \mathcal{P}_K)$ -almost periodic due to the Weierstrass approximation theorem).

The following result can be proven in a similar fashion to [14, proposition 2.3]:

**Proposition 2.2** Suppose that  $\emptyset \neq \Lambda \subseteq \mathbb{R}^n$ ,  $F : \Lambda \times X \rightarrow Y$ ,  $h : Y \rightarrow Z$  is Lipschitz continuous,  $\varphi(\cdot)$  is monotonically increasing and there exists a function  $\varphi : [0, \infty) \rightarrow [0, \infty)$  such that  $\varphi(xy) \leq \varphi(x)\varphi(y)$  for all  $x, y \geq 0$ . Let the assumptions (C0)-(C1) hold for every nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$ , where:

(C0-K) The assumptions  $0 \leq f \leq g$  and  $g \in P_K$  imply  $f \in P_K$  and  $\|f\|_{P_K} \leq \|g\|_{P_K}$ .

(C1-K) If  $f \in P_K$ , then  $d'f \in P_K$  for all reals  $d' \geq 0$ , and there exists a finite real constant  $d_K > 0$  such that  $\|d'f\|_{P_K} \leq d_K(1 + d')\|f\|_{P_K}$  for all reals  $d' \geq 0$  and all functions  $f \in P_K$ .

Then we have the following:

- (i) Suppose that  $F(\cdot; \cdot)$  is Levitan semi- $(\varphi, \rho, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \mathcal{P}_{\mathbf{K}})$ -periodic (Levitan semi- $(\varphi, \rho_j, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \mathcal{P}_{\mathbf{K}})_{j \in \mathbb{N}_n}$ -periodic), and  $h \circ \rho \subseteq \rho \circ h$  ( $h \circ \rho_j \subseteq \rho_j \circ h$  for  $1 \leq j \leq n$ ). Then the function  $h \circ F : \Lambda \times X \rightarrow Y$  is likewise Levitan semi- $(\varphi, \rho, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \mathcal{P}_{\mathbf{K}})$ -periodic (Levitan semi- $(\varphi, \rho_j, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \mathcal{P}_{\mathbf{K}})_{j \in \mathbb{N}_n}$ -periodic).
- (ii) Suppose that  $X = \{0\}$ , there exists a finite real constant  $c > 0$  such that  $\varphi(x + y) \leq c[\varphi(x) + \varphi(y)]$  for all  $x, y \geq 0$ ,  $\varphi(\cdot)$  is continuous at the point zero and, for every nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$ , we have

$$\mathbb{F}_K \in P_K \quad \text{and} \quad \lim_{\epsilon \rightarrow 0^+} \|\epsilon \mathbb{F}_K(\cdot)\|_{P_K} = 0.$$

Suppose, further, that the function  $F(\cdot)$  is Levitan strongly  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \mathcal{P}_{\mathbf{K}})$ -almost periodic and the assumption (C2-K) holds, where:

(C2-K) There exists a finite real constant  $e_K > 0$  such that the assumptions  $f, g \in P_K$  and  $0 \leq w \leq d'[f + g]$  for some finite real constant  $d' > 0$  imply  $w \in P_K$  and  $\|w\|_{P_K} \leq e_K(1 + d')[\|f\|_{P_K} + \|g\|_{P_K}]$ .

Then the function  $(h \circ F)(\cdot)$  is likewise Levitan strongly  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \mathcal{P}_{\mathbf{K}})$ -almost periodic.

We continue by providing the following illustrative example:

**Example 2.3** Let us recall that A. Haraux and P. Souplet have proven, in [27, theorem 1.1], that the function

$$f(t) := \sum_{m=1}^{\infty} \frac{1}{m} \sin^2\left(\frac{t}{2^m}\right), \quad t \in \mathbb{R}, \tag{2.1}$$

is not Besicovitch- $p$ -almost periodic for any finite exponent  $p \geq 1$ , as well as that  $f(\cdot)$  is uniformly recurrent and uniformly continuous (cf. also [31]). We also know that the function  $f(\cdot)$  is slowly  $p$ -semiperiodic in variation ( $1 \leq p < +\infty$ ); see [14] for the notion and more details. Let  $\epsilon_0 > 0$  be a fixed real number; using the elementary inequality  $|\sin t| \leq |t|$ ,  $t \in \mathbb{R}$ , it readily follows that the function  $f(\cdot)$  is Levitan semi- $(x, \mathbb{I}, \mathbb{F}_{\mathbf{K}}, \mathcal{P}_K)$ -semiperiodic with  $\mathbb{F}_{[-N, N]} \equiv N^{-2-\epsilon_0}$  and  $\mathcal{P}_{[-N, N]} := C[-N, N]$  for  $N > 0$ .

Similarly, any continuous function  $F : \mathbb{R}^n \rightarrow \mathbb{C}$  given by  $F(\mathbf{t}) = \sum_{m=1}^{\infty} P_m(\mathbf{t})$ ,  $\mathbf{t} \in \mathbb{R}^n$ , where  $P_m(\cdot)$  is a trigonometric polynomial ( $m \in \mathbb{N}$ ), is Levitan semi- $(x, \mathbb{I}, \mathbb{F}_{\mathbf{K}}, \mathcal{P}_K)$ -semiperiodic with the constant function  $\mathbb{F}_{[-N, N]^n} \equiv c_N > 0$  appropriately chosen and  $\mathcal{P}_{[-N, N]^n} := C([-N, N]^n)$  for  $N > 0$ , provided that there exists a summable sequence of nonnegative real numbers  $(a_m)$  and a continuous function  $G : \mathbb{R}^n \rightarrow \mathbb{C}$  such that  $|F(\mathbf{t})| \leq |G(\mathbf{t})| \cdot \sum_{m=1}^{\infty} a_m$ ,  $\mathbf{t} \in \mathbb{R}^n$ .

We continue by introducing the following notion (cf. also [14, definitions 2.5 and 2.8]):

**Definition 2.4** Suppose that  $\mathbb{R}$  is any collection of sequences in  $\Lambda''$ ,  $F : \Lambda \times X \rightarrow Y$  and  $\varphi : [0, \infty) \rightarrow [0, \infty)$ . Then we say that the function  $F(\cdot; \cdot)$  is Levitan  $(\varphi, \mathbb{R}, \mathcal{B}, \mathbb{F}_{\mathbf{K}}, \mathcal{P}_{\mathbf{K}})$ -normal if for every set  $B \in \mathcal{B}$ , for every nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$  and for every sequence  $(\mathbf{b}_k)_{k \in \mathbb{N}}$  in  $\mathbb{R}$ , there exists a subsequence  $(\mathbf{b}_{k_m})_{m \in \mathbb{N}}$  of  $(\mathbf{b}_k)_{k \in \mathbb{N}}$  such that, for every  $\epsilon > 0$ , there exists an integer  $m_0 \in \mathbb{N}$  such that, for every integers  $m, m' \geq m_0$ , we have  $\mathbb{F}_K(\cdot)[\varphi(\|F(\cdot + \mathbf{b}_{k_m}; x) - F(\cdot + \mathbf{b}_{k_{m'}}; x)\|_Y)]|_{K \cap \Lambda} \in P_K$  for all  $x \in X$ , and

$$\sup_{x \in B} \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \|F(\cdot + \mathbf{b}_{k_m}; x) - F(\cdot + \mathbf{b}_{k_{m'}}; x)\|_Y \right) \right] \Big|_{K \cap \Lambda} \right\|_{P_K} < \epsilon.$$

**Definition 2.5** Suppose that  $\emptyset \neq \Lambda' \subseteq \mathbb{R}^n$ ,  $\emptyset \neq \Lambda \subseteq \mathbb{R}^n$ ,  $F : \Lambda \times X \rightarrow Y$  is a given function,  $\rho$  is a binary relation on  $Y$ , and  $\Lambda' \subseteq \Lambda''$ . Then we say that:

(i)  $F(\cdot; \cdot)$  is Levitan  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \Lambda', \rho, \mathcal{P}_{\mathbf{K}})$ -almost periodic if for every  $B \in \mathcal{B}$ ,  $\epsilon > 0$  and for every nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$ , there exists  $l > 0$  such that for each  $\mathbf{t}_0 \in \Lambda'$  there exists  $\tau \in B(\mathbf{t}_0, l) \cap \Lambda'$  such that, for every  $\mathbf{t} \in \Lambda$  and  $x \in B$ , there exists an element  $y_{\mathbf{t};x} \in \rho(F(\mathbf{t}; x))$  such that  $\mathbb{F}_K(\cdot)[\varphi(\|F(\cdot + \tau; x) - y_{\cdot;x}\|_Y)]|_{K \cap \Lambda} \in P_K$  for all  $x \in X$ , and

$$\sup_{x \in B} \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \|F(\cdot + \tau; x) - y_{\cdot;x}\|_Y \right) \right] \right\|_{|K \cap \Lambda|_{P_K}} \leq \epsilon.$$

(ii)  $F(\cdot; \cdot)$  is Bebutov  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \Lambda', \rho, \mathcal{P}_{\mathbf{K}})$ -uniformly recurrent if for every  $B \in \mathcal{B}$  and for every nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$ , there exists a sequence  $(\tau_k)$  in  $\Lambda'$  such that  $\lim_{k \rightarrow +\infty} |\tau_k| = +\infty$  and that, for every  $\mathbf{t} \in \Lambda$  and  $x \in B$ , there exists an element  $y_{\mathbf{t};x} \in \rho(F(\mathbf{t}; x))$  such that  $\mathbb{F}_K(\cdot)[\varphi(\|F(\cdot + \tau_k; x) - y_{\cdot;x}\|_Y)]|_{K \cap \Lambda} \in P_K$  for all  $x \in X$ , and

$$\lim_{k \rightarrow +\infty} \sup_{x \in B} \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \|F(\cdot + \tau_k; x) - y_{\cdot;x}\|_Y \right) \right] \right\|_{|K \cap \Lambda|_{P_K}} = 0.$$

If the sequence  $(\tau_k)$  in  $\Lambda'$  is independent of the choice of a nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$ , for a set  $B \in \mathcal{B}$  given in advance, then we say that the function  $F(\cdot; \cdot)$  is Bebutov  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \Lambda', \rho, \mathcal{P}_{\mathbf{K}})$ -uniformly recurrent of type 1.

In any normal situation, a Bebutov  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \Lambda', \rho, \mathcal{P}_{\mathbf{K}})$ -uniformly recurrent function is already Bebutov  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \Lambda', \rho, \mathcal{P}_{\mathbf{K}})$ -uniformly recurrent function of type 1:

**Proposition 2.6** Suppose that  $\emptyset \neq \Lambda' \subseteq \mathbb{R}^n$  and  $\emptyset \neq \Lambda \subseteq \mathbb{R}^n$  are unbounded sets,  $F : \Lambda \times X \rightarrow Y$  is a given function,  $\rho$  is a binary relation on  $Y$ , and  $\Lambda' \subseteq \Lambda''$ . If  $F(\cdot; \cdot)$  is Bebutov  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \Lambda', \rho, \mathcal{P}_{\mathbf{K}})$ -uniformly recurrent, then  $F(\cdot; \cdot)$  is Bebutov  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \Lambda', \rho, \mathcal{P}_{\mathbf{K}})$ -uniformly recurrent of type 1, provided that for each compact set  $K \subseteq \mathbb{R}^n$  such that  $K \cap \Lambda \neq \emptyset$ , there exists a finite real constant  $c_K > 0$  such that, for every  $x \in X$ ,  $\tau \in \Lambda'$ , and for every compact set  $K' \subseteq \mathbb{R}^n$  which contains  $K$ , we have

$$\begin{aligned} & \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \|F(\cdot + \tau; x) - y_{\cdot;x}\|_Y \right) \right] \right\|_{|K \cap \Lambda|_{P_K}} \\ & \leq c_K \left\| \mathbb{F}_{K'}(\cdot) \left[ \varphi \left( \|F(\cdot + \tau; x) - y_{\cdot;x}\|_Y \right) \right] \right\|_{|K' \cap \Lambda|_{P_{K'}}}, \quad y_{\cdot;x} \in \rho(F(\cdot; x)). \end{aligned} \tag{2.2}$$

**Proof** Let the set  $B \in \mathcal{B}$  be given. We know that there exists a natural number  $N_0 \in \mathbb{N}$  such that  $[-N, N]^n \cap \Lambda \neq \emptyset$  for every natural number  $N \geq N_0$ . Choose a point  $\tau_N$  in  $\Lambda'$  such that  $|\tau_N| \geq N$  and

$$\sup_{x \in B} \left\| \mathbb{F}_{[-N, N]^n}(\cdot) \left[ \varphi \left( \|F(\cdot + \tau_N; x) - y_{\cdot;x}\|_Y \right) \right] \right\|_{|[-N, N]^n \cap \Lambda|_{P_{[-N, N]^n}}} \leq 1/N.$$

Then the sequence  $(\tau_N)$  in  $\Lambda'$  satisfies the desired requirements since we have assumed (2.2). □



The notion introduced in [14, definitions 2.1, 2.5, and 2.8] is a special case of the notion introduced in the previous three definitions, provided that for each nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$  we have the existence of a finite real constant  $c_K > 0$  such that  $d(f|_{K \cap \Lambda}, g|_{K \cap \Lambda}) \leq c_K d(f, g)$  for all  $f, g \in P$ ,  $\mathbb{F}_K = \mathbb{F}|_{K \cap \Lambda}$ , and  $P_K = \{f|_{K \cap \Lambda}; f \in P\}$ ; here and hereafter,  $\mathcal{P} = (P, d)$  is a pseudometric space and  $P \subseteq [0, \infty)^Y$  contains the zero function.

Usually, we plug  $\mathbb{F}_K(\cdot) \equiv 1$ ,  $\varphi(x) \equiv x$ ,  $\rho = I$  and  $P_K = C(\Lambda \cap K : Y)$  in Definitions 2.4 and 2.5. For some examples of the one-dimensional uniformly Poisson stable functions, we refer the reader to [53] and the research monograph [23, pp. 219-223] by J. de Vries (the function constructed by B. A. Shcherbakov in [53] is uniformly recurrent, in fact, which simply follows from an application of [53, lemma, p. 324]). Any compactly almost automorphic function  $f : \mathbb{R} \rightarrow X$  is bounded, uniformly continuous and Levitan  $N$ -almost periodic, which simply implies that  $F(\cdot)$  is uniformly Poisson stable; on the other hand, we know that there exists a compactly almost automorphic function  $f : \mathbb{R} \rightarrow \mathbb{R}$  which is not (asymptotically) uniformly recurrent (see, e.g., [33, example 2.4.35]). Therefore, the class of compactly almost automorphic functions seems to be ideal for finding certain functions which are uniformly Poisson stable but not uniformly recurrent (the function analyzed in Example 2.3 is uniformly recurrent but not Stepanov almost automorphic; cf. [33] for the notion). Another class of functions which are always uniformly Poisson stable but not necessarily uniformly recurrent can be simply constructed by taking the compositions of the uniformly recurrent functions which do not have relatively compact range with the functions which are only continuous but not uniformly continuous.

**Example 2.7** Let  $c_0$  be the Banach space of all numerical sequences vanishing at infinity, equipped with the sup-norm. Define  $f : [0, \infty) \rightarrow c_0$  by

$$f(t) := \left( \frac{4n^2 t^2}{(t^2 + n^2)^2} \right)_{n \in \mathbb{N}}, \quad t \geq 0.$$

Besides many other features, we know that the function  $f(\cdot)$  is bounded, uniformly continuous, quasiasymptotically almost periodic, and not almost automorphic; see, e.g., [32, example 3.11.14]. Here we will prove that the function  $f(\cdot)$  is not uniformly Poisson stable. Let us assume the contrary, and let

$$0 < \epsilon < \inf_{n \in \mathbb{N}} \frac{4n^4}{(2n^2 + 2n + 1)^2}; \tag{2.3}$$

then there exists a strictly increasing sequence  $(\tau_k)$  of positive real numbers such that  $|f(\tau_k) - f(0)| \leq \epsilon$ ,  $k \geq k_0$  for some positive integer  $k_0 \in \mathbb{N}$ . This implies

$$\sup_{n \in \mathbb{N}} \frac{4n^2 \tau_k^2}{(\tau_k^2 + n^2)^2} \leq \epsilon, \quad k \geq k_0.$$

Let  $\tau_k \in [n_k, n_k + 1)$  for some  $n_k \in \mathbb{N}$  ( $k \geq k_0$ ). Then the previous estimate implies

$$\epsilon \geq \sup_{n \in \mathbb{N}} \frac{4n^2 \tau_k^2}{(\tau_k^2 + n^2)^2} \geq \frac{4n_k^2 n_k^2}{((n_k + 1)^2 + n_k^2)^2} > \epsilon,$$

which is a contradiction; see (2.3).

In the following example, the considered metric space  $P_K$  is different from  $C(\Lambda \cap K : Y)$  :

**Example 2.8** Suppose that  $(T(t)) \subseteq L(X, Y)$  is a strongly continuous operator family and  $\mathcal{B}$  denotes the collection of all bounded subsets of  $X$ . Define

$$F(t, s; x) := e^{\int_s^t \varphi(\tau) d\tau} T(t - s)x, \quad (t, s) \in \mathbb{R}^2, \quad x \in X.$$

Then our analysis from [33, example 8.1.5] shows the following: If the function  $\varphi(\cdot)$  is bounded and Levitan  $(x, \mathbb{R}, 1, \mathcal{P}_K)$ -normal, where  $\mathbb{R}$  denotes the collection of all sequences in  $\mathbb{R}$  and  $P_K = L^1(\mathbb{R})$  for each compact set  $K \subseteq \mathbb{R}$ , then the function  $F(\cdot, \cdot; \cdot)$  is Levitan  $(x, \mathbb{R}_1, \mathcal{B}, 1, \mathcal{P}_{1,K})$ -normal, where  $\mathbb{R}_1$  denotes the collection of all sequences in  $\{(x, x) : x \in \mathbb{R}\}$  and  $P_{1,K} = C(\mathbb{R})$  for each compact set  $K \subseteq \mathbb{R}^2$ .

The usually considered spaces of Levitan  $N$ -almost periodic functions and uniformly Poisson stable functions are translation invariant; the basic properties of multidimensional  $\rho$ -almost periodic functions clarified in [25, proposition 2.11] can be formulated in our new setting, as well. Further on, the qualitative behavior of the limit function of a uniformly convergent sequence of almost periodic type functions has been analyzed many times before. For example, using the same argumentation as in the proof of [14, proposition 2.6], we can deduce the following result:

**Proposition 2.9** Suppose that  $\mathbb{R}$  is any collection of sequences in  $\Lambda'$ ,  $F_j : \Lambda \times X \rightarrow Y$ , and the function  $F_j(\cdot; \cdot)$  is Levitan  $(\varphi, \mathbb{R}, \mathcal{B}, \mathbb{F}_K, \mathcal{P}_K)$ -normal for all  $j \in \mathbb{N}$ . If  $F : \Lambda \times X \rightarrow Y$  and, for every set  $B \in \mathcal{B}$ , for every sequence  $(\mathbf{b}_k)_{k \in \mathbb{N}}$  in  $\mathbb{R}$  and for every nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$ , we have

$$\lim_{(j,k) \rightarrow +\infty} \sup_{x \in B} \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \|F_j(\cdot + \mathbf{b}_j; x) - F(\cdot + \mathbf{b}_k; x)\|_Y \right) \right]_{K \cap \Lambda} \right\|_{P_K} = 0,$$

then the function  $F(\cdot; \cdot)$  is likewise Levitan  $(\varphi, \mathbb{R}, \mathcal{B}, \mathbb{F}_K, \mathcal{P}_K)$ -normal, provided that:

(i) The function  $\varphi(\cdot)$  is monotonically increasing and there exists a finite real constant  $c > 0$  such that  $\varphi(x + y) \leq c[\varphi(x) + \varphi(y)]$  for all  $x, y \geq 0$ .

(ii) Condition (C3-K) holds, where:

(C3-K) For every nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$ , there exists a finite real constant  $f_K > 0$  such that the assumptions  $f, g, h \in P_K$  and  $0 \leq w \leq d'[f + g + h]$  for some finite real constant  $d' > 0$  imply  $w \in P_K$  and  $\|w\|_{P_K} \leq f_K(1 + d')[\|f\|_{P_K} + \|g\|_{P_K} + \|h\|_{P_K}]$ .

Without going into full details, we will only note here that the supremum formula can be formulated in our new framework; for example, if a function  $F : \mathbb{R}^n \rightarrow X$  is uniformly Poisson stable, then for each positive real number  $a > 0$  we have  $\sup_{\mathbf{t} \in \mathbb{R}^n} \|F(\mathbf{t})\| = \sup_{\mathbf{t} \in \mathbb{R}^n; |\mathbf{t}| \geq a} \|F(\mathbf{t})\| \in [0, \infty]$ ; cf. also [33, propositions 2.4.13, 6.1.6, and 8.1.15] for the case that  $\rho = I$ , and [25, proposition 2.13] for the case in which  $\rho = T \in L(Y)$  is a linear isomorphism (the statement of [25, proposition 2.20] concerning multidimensional  $\rho$ -almost periodic functions in the finite-dimensional spaces can be formulated in our new setting, as well).

We continue with some examples:

**Example 2.10** (i) The spaces of Levitan  $N$ -almost periodic functions (Bebutov uniformly recurrent functions)  $F : \mathbb{R}^n \rightarrow Y$  can be constricted if we use the pseudometric spaces  $\mathcal{P}_K$  such that  $\mathcal{P}_K$  is continuously embedded into the space  $C(K : Y)$  for every nonempty compact set  $K \subseteq \mathbb{R}^n$ . For instance, the function  $f(t) := 1/(2 + \cos t + \cos(\sqrt{2}t))$ ,  $t \in \mathbb{R}$  is unbounded, continuous and Levitan  $(x, 1, \mathcal{P}_K)$ -almost periodic, where  $\mathcal{P}_K = C(K)$ ; see [41, pp. 58-59] for more details. Therefore, the function  $f(\cdot)$  is uniformly Poisson stable, as well; before going any further, we would like to ask whether the function  $f(\cdot)$  is uniformly recurrent or Besicovitch  $p$ -bounded for some finite exponent  $p \geq 1$  (cf. [33] for the notion)?

Here, we can also use the metric spaces  $\mathcal{P}_K$  equipped with the distance of the form

$$d(f|_K, g|_K) := \sup_{\mathbf{t} \in K} \|f(\mathbf{t}) - g(\mathbf{t})\| + d_1(f|_K, g|_K), \quad f, g \in P,$$

where  $P$  is a subspace of the space  $C_b(\mathbb{R}^n : [0, \infty))$ ,  $d_1(\cdot; \cdot)$  is a pseudometric on  $P_K$ ,  $\mathbb{F}_K = \mathbb{F}_{|K \cap \Lambda}$  and  $P_K = \{f|_{K \cap \Lambda}; f \in P\}$ .

(ii) As already emphasized in [14, example 2.4(iii)], the spaces of Levitan  $N$ -almost periodic functions (Bebutov uniformly recurrent functions)  $F : \mathbb{R}^n \rightarrow Y$  can be extended if we use the pseudometric spaces  $\mathcal{P}_K$  such that the space  $C(K : Y)$  is continuously embedded into  $\mathcal{P}_K$  for every nonempty compact set  $K \subseteq \mathbb{R}^n$ . The use of the incomplete metric space  $P$  consisting of all continuous functions from  $\mathbb{R}^n$  into  $[0, \infty)$ , equipped with the distance

$$d(f, g) := \sup_{x \in \mathbb{R}^n} |\arctan(f(x)) - \arctan(g(x))|, \quad f, g \in P,$$

has been proposed in [14]. This concept allows one to consider the generalized almost periodicity of functions  $f(\cdot)$  which are not locally integrable. We can similarly analyze the generalized Levitan  $N$ -almost periodicity and the generalized Bebutov uniform recurrence of the functions  $f(\cdot)$  which are not locally integrable by replacing the term  $\|\cdot\|_{P_K}$  by  $\|\cdot\|_{\overline{P_K}}$ , where  $\overline{P_K} = (\overline{P_K}, \overline{d_K})$  is the completion of the metric space  $\mathcal{P}_K$ .

(iii) Define the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  by  $f(x) := n3^{n+1} \sin(2\pi x)$  if  $x \in [3^n, 3^n + 1] + 2 \cdot 3^{n+1}\mathbb{Z}$  for some  $n \in \mathbb{N}$ , and  $f(x) := 0$ , otherwise. Then  $f(\cdot)$  is clearly unbounded; moreover, we know that  $f(\cdot)$  is Levitan  $N$ -almost periodic (see A. Nawrocki [45, lemma 2.18, example 2.19]).

(iv) Suppose that  $(\alpha_k)$  is a fixed sequence of positive real numbers such that  $\lim_{k \rightarrow +\infty} \alpha_k = +\infty$ . If  $f_i : \mathbb{R} \rightarrow \mathbb{C}$  is a bounded, uniformly Poisson stable function such that  $\lim_{k \rightarrow +\infty} f_i(\cdot + \alpha_k) = f_i(\cdot)$ , uniformly on compacts of  $\mathbb{R}$  ( $1 \leq i \leq n$ ), then the function  $F(\cdot)$ , given by

$$F(t_1, \dots, t_n) := f_1(t_1) \cdot \dots \cdot f_n(t_n), \quad \mathbf{t} = (t_1, \dots, t_n) \in \mathbb{R}^n,$$

is bounded, uniformly Poisson stable and satisfies  $\lim_{k \rightarrow +\infty} F(\cdot + \beta_k) = F(\cdot)$ , uniformly on compacts of  $\mathbb{R}^n$ , where  $\beta_k = (\alpha_k, \dots, \alpha_k)$  for all  $k \in \mathbb{N}$ .

(v) The introduced notion has the meaning even if  $\rho = 0 \in L(Y)$ . In this case, we can simply construct a Bebutov  $(x, 1, 0, \mathcal{P}_K)$ -uniformly recurrent function  $f : \mathbb{R} \rightarrow \mathbb{R}$  which does not vanish as  $|t| \rightarrow +\infty$ ; here,  $P_K = C(K)$  for each nonempty compact subset of  $\mathbb{R}$ .

(vi) In order to avoid any form of plagiarism, we will only note here that examples 6.1.13, 6.1.15, and 6.1.16 of [33] can be formulated in our new framework; these examples justify the introduction of our concepts with the set  $\Lambda'$  being not equal to the set  $\Lambda$  or some of its proper subsets.

In the usually considered situation, the notion introduced in Definition 2.1 is more specific than the notion introduced in Definitions 2.4 and 2.5. Concerning this issue, we will clarify the following result (cf. also [14, proposition 2.9]):

**Proposition 2.11** (i) Suppose that  $R$  is any collection of sequences in  $\Lambda''$ ,  $F : \Lambda \times X \rightarrow Y$ , and  $\varphi : [0, \infty) \rightarrow [0, \infty)$ . Let the following conditions hold:

- (a) The function  $\varphi(\cdot)$  is monotonically increasing, continuous at the point zero, and there exists a finite real constant  $c > 0$  such that  $\varphi(x + y) \leq c[\varphi(x) + \varphi(y)]$  for all  $x, y \geq 0$ .
- (b) Condition (C3-K) holds.
- (c) For every nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$ , we have that  $\mathbb{F}_K(\cdot)\varphi(\|P(\cdot; x)\|_Y) \in P_K$  for any trigonometric polynomial (periodic function)  $P(\cdot; \cdot)$  and  $x \in X$ .
- (d) For every nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$ , there exists a finite real constant  $g_K > 0$  such that

$$\left\| \mathbb{F}_K(\cdot) \left[ \varphi(\|P(\cdot; x)\|_Y) \right] \Big|_{K \cap \Lambda} \right\|_{P_K} \leq g_K \sup_{t \in K \cap \Lambda} \varphi(\|P(t; x)\|_Y),$$

for any any trigonometric polynomial (periodic function)  $P(\cdot; \cdot)$  and  $x \in X$ .

- (e) For every nonempty compact set  $K \subseteq \mathbb{R}^n$  with  $K \cap \Lambda \neq \emptyset$ , there exists a finite real constant  $h_K > 0$  such that, for every  $x \in X$  and  $\tau \in \Lambda''$ , the assumptions  $H : \Lambda \times X \rightarrow Y$  and  $\mathbb{F}_K(\cdot)[\varphi(\|H(\cdot; x)\|_Y)] \Big|_{K \cap \Lambda} \in P_K$  imply  $\mathbb{F}_K(\cdot)[\varphi(\|H(\cdot + \tau; x)\|_Y)] \Big|_{K \cap \Lambda} \in P_K$  and

$$\left\| \mathbb{F}_K(\cdot) \left[ \varphi(\|H(\cdot + \tau; x)\|_Y) \right] \Big|_{K \cap \Lambda} \right\|_{P_K} \leq h_K \left\| \mathbb{F}_K(\cdot) \left[ \varphi(\|H(\cdot; x)\|_Y) \right] \Big|_{K \cap \Lambda} \right\|_{P_K}.$$

- (f) Any set  $B$  of collection  $\mathcal{B}$  is bounded.

If the function  $F(\cdot; \cdot)$  is Levitan strongly  $(\varphi, \mathbb{F}_K, \mathcal{B}, \mathcal{P}_K)$ -almost periodic (Levitan semi- $(\varphi, I, \mathbb{F}_K, \mathcal{B}, \mathcal{P}_K)_{j \in \mathbb{N}_n}$ -periodic), then the function  $F(\cdot; \cdot)$  is Levitan  $(\varphi, R, \mathcal{B}, \varphi, \mathbb{F}_K, \mathcal{P}_K)$ -normal.

- (ii) Suppose that  $\emptyset \neq \Lambda' \subseteq \mathbb{R}^n$ ,  $\emptyset \neq \Lambda \subseteq \mathbb{R}^n$ ,  $F : \Lambda \times X \rightarrow Y$  is a given function and  $\Lambda' \subseteq \Lambda''$ . If the function  $F(\cdot; \cdot)$  is Levitan strongly  $(\varphi, \mathbb{F}_K, \mathcal{B}, \mathcal{P}_K)$ -almost periodic (Levitan semi- $(\varphi, I, \mathbb{F}_K, \mathcal{B}, \mathcal{P}_K)_{j \in \mathbb{N}_n}$ -periodic), then the function  $F(\cdot; \cdot)$  is Levitan  $(\varphi, \mathbb{F}_K, \mathcal{B}, \Lambda', I, \mathcal{P}_K)$ -almost periodic, provided that the assumptions (a)-(f) given in the formulation of (i) hold.

**Proof** We will include all details of the proof of (i) for the sake of completeness, considering the class of Levitan strongly  $(\varphi, \mathbb{F}_K, \mathcal{B}, \mathcal{P}_K)$ -almost periodic functions, only. Let  $F(\cdot; \cdot)$  be such a function, let  $\epsilon > 0$ ,  $B \in \mathcal{B}$ , and let  $(\mathbf{b}_k)_{k \in \mathbb{N}}$  belong to  $R$ . Suppose, further, that  $K$  is a nonempty compact set of  $\mathbb{R}^n$  such that  $K \cap \Lambda \neq \emptyset$ . Then there exists a trigonometric polynomial  $P_k^{B,K}(\cdot; \cdot)$  such that  $\mathbb{F}_K(\cdot)[\varphi(\|P_k^{B,K}(\cdot; x) - F(\cdot; x)\|_Y)] \Big|_{K \cap \Lambda} \in P_K$

for all  $x \in X$ , and

$$\sup_{x \in B} \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \| P_k^{B,K}(\cdot; x) - F(\cdot; x) \|_Y \right) \right]_{|K \cap \Lambda} \right\|_{P_K} < \epsilon/3.$$

Using (e), we get:

$$\sup_{x \in B} \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \| P_k^{B,K}(\cdot + \tau; x) - F(\cdot + \tau; x) \|_Y \right) \right]_{|K \cap \Lambda} \right\|_{P_K} < h_K \epsilon/3 \tag{2.4}$$

for all  $\tau \in \Lambda''$ . The set  $B$  is bounded due to (f), so that the Bochner criterion for almost periodic functions in  $\mathbb{R}^n$  ensures that there exist a subsequence  $(\mathbf{b}_{k_m})_{m \in \mathbb{N}}$  of  $(\mathbf{b}_k)_{k \in \mathbb{N}}$  and a natural number  $m_0 \in \mathbb{N}$  such that, for every positive integers  $m', m'' \geq m_0$ , we have

$$\sup_{x \in B} \left\| P_k^{B,K}(\mathbf{t} + \mathbf{b}_{k_{m'}}; x) - P_k^{B,K}(\mathbf{t} + \mathbf{b}_{k_{m''}}; x) \right\|_Y < \epsilon/3, \quad \mathbf{t} \in \mathbb{R}^n.$$

Using (c)-(d), we obtain:

$$\sup_{x \in B} \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \| P_k^{B,K}(\cdot + \mathbf{b}_{k_{m'}}; x) - P_k^{B,K}(\cdot + \mathbf{b}_{k_{m''}}; x) \|_Y \right) \right]_{|K \cap \Lambda} \right\|_{P_K} \leq g_K \varphi(\epsilon/3),$$

for all positive integers  $m', m'' \geq m_0$ . Then the final conclusion follows from conditions (a)-(b), the estimate (2.4) and the following decomposition:

$$\begin{aligned} & \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \| F(\cdot + \mathbf{b}_{k_{m'}}; x) - F(\cdot + \mathbf{b}_{k_{m''}}; x) \|_Y \right) \right]_{|K \cap \Lambda} \right\|_{P_K} \\ & \leq c'_K \left[ \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \| F(\cdot + \mathbf{b}_{k_{m'}}; x) - P(\cdot + \mathbf{b}_{k_{m'}}; x) \|_Y \right) \right]_{|K \cap \Lambda} \right\|_{P_K} \right. \\ & \quad + \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \| P(\cdot + \mathbf{b}_{k_{m'}}; x) - P(\cdot + \mathbf{b}_{k_{m''}}; x) \|_Y \right) \right]_{|K \cap \Lambda} \right\|_{P_K} \\ & \quad \left. + \left\| \mathbb{F}_K(\cdot) \left[ \varphi \left( \| P(\cdot + \mathbf{b}_{k_{m''}}; x) - F(\cdot + \mathbf{b}_{k_{m''}}; x) \|_Y \right) \right]_{|K \cap \Lambda} \right\|_{P_K} \right], \end{aligned}$$

holding with a certain positive real constant  $c'_K > 0$ . □

The method proposed in the proof of [33, theorem 6.1.37] does not work for Levitan  $N$ -almost periodic functions and uniformly Poisson stable functions and, because of that, we will omit here all details regarding the extensions of Levitan  $N$ -almost periodic functions and the extensions of uniformly Poisson stable functions. Further on, the statement of [14, proposition 2.10] can be formulated in our new framework under certain very restrictive assumptions. Before proceeding further, we will also note that the class of metrical semi- $(c_j, \mathcal{B})_{j \in \mathbb{N}_n}$ -periodic functions has been analyzed in [14, subsection 2.2] following the initial analysis of J. Andres and D. Pennequin in [7]. The Levitan and Bebutov classes of metrical semi- $(c_j, \mathcal{B})_{j \in \mathbb{N}_n}$ -periodic functions can be considered similarly; we leave all details concerning these topics to the interested readers.

**2.1. Levitan  $(N, c)$ -almost periodic functions and uniformly Poisson  $c$ -stable functions**

In this subsection, we will provide some examples and open questions concerning the Levitan  $(N, c)$ -almost periodic functions and the uniformly Poisson  $c$ -stable functions, where  $c \in \mathbb{C} \setminus \{0\}$ ; for simplicity, we will always consider here the case in which  $\Lambda = \mathbb{R}^n$ ,  $\varphi(x) \equiv x$ ,  $\mathbb{F}_{\mathbf{K}} \equiv 1$ , and  $P_K = C(K)$  for each nonempty compact set  $K \subseteq \mathbb{R}^n$ . Let  $\rho = cI$ ; then any Levitan  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \Lambda', \rho, \mathcal{P}_{\mathbf{K}})$ -almost periodic function is simply called Levitan  $(N, \Lambda', c)$ -almost periodic and any Bebutov  $(\varphi, \mathbb{F}_{\mathbf{K}}, \mathcal{B}, \Lambda', \rho, \mathcal{P}_{\mathbf{K}})$ -uniformly recurrent function is simply called uniformly Poisson  $(\Lambda', c)$ -stable [we omit the term “ $\Lambda'$ ” from the notation if  $\Lambda' = \mathbb{R}^n$ ].

The proofs of [33, propositions 4.2.14 and 7.1.13] do not work for the Levitan/Bebutov concepts; because of that, we would like to ask the following:

**Problem.** Suppose that  $c \in \mathbb{C} \setminus \{0, 1\}$ . Can we find a bounded continuous function  $F : \mathbb{R}^n \rightarrow \mathbb{C}$  which is Levitan  $(N, c)$ -almost periodic (uniformly  $c$ -Poisson stable) but not Levitan  $(N, c^2)$ -almost periodic (uniformly  $c^2$ -Poisson stable)?

The following important counterexample shows that there exists a nontrivial uniformly Poisson  $c$ -stable function  $F : \mathbb{R} \rightarrow \mathbb{R}$  for any complex number  $c \neq 0$ ; this counterexample is based on the conclusions established in [21, lemma 3.5] and particularly shows that the requirements of [33, proposition 7.1.9] do not imply  $c = \pm 1$  for uniformly Poisson  $c$ -stable functions:

**Example 2.12** Consider the function  $\varphi : \mathbb{R} \rightarrow [0, \infty)$  given by

$$\varphi(t) := \sum_{k=1}^{\infty} \sin^2\left(\frac{\pi t}{2^k}\right), \quad t \in \mathbb{R}.$$

In the above-mentioned lemma, E. Ait Dads, B. Es-sebbar, and L. Lhachimi have proved that the function  $\varphi(\cdot)$  is continuous and

$$\lim_{l \rightarrow +\infty} \varphi(t + 2^l) = \lim_{l \rightarrow +\infty} \varphi(t - 2^l) = \varphi(t) + \varphi(1), \tag{2.5}$$

pointwisely on  $\mathbb{R}$ . We will first prove that the function  $\varphi(\cdot)$  is Lipschitz continuous as well as that the convergence in (2.5) is uniform on compact subsets of  $\mathbb{R}$ . The uniform continuity of function  $\varphi(\cdot)$  simply follows from the fact that the function  $\sin^2 \cdot$  is Lipschitz with the corresponding Lipschitz constant  $L = 1$  and the Lagrange mean value theorem, which shows that  $|\varphi(x) - \varphi(y)| \leq \pi|x - y|$ ,  $x, y \in \mathbb{R}$ . Let  $K = [a, b] \subseteq \mathbb{R}$  be a compact set. Then we have:

$$\varphi(t + 2^l) = \sum_{k=1}^l \sin^2\left(\frac{\pi t}{2^k}\right) + \sum_{k=1}^{\infty} \sin^2\left(\frac{\pi t}{2^{k+l}} + \frac{\pi}{2^k}\right), \quad t \in \mathbb{R}.$$

Using this equality, the above-mentioned fact that the function  $\sin^2 \cdot$  is Lipschitz continuous with the corresponding Lipschitz constant  $L = 1$  and the Langrange mean value theorem, we simply get that the convergence in (2.5) is uniform in  $t \in K$  since:

$$\sum_{k=l+1}^{\infty} \sin^2\left(\frac{\pi t}{2^k}\right) \leq \sum_{k=l+1}^{\infty} \frac{\pi^2(|a| + |b|)^2}{4^k}, \quad k, l \in \mathbb{N}$$

and

$$\left| \sum_{k=1}^{\infty} \sin^2\left(\frac{\pi t}{2^{k+l}} + \frac{\pi}{2^k}\right) - \sum_{k=1}^{\infty} \sin^2\left(\frac{\pi}{2^k}\right) \right| \leq \pi(|a| + |b|) \sum_{k=l+1}^{\infty} 2^{-k}, \quad k, l \in \mathbb{N};$$

here we have also used the elementary inequality  $|\sin t| \leq |t|$ ,  $t \in \mathbb{R}$ . Therefore, we have that the function  $\varphi(\cdot)$  is Bebutov  $(x, 1, \{2^l : l \in \mathbb{N}\}, \rho, \mathcal{P}_K)$ -uniformly recurrent, where  $P_K = C(K)$  for each nonempty compact subset  $K$  of  $\mathbb{R}$ ,  $D(\rho) := [0, \infty)$  and  $\rho(\varphi(t)) := \varphi(t) + \varphi(1)$  for all  $t \in \mathbb{R}$  (note that the function  $\varphi(\cdot)$  is surjective since it is continuous,  $\varphi(0) = 0$  and the function  $\varphi(\cdot)$  is Besicovitch unbounded, which follows from the inequality  $\varphi(t) \geq f(\pi t)$ ,  $t \in \mathbb{R}$  with the function  $f(\cdot)$  being defined through (2.1)).

Suppose now that  $c \in \mathbb{C} \setminus \{0\}$  is fixed and consider any nontrivial continuous  $(\varphi(1), c)$ -periodic function  $p : \mathbb{R} \rightarrow \mathbb{C}$ , i.e. any nontrivial continuous function  $p : \mathbb{R} \rightarrow \mathbb{C}$  such that  $p(t + \varphi(1)) = cp(t)$  for all  $t \in \mathbb{R}$ . Then an elementary argumentation involving the equation (2.5) and the uniform continuity of function  $p(\cdot)$  on the interval  $[-a, a]$ , where  $a = \max_{t \in K}(\varphi(t) + \varphi(1) + 1)$ , shows that the function  $(p \circ \varphi)(\cdot)$  is uniformly Poisson  $c$ -stable, nontrivial and  $\lim_{l \rightarrow +\infty} (p \circ \varphi)(\cdot + 2^l) = (p \circ \varphi)(\cdot)$ , uniformly on compacts of  $\mathbb{R}$ .

Let us finally consider the case  $c = 1$  and the function  $p(t) := \sin(2\pi\varphi(t)/\varphi(1))$ ,  $t \in \mathbb{R}$ . Then  $p(\cdot)$  is of period  $\varphi(1)$ , and we know that the function  $(p \circ \varphi)(\cdot)$  is uniformly Poisson stable. At the end of this example, we would like to ask whether the function  $(p \circ \varphi)(\cdot)$  is uniformly recurrent (cf. also [21, proposition 3.6])?

Before proceeding to the next subsection, we would also like to ask whether we can construct a continuous Levitan  $(N, c)$ -almost periodic function  $f : \mathbb{R} \rightarrow \mathbb{R}$  for any complex number  $c \neq 0$ ?

### 2.2. Multidimensional Levitan $N$ -almost periodic functions

Let us recall that a continuous function  $F : \mathbb{R}^n \rightarrow X$  is almost periodic if for each finite real number  $\epsilon > 0$  there exist relatively dense sets  $E^j$  in  $\mathbb{R}$  ( $1 \leq j \leq n$ ) such that the set  $E \equiv \prod_{j=1}^n E^j$  is contained in the set of all  $\epsilon$ -almost periods of  $F(\cdot)$ ; see, e.g., [33, introduction, pp. 31–32]. Motivated by this fact, we will first introduce the following notion (let us only note here that we can similarly reconsider the notion of pre-Levitan almost periodicity):

**Definition 2.13** Suppose that  $F : \mathbb{R}^n \rightarrow X$  is a continuous function. Then we say that the function  $F(\cdot)$  is strongly Levitan  $N$ -almost periodic if  $F(\cdot)$  is pre-Levitan almost periodic and, for every real numbers  $N > 0$  and  $\epsilon > 0$ , there exist a finite real number  $\eta > 0$  and the relatively dense sets  $E_{\eta;N}^j$  in  $\mathbb{R}$  ( $1 \leq j \leq n$ ) such that the set  $E_{\eta;N} \equiv \prod_{j=1}^n E_{\eta;N}^j$  consists solely of  $(\eta, N)$ -almost periods of  $F(\cdot)$  and  $E_{\eta;N} \pm E_{\eta;N} \subseteq E(\epsilon, N)$ .

It is clear that any strongly Levitan  $N$ -almost periodic function  $F : \mathbb{R}^n \rightarrow X$  is Levitan  $N$ -almost periodic as well as that both notions coincide in the one-dimensional setting.

We continue this section by introducing the following notion, which has been proposed for the first time by V. A. Marchenko in [44] (1950, the one-dimensional setting):

**Definition 2.14** Suppose that  $F : \mathbb{R}^n \rightarrow X$  is a continuous function. Then we say that  $F(\cdot)$  is Levitan  $N$ -almost periodic of type 1 if for each  $j \in \mathbb{N}_n$  there exists a real sequence  $(\lambda_k^j)$  such that, for every  $N > 0$  and  $\epsilon > 0$ , there exist numbers  $k \in \mathbb{N}$  and  $\delta > 0$  such that any point  $\tau = (\tau_1, \tau_2, \dots, \tau_n) \in \mathbb{R}^n$  satisfying  $|\lambda_l^j \tau_j| \leq \delta \pmod{2\pi}$  for all  $l \in \mathbb{N}_k$  and  $j \in \mathbb{N}_n$  is an  $\epsilon, N$ -almost period of the function  $F(\cdot)$ .

Now we will clarify the following result:

**Proposition 2.15** *Consider the following statements:*

(i)  $F(\cdot)$  is Levitan  $N$ -almost periodic of type 1.

(ii)  $F(\cdot)$  is strongly Levitan  $N$ -almost periodic.

(iii)  $F(\cdot)$  is Levitan  $N$ -almost periodic.

Then (i)  $\Leftrightarrow$  (ii) and (ii)  $\Rightarrow$  (iii).

**Proof** Using Kronecker’s theorem, the Bogolyubov theorem and the argumentation given in [40, pp. 54-55, 57] we can simply prove, as in the one-dimensional setting, that the statements (i) and (ii) are equivalent. The statement (ii)  $\Rightarrow$  (iii) is trivial.  $\square$

Further on, we can simply show that the Levitan  $N$ -almost periodic functions of type 1 form a vector space with the usual operations. Due to Proposition 2.15, it follows that the strongly Levitan  $N$ -almost periodic form a vector space with the usual operations. Concerning this result, it is logical to ask whether the Levitan  $N$ -almost periodic functions form a vector space with the usual operations as well as whether the all statements from the formulation of Theorem 2.15 are mutually equivalent, i.e. whether the implication (iii)  $\Rightarrow$  (i) is true? Concerning the last question, we would like to notice that the arguments contained on p. 57 of [40] can be repeated once more in order to see that this is true, provided that the following generalization of the Bogolyubov theorem holds true in the higher-dimensional setting:

(S) Suppose that  $A$  is a relatively dense set in  $\mathbb{R}^n$  and  $\delta > 0$ . Then there exist a real number  $\eta > 0$  and a finite set  $\{\omega_1, \dots, \omega_k\}$  in  $\mathbb{R}^n$  such that

$$\left\{ \tau \in \mathbb{R}^n : |e^{i\omega_l \tau} - 1| < \eta \text{ for all } l = 1, \dots, k \right\} \subseteq A - A + A - A + B_\delta.$$

Unfortunately, the statement (S) does not hold in the case that  $n \geq 2$ , as the following trivial counterexample shows:

**Example 2.16** *Suppose that  $A = \mathbb{Z}^n$ ,  $n \geq 2$ , and  $\delta \in (0, 1/4)$ ; then  $A$  is relatively dense in  $\mathbb{R}^n$ . If (S) is true, then there exist a sufficiently small number  $\eta_0 > 0$  and two real numbers  $\omega_1$  and  $\omega_2$  (take  $k = 1$  in (S)) such that  $W = \{\tau \in \mathbb{R}^n : |\omega_1 \tau_1 + \dots + \omega_n \tau_n| \leq \eta_0 \pmod{2\pi}\} \subseteq A - A + A - A + B_\delta = A + B_\delta$ . If  $\omega_1 = \omega_2 = 0$ , then  $W = \mathbb{R}^n$  and the contradiction is obvious; otherwise,  $W$  is a hyperplane in  $\mathbb{R}^n$  and this clearly implies that  $W$  cannot be contained in  $A + B_\delta = (A - A) + \dots + (A - A) + B_\delta$ , where we have exactly  $l$  addends in the above sum with the number  $l \in \mathbb{N} \setminus \{1\}$  being arbitrarily chosen.*

Therefore, it is also important to ask whether the Bogolyubov theorem can be satisfactorily formulated in the higher-dimensional setting?

We close this subsection with the observation that we will not reconsider the approximation theorem for multidimensional Levitan  $N$ -almost periodic functions and the uniqueness theorem for multidimensional Levitan  $N$ -almost periodic functions here; see [41, pp. 60–62] for the corresponding results established in the one-dimensional setting.



### 3. Applications to the abstract Volterra integro-differential equations

In this section, we will present several new theoretical results and their applications to the abstract Volterra integro-differential equations and the classical partial differential equations. We start with the following theme:

#### 3.1. The actions of infinite convolution products

In the one-dimensional setting, the analysis of existence and uniqueness of almost periodic type solutions for various classes of the abstract Volterra integro-differential equations without initial conditions leans heavily on the analysis of the qualitative features of the infinite convolution product

$$t \mapsto F(t) := \int_{-\infty}^t R(t-s)f(s) ds, \quad t \in \mathbb{R}. \tag{3.1}$$

In [1]-[4], M. Akhmet et al. have proposed a new method in the analysis of the existence and uniqueness of unpredictable solutions of linear differential and discrete equations which is called the method of included intervals (see [5]-[6]). The proof of the following result is almost entirely based on the method of included intervals (cf. also [51]-[52], where B. A. Shcherbakov have developed the so-called comparability method by character of recurrence, which is not so easy for comprehension and applications):

**Proposition 3.1** *Let  $\emptyset \neq \Lambda' \subseteq \mathbb{R}$ , let  $A$  be a closed linear operator on  $X$ , and let  $(R(t))_{t>0} \subseteq L(X)$  be a strongly continuous operator family such that  $R(t)A \subseteq AR(t)$  for all  $t > 0$  and  $\int_0^\infty \|R(t)\| dt < \infty$ . Let  $\mathcal{P}_K = C(K)$  for each nonempty compact set  $K$  of  $\mathbb{R}$ . If  $f : \mathbb{R} \rightarrow X$  is a bounded, continuous and Levitan  $(x, \Lambda', A, \mathcal{P}_K)$ -almost periodic, resp. bounded, continuous and Bebutov  $(x, \Lambda', A, \mathcal{P}_K)$ -uniformly recurrent, and the function  $Af : \mathbb{R} \rightarrow X$  is well defined and bounded, then the function  $F : \mathbb{R} \rightarrow X$ , given by (3.1), is bounded, continuous and Levitan  $(x, \Lambda', A, \mathcal{P}_K)$ -almost periodic, resp. bounded, continuous and Bebutov  $(x, \Lambda', A, \mathcal{P}_K)$ -uniformly recurrent.*

**Proof** We will consider only Bebutov  $(x, \Lambda', A, \mathcal{P}_K)$ -uniformly recurrent functions. It is clear that the function  $F(\cdot)$  is well-defined and bounded since

$$F(t) := \int_0^\infty R(s)f(t-s) ds, \quad t \in \mathbb{R},$$

as well as  $\int_0^\infty \|R(t)\| dt < \infty$  and the function  $f(\cdot)$  is bounded. The continuity of  $F(\cdot)$  simply follows from the dominated convergence theorem. Since  $R(t)A \subseteq AR(t)$  for all  $t > 0$  and the function  $Af(\cdot)$  is bounded, we have that  $AF(t) = \int_0^\infty R(s)Af(t-s) ds$  for all  $t \in \mathbb{R}$ . Let  $\epsilon > 0$ , let  $-\infty < a < b < +\infty$ , and let  $K = [a, b]$ . Further on, let  $-\infty < c < a$  be such that

$$(\|f\|_\infty + \|Af\|_\infty) \cdot \int_{a-c}^\infty \|R(s)\| ds < \epsilon/2. \tag{3.2}$$

Then there exists a sequence  $(\tau_k)$  in  $\Lambda'$  such that  $\lim_{k \rightarrow +\infty} |\tau_k| = +\infty$  and

$$\left( \int_0^\infty \|R(t)\| dt \right) \cdot \sup_{t \in [c, b]} \|f(t + \tau_k) - Af(t)\| < \epsilon/2. \tag{3.3}$$

The final conclusion simply follows from the estimates (3.2)-(3.3) and the next computation, which holds for every  $k \in \mathbb{N}$  and  $t \in K$  :

$$\begin{aligned} \|F(t + \tau_k) - AF(t)\| &= \left\| \int_{-\infty}^t R(t-s)[f(s + \tau_k) - Af(s)] ds \right\| \\ &\leq \int_{-\infty}^c \|R(t-s)\| \cdot \|f(s + \tau_k) - Af(s)\| ds + \int_c^t \|R(t-s)\| \cdot \|f(s + \tau_k) - Af(s)\| ds \\ &\leq (\|f\|_\infty + \|Af\|_\infty) \cdot \int_{a-c}^\infty \|R(s)\| ds \\ &+ \left( \int_t^c \|R(t-s)\| ds \right) \cdot \sup_{s \in [c,b]} \|f(s + \tau_k) - Af(s)\| \leq (\epsilon/2) + (\epsilon/2) = \epsilon. \end{aligned}$$

□

**Remark 3.2** (i) Suppose that  $\nu_K : K \rightarrow (0, \infty)$  satisfies that the function  $1/\nu_K(\cdot)$  is bounded. Then the use of weighted function space  $C_{b,\nu_K}(K)$  seems to be discutable here since the argumentation contained in the proof given above indicates that the function  $\nu_K(\cdot)$  has to be bounded, as well, unless we assume certain very restrictive conditions.

(ii) The statement of Proposition 3.1 can be transferred to the multidimensional setting without any serious difficulty; cf. [33] for more details.

(iii) For simplicity, let us consider here the class of Levitan  $(x, \mathbb{R}, I, \mathcal{P}_K)$ -almost periodic functions. If we additionally suppose that for each  $N > 0$  and  $\epsilon > 0$ , there exists a relatively dense set  $E_{\eta,N}$  of  $(\eta, N)$ -almost periods of  $f(\cdot)$  such that  $E_{\eta,N} \pm E_{\eta,N} \subseteq E(\epsilon, N)$ , then the same property holds for the function  $F(\cdot)$ .

As mentioned many times before, our results about the convolution invariance of generalized metrical Levitan almost periodicity and generalized metrical Bebutov uniform recurrence under the actions of the infinite convolution product (3.1) can be successfully applied in the qualitative analysis of solutions for a large class of the abstract (degenerate) Volterra integro-differential equations without initial conditions. For example, Proposition 3.1 is clearly applicable in the analysis of the existence and uniqueness of Bebutov uniformly recurrent solutions of the initial value problems with constant coefficients

$$D_{t,+}^\gamma u(t, x) = \sum_{|\alpha| \leq k} a_\alpha f^{(\alpha)}(t, x) + f(t, x), \quad t \in \mathbb{R}, \quad x \in \mathbb{R}^n$$

in the space  $L^p(\mathbb{R}^n)$ , where  $\gamma \in (0, 1)$ ,  $D_{t,+}^\gamma u(t)$  denotes the Weyl-Liouville fractional derivative of order  $\gamma$  and  $1 \leq p < \infty$ ; cf. also [13, 31] for many similar applications of this type.

### 3.2. The convolution invariance and some applications

In this subsection, we continue our investigation from the previous subsection by examining the convolution invariance of Levitan almost like periodicity. In actual fact, we would like to observe that the method of included intervals [1]-[4] enables one to simply deduce the following result:

**Proposition 3.3** *Let  $\emptyset \neq \Lambda' \subseteq \mathbb{R}^n$ , let  $A$  be a closed linear operator on  $X$ ,  $h \in L^1(\mathbb{R}^n)$ , and let  $\mathcal{P}_K = C(K)$  for each nonempty compact set  $K$  of  $\mathbb{R}$ . If  $f : \mathbb{R}^n \rightarrow X$  is bounded, continuous and Levitan  $(x, \Lambda', A, \mathcal{P}_K)$ -almost periodic, resp. bounded, continuous and Bebutov  $(x, \Lambda', A, \mathcal{P}_K)$ -uniformly recurrent, and the function  $Af : \mathbb{R}^n \rightarrow X$  is well defined and bounded, then the function  $H : \mathbb{R}^n \rightarrow X$ , given by*

$$H(\mathbf{t}) := (h * f)(\mathbf{t}) := \int_{\mathbb{R}^n} h(\mathbf{t} - \mathbf{s})f(\mathbf{s}) \, d\mathbf{s}, \quad \mathbf{t} \in \mathbb{R}^n,$$

*is bounded, continuous and Levitan  $(x, \Lambda', A, \mathcal{P}_K)$ -almost periodic, resp. bounded, continuous and Bebutov  $(x, \Lambda', A, \mathcal{P}_K)$ -uniformly recurrent.*

**Remark 3.4** (i) *Unfortunately, we cannot formulate a satisfactory analogue of Proposition 3.3 in the case that there exist two finite real numbers  $b > 0$  and  $c > 0$  such that  $\|f(\mathbf{t})\| \leq c(1 + |\mathbf{t}|)^b$ ,  $\mathbf{t} \in \mathbb{R}^n$  and  $h(\cdot)[1 + |\cdot|]^b \in L^1(\mathbb{R}^n)$ , when the function  $H(\cdot)$  is clearly well-defined.*

(ii) *Let us consider the class of Levitan  $(x, \mathbb{R}, I, \mathcal{P}_K)$ -almost periodic functions. If we suppose, additionally, that for each  $N > 0$  and  $\epsilon > 0$  there exists a relatively dense set  $E_{\eta;N}$  of  $(\eta, N)$ -almost periods of  $f(\cdot)$  such that  $E_{\eta;N} \pm E_{\eta;N} \subseteq E(\epsilon, N)$ , then the same property holds for the function  $H(\cdot)$ . Taken together with the statement of Proposition 3.3, this provides a proper extension of the well known result [11, theorem 13], proved for the first time by D. Bugajewski, X. Gan, and P. Kasprzak.*

Now we will briefly explain how we can incorporate Proposition 3.3 in the analysis of the existence and uniqueness of the Levitan almost periodic type solutions for some classes of (fractional) partial differential equations:

1. We will consider first the inhomogeneous heat equation in  $\mathbb{R}^n$ . Recall, a unique solution of the heat equation  $u_t(t, x) = u_{xx}(t, x)$ ,  $t \geq 0$ ,  $x \in \mathbb{R}^n$ ;  $u(0, x) = F(x)$ ,  $x \in \mathbb{R}^n$  is given by the action of Gaussian semigroup

$$F \mapsto (G(t)F)(x) \equiv (4\pi t)^{-n/2} \int_{\mathbb{R}^n} e^{-|y|^2/4t} F(x - y) \, dy, \quad t > 0, \, x \in \mathbb{R}^n.$$

Suppose now that  $c \in \mathbb{C} \setminus \{0\}$ , a number  $t > 0$  is fixed and the function  $F(\cdot)$  is bounded, continuous and Levitan  $(x, \Lambda', c, \mathcal{P}_K)$ -almost periodic, resp. bounded, continuous and Bebutov  $(x, \Lambda', c, \mathcal{P}_K)$ -uniformly recurrent. Then an application of Proposition 3.3 shows that the function  $x \mapsto (G(t)F)(x)$ ,  $x \in \mathbb{R}^n$  is likewise bounded, continuous and Levitan  $(x, \Lambda', c, \mathcal{P}_K)$ -almost periodic, resp. bounded, continuous and Bebutov  $(x, \Lambda', c, \mathcal{P}_K)$ -uniformly recurrent (cf. also S. Zaidman [60, examples 4, 5, 7, 8; pp. 32-34], which can be also used for our purposes).

2. Suppose that  $1 < \beta < 2$ . The fractional diffusion-wave equation

$$\mathbb{D}_t^{(\beta)} u(t, x) = \Delta u(t, x), \quad t > 0, \, x \in \mathbb{R}^n,$$

subjected with the initial conditions

$$u(0, x) = u_0(x), \quad u_t(0, x) = 0,$$

where  $\mathbb{D}_t^{(\beta)}u(t, x)$  is the Caputo-Dzhrbashyan fractional derivative, given by

$$\mathbb{D}_t^{(\beta)}u(t, x) := \frac{1}{\Gamma(2-\beta)} \frac{\partial^2}{\partial^2 t} \int_0^t (t-\tau)^{1-\beta} u(\tau, x) d\tau - t^{1-\beta} \frac{u_t(0, x)}{\Gamma(2-\beta)} - t^{1-\beta} \frac{u(0, x)}{\Gamma(1-\beta)}, \quad t > 0,$$

was studied by many authors (see, e.g., the research article [30] by A. N. Kochubei and references cited therein). A unique solution  $u(t, x)$  of this problem is obtained as the convolution with a Green kernel

$$u(t, x) = \int_{\mathbb{R}^n} G(t, x - \xi) u_0(\xi) d\xi = \int_{\mathbb{R}^n} G(t, \xi) u_0(x - \xi) d\xi.$$

Since the Green kernel  $G(t, x)$  satisfies the estimate (cf. the research article [48] by A. V. Pskhu for more details;  $c > 0$  and  $a > 0$  denote the positive real constants here):

$$|G(t, x)| \leq ct^{-n\beta/2} \gamma_n(|x|t^{-\beta/2}) E(|x|t^{-\beta/2}),$$

where  $\gamma_n(z) := 1$ , if  $n = 1$  [ $\gamma_n(z) := |\ln z|$ , if  $n = 2$ ;  $\gamma_n(z) := z^{2-n}$ , if  $n \geq 3$ ] and  $E(z) := \exp(-az^{2/(2-\beta)})$ , the following can be easily shown: If the function  $x \mapsto u_0(x)$ ,  $x \in \mathbb{R}^n$  is bounded, continuous and  $(I', c)$ -almost periodic ( $(I', c)$ -uniformly recurrent; Levitan  $(x, \Lambda', c, \mathcal{P}_K)$ -almost periodic, Bebutov  $(x, \Lambda', c, \mathcal{P}_K)$ -uniformly recurrent) for some  $c \in \mathbb{C}$ , then the solution  $u(t, x)$  is bounded, continuous and  $(I', c)$ -almost periodic ( $(I', c)$ -uniformly recurrent; Levitan  $(x, \Lambda', c, \mathcal{P}_K)$ -almost periodic, Bebutov  $(x, \Lambda', c, \mathcal{P}_K)$ -uniformly recurrent) in the variable  $x$  for every fixed value of variable  $t > 0$ ; cf. [25] for the notion used.

3. Using the Laplace transform of vector-valued distributions, M. Kunzinger, E. A. Nigsch, and N. Ortner [38] have presented the explicit solutions for the Cauchy-Dirichlet problem of the iterated wave operator  $(\Delta_n + \partial_y^2 - \partial_t^2)^m$ , the iterated Klein-Gordon operator  $(\Delta_{2n+1} + \partial_y^2 - \partial_t^2 - \xi^2)^m$ , the iterated metaharmonic operator  $(\Delta_n + \partial_y^2 - p^2)^m$ , and the iterated heat operator  $(\Delta_n + \partial_y^2 - \partial_t)^m$ , in the half-space  $y > 0$ . In particular, the authors have shown that, under certain logical assumptions, the Cauchy problem

$$\begin{aligned} (\partial_{x_1}^2 + \dots + \partial_{x_n}^2 + \partial_y^2)^2 u &= 0 \quad \text{in } (x, y) \in \mathbb{R}^n \times (0, \infty); \\ u|_{y=0} &= g_0(x), \quad (\partial u_y)|_{y=0} = g_1(x) \end{aligned}$$

has a unique solution

$$\begin{aligned} u(x, y) &= 2 \frac{\Gamma((n+3)/2)}{\pi^{(n+1)/2}} y^3 \int_{\mathbb{R}^n} g_0(x - \xi) \frac{d\xi}{(|\xi|^2 + y^2)^{(n+3)/2}} \\ &\quad + \frac{\Gamma((n+1)/2)}{\pi^{(n+1)/2}} y^2 \int_{\mathbb{R}^n} g_1(x - \xi) \frac{d\xi}{(|\xi|^2 + y^2)^{(n+1)/2}}. \end{aligned}$$

Suppose now that the function  $x \mapsto (g_0(x), g_1(x))$ ,  $x \in \mathbb{R}^n$  is bounded, continuous and  $(I', c)$ -almost periodic ( $(I', c)$ -uniformly recurrent; Levitan  $(x, \Lambda', c, \mathcal{P}_K)$ -almost periodic, Bebutov  $(x, \Lambda', c, \mathcal{P}_K)$ -uniformly recurrent) for some  $c \in \mathbb{C}$ . Then it can be easily shown that the solution  $u(x, y)$  is bounded, continuous and  $(I', c)$ -almost periodic ( $(I', c)$ -uniformly recurrent; Levitan  $(x, \Lambda', c, \mathcal{P}_K)$ -almost periodic, Bebutov  $(x, \Lambda', c, \mathcal{P}_K)$ -uniformly recurrent) in the variable  $x$  for every fixed value of variable  $y > 0$ . This example and the first example given in this subsection can be also used to justify the introduction of multidimensional Levitan  $\rho$ -almost periodic

type functions with  $\rho$  being not a function but a general binary relation; cf. [25] for more details concerning this issue.

4. We can similarly consider the existence and uniqueness of Levitan almost periodic type solutions of the inhomogeneous abstract Cauchy problems considered in [15, Example 1.1], the backward wave equation considered in [60, Example 8, p. 33], and provide certain applications to the abstract ill-posed Cauchy problems considered on [33, pp. 543-545].

5. Concerning semilinear Cauchy problems, we will only note that the composition principle clarified in [33, theorem 7.1.18] can be simply formulated for the uniformly Poisson  $c$ -stable functions. Keeping in mind Proposition 3.3, we can simply consider the existence and uniqueness of the uniformly Poisson  $c$ -stable solutions for the class of semilinear Hammerstein integral equation of convolution type on  $\mathbb{R}^n$ ; see [33, p. 470] and the corresponding application made in [29, section 3]. Without going into full details, we will only note that the statement of [11, theorem 12] and some composition principles clarified in [33, subsection 6.1.5] can probably be formulated in our new framework.

6. Before proceeding to the next subsection, we would like to notice that the convolution invariance of Levitan  $N$ -almost periodicity has recently been considered by A. Nawrocki in [45, theorems 4.2, 4.4, and 4.8]. Concerning these statements, clarified in the one-dimensional setting, we would like to make the following comments:

- (i) Theorem 4.2 can be formulated for the uniformly Poisson stable (Levitan  $N$ -almost periodic) functions  $f : \mathbb{R} \rightarrow Y$ . Then the convolution  $f * g$  exists and it is a uniformly Poisson stable (Levitan  $N$ -almost periodic) function.
- (ii) Theorem 4.4 can be formulated for the Stepanov bounded, uniformly Poisson stable (Levitan  $N$ -almost periodic) functions  $f : \mathbb{R} \rightarrow Y$ . The resulting convolution  $f * g_\lambda$  exists and it is a bounded, uniformly Poisson stable (Levitan  $N$ -almost periodic) function.
- (iii) Concerning Theorem 4.8, we will only note that the Stepanov unboundedness condition [45, (4.2)] is not suitable for the work with the uniformly Poisson stable functions. Let us assume, in place of this condition, that for each real number  $\omega > 0$  and for each strictly increasing sequence  $(\alpha_k)$  of positive integers, we have

$$\limsup_{k \rightarrow +\infty} \int_{\alpha_k \omega}^{(\alpha_k + 1)\omega} f(t) dt = +\infty.$$

Then the resulting convolution  $f * g_\lambda$  exists and it is a bounded, uniformly Poisson stable function. The statement can be also formulated for the vector-valued functions.

In [45, Section 5], the author has applied the above results (cf. also [45, Corollaries 4.11–4.14]) in the analysis of the existence and uniqueness of Levitan  $N$ -almost periodic solutions of the linear ordinary differential equation of first order

$$y'(x) = \lambda y(x) + f(x), \quad x \in \mathbb{R}.$$

The statements of [45, lemma 5.5, theorem 5.10(i)] can be formulated for the uniformly Poisson stable solutions of this equation.

**3.3. On the wave equation in  $\mathbb{R}^n$**

In this subsection, we will revisit our recent investigations of the wave equation in  $\mathbb{R}^n$ , where  $n \leq 3$ . Let us consider first the following wave equation in  $\mathbb{R}^3$  :

$$u_{tt}(t, x) = d^2 \Delta_x u(t, x), \quad x \in \mathbb{R}^3, \quad t > 0; \quad u(0, x) = g(x), \quad u_t(0, x) = h(x), \tag{3.4}$$

where  $d > 0$ ,  $g \in C^3(\mathbb{R}^3 : \mathbb{R})$  and  $h \in C^2(\mathbb{R}^3 : \mathbb{R})$ . By the Kirchoff formula (see e.g., [50, theorem 5.4, pp. 277-278]), the function

$$\begin{aligned} u(t, x) &:= \frac{\partial}{\partial t} \left[ \frac{1}{4\pi d^2 t} \int_{\partial B_{dt}(x)} g(\sigma) \, d\sigma \right] + \frac{1}{4\pi d^2 t} \int_{\partial B_{dt}(x)} g(\sigma) \, d\sigma \\ &= \frac{1}{4\pi} \int_{\partial B_1(0)} g(x + dt\omega) \, d\omega + \frac{dt}{4\pi} \int_{\partial B_1(0)} \nabla g(x + dt\omega) \cdot \omega \, d\omega \\ &\quad + \frac{t}{4\pi} \int_{\partial B_1(0)} h(x + dt\omega) \, d\omega, \quad t \geq 0, \quad x \in \mathbb{R}^3, \end{aligned}$$

is a unique solution of problem (3.4) which belongs to the class  $C^2([0, \infty) \times \mathbb{R}^3)$ . Fix now a number  $t_0 > 0$ . Then the function  $x \mapsto u(t_0, x)$ ,  $x \in \mathbb{R}^3$  will be Levitan  $N$ -almost periodic (uniformly Poisson stable) provided that the function  $(g(\cdot), \nabla g(\cdot), h(\cdot))$  is Levitan  $N$ -almost periodic (uniformly Poisson stable), for example.

Let us consider now the following wave equation in  $\mathbb{R}^2$  :

$$u_{tt}(t, x) = d^2 \Delta_x u(t, x), \quad x \in \mathbb{R}^2, \quad t > 0; \quad u(0, x) = g(x), \quad u_t(0, x) = h(x), \tag{3.5}$$

where  $d > 0$ ,  $g \in C^3(\mathbb{R}^2 : \mathbb{R})$ , and  $h \in C^2(\mathbb{R}^2 : \mathbb{R})$ . By the Poisson formula (see e.g., [50, theorem 5.5, pp. 280-281]), we know that the function

$$\begin{aligned} u(t, x) &:= \frac{\partial}{\partial t} \left[ \frac{1}{2\pi d} \int_{\partial B_{dt}(x)} \frac{g(\sigma)}{\sqrt{d^2 t^2 - |x - y|^2}} \, d\sigma \right] + \frac{1}{2\pi d} \int_{\partial B_{dt}(x)} \frac{h(\sigma)}{\sqrt{d^2 t^2 - |x - y|^2}} \, d\sigma \\ &= d \int_{B_1(0)} \frac{g(x + dt\sigma)}{\sqrt{1 - |\sigma|^2}} \, d\sigma + d^2 t \int_{B_1(0)} \frac{\nabla g(x + dt\sigma) \cdot \sigma}{\sqrt{1 - |\sigma|^2}} \, d\sigma \\ &\quad + dt \int_{B_1(0)} \frac{h(x + dt\sigma)}{\sqrt{1 - |\sigma|^2}} \, d\sigma, \quad t \geq 0, \quad x \in \mathbb{R}^2, \end{aligned}$$

is a unique solution of problem (3.5) which belongs to the class  $C^2([0, \infty) \times \mathbb{R}^3)$ . Fix now a number  $t_0 > 0$ . Similarly as above, we have that the function  $x \mapsto u(t_0, x)$ ,  $x \in \mathbb{R}^2$  will be Levitan  $N$ -almost periodic (uniformly Poisson stable) provided that the function  $(g(\cdot), \nabla g(\cdot), h(\cdot))$  is Levitan  $N$ -almost periodic (uniformly Poisson stable).

Let us finally consider the wave equation  $u_{tt} = a^2 u_{xx}$  in domain  $\{(x, t) : x \in \mathbb{R}, t > 0\}$ , equipped with the initial conditions  $u(x, 0) = f(x) \in C^2(\mathbb{R})$  and  $u_t(x, 0) = g(x) \in C^1(\mathbb{R})$ . As is well known, its unique regular solution is given by the famous d'Alembert formula

$$u(x, t) = \frac{1}{2} [f(x - at) + f(x + at)] + \frac{1}{2a} \int_{x-at}^{x+at} g(s) \, ds, \quad x \in \mathbb{R}, \quad t > 0.$$

Our consideration from [15, example 1.2] shows that the solution  $u(x, t)$  can be extended to the whole real line in time variable and that  $u(x, t)$  will be uniformly  $c$ -Poisson stable in  $(x, t) \in \mathbb{R}^2$ , provided that the function  $t \mapsto (f(\cdot), \int_0^t g(s) ds)$ ,  $t \in \mathbb{R}$  is uniformly  $c$ -Poisson stable, with the meaning clear ( $c \in \mathbb{C} \setminus \{0\}$ ). We would like to emphasize that it is not clear how one can prove that the solution  $u(x, t)$  will be Levitan  $N$ -almost periodic in  $(x, t) \in \mathbb{R}^2$ , provided that the functions  $f(\cdot)$  and  $\int_0^t g(s) ds$  are Levitan  $N$ -almost periodic.

#### 4. Conclusions and final remarks

In this paper, we have investigated the Levitan and Bebutov approaches to the metrical approximations by trigonometric polynomials and  $\rho$ -periodic type functions, providing also certain applications to the abstract Volterra integro-differential equations and the partial differential equations. We have proved many structural results for the introduced classes of functions and propose several open problems, remarks, and illustrative examples.

Finally, we would like to mention some topics not considered in our former work:

1. The Levitan and Bebutov approaches to the metrical approximations by trigonometric polynomials and  $\rho$ -periodic type functions can be further generalized using the approaches of Stepanov, Weyl, and Besicovitch. Many structural results established in [14] can be reformulated in this context.

2. It is well known that every Levitan  $N$ -almost periodic function  $f : \mathbb{R} \rightarrow \mathbb{C}$  can be represented by the uniform limit of ratios of scalar-valued almost periodic functions. We will not discuss here the question whether this result continues to hold for multidimensional Levitan  $N$ -almost periodic functions.

3. In this paper, we have not analyzed  $\mathbb{D}$ -asymptotically Levitan (Bebutov) almost periodic type functions in general metric. Concerning this issue, we will only note that we have recently provided some new results about the existence and uniqueness of  $\mathbb{D}$ -asymptotically almost automorphic solutions of the inhomogeneous wave equation

$$u_{tt}(t, x) - d^2 \Delta_x u(t, x) = f(t, x), \quad x \in \mathbb{R}^2, \quad t > 0;$$

$$u(0, x) = g(x), \quad u_t(0, x) = h(x),$$

where  $d > 0$ ,  $f(t, x)$  is continuously differentiable in the variable  $t \in \mathbb{R}$  and continuous in the variable  $x \in \mathbb{R}$ ,  $g \in C^2(\mathbb{R}^2 : \mathbb{R})$  and  $h \in C^1(\mathbb{R}^2 : \mathbb{R})$ . It is well known that the unique solution of this problem is given by

$$u(x, t) = \frac{1}{2} [g(x - at) + g(x + at)] + \frac{1}{2a} \int_{x-at}^{x+at} h(s) ds$$

$$+ \frac{1}{2d} \int_0^t \left[ \int_{x-d(t-s)}^{x+d(t-s)} f(r, s) dr \right] ds, \quad x \in \mathbb{R}, \quad t > 0,$$

so that the conclusions established in [33, pp. 541-542] can be also formulated for  $\mathbb{D}$ -asymptotically uniformly Poisson stable solutions (defined in the obvious way), for example. Details can be left to the interested readers.

#### Acknowledgments

This research is partially supported by grant 174024 of Ministry of Science and Technological Development, Republic of Serbia and Bilateral project between Academy of Sciences and Arts of Macedonia and Serbian Academy of Sciences and Arts.

## References

- [1] Akhmet M, Fen MO, Tleubergenova M, Zhamanshin A. Unpredictable solutions of linear differential and discrete equations. *Turkish Journal of Mathematics* 2019; 43 (5): 2377-2389. <https://doi.org/10.3906/mat-1810-86>
- [2] Akhmet M, Fen MO, Tleubergenova M, Zhamanshin A. Poincare chaos for a hyperbolic quasilinear system. *Miskolc Mathematical Notes* 2019; 20 (1): 33-44. <https://doi.org/10.18514/MMN.2019.2826>
- [3] Akhmet M, Tleubergenova M, Zhamanshin A. Modulo periodic Poisson stable solutions of quasilinear differential equations. *Entropy* 2021; 23: 1535. <https://doi.org/10.3390/e23111535>
- [4] Akhmet M, Tleubergenova M, Zhamanshin A. Quasilinear differential equations with strongly unpredictable solutions. *Carpathian Journal of Mathematics* 2020; 36 (3): 341-349. <https://doi.org/10.37193/CJM.2020.03.02>
- [5] Akhmet M, Tleubergenova M, Zhamanshin A. Dynamics of shunting inhibitory cellular neural networks with variable two-component Passive decay rates and Poisson stable inputs. *Symmetry* 2022; 14 (6): 1162. <https://doi.org/10.3390/sym14061162>
- [6] Akhmet M, Tleubergenova M, Seilova R, Nugayeva Z. Poisson stability in symmetrical impulsive shunting inhibitory cellular neural networks with generalized piecewise constant argument. *Symmetry* 2022; 14 (9): 1754. <https://doi.org/10.3390/sym14091754>
- [7] Andres J, Pennequin D. Semi-periodic solutions of difference and differential equations. *Boundary Value Problems* 2012; 141: 1-16. <https://doi.org/10.1186/1687-2770-2012-141>
- [8] Basit B, Günzler H. Difference property for perturbations of vector-valued Levitan almost periodic functions and their analogs. *Russian Journal of Mathematical Physics* 2005; 12 (4): 424-438.
- [9] Bertotti G, Mayergoz ID. *The Science of Hysteresis*. New York, USA: Academic Press, 2005.
- [10] Besicovitch AS. *Almost Periodic Functions*. New York, USA: Dover Publications, 1954.
- [11] Bugajewski D, Gan X, Kasprzak P. Mappings of higher order and nonlinear equations in some spaces of almost periodic functions. *Nonlinear Analysis* 2012; 75 (13): 5294-5310. <https://doi.org/10.1016/j.na.2012.04.046>
- [12] Caraballo T, Cheban D. Levitan/Bohr almost periodic and almost automorphic solutions of second order monotone differential equations. *Journal of Differential Equations* 2011; 251 (3): 708-727. <https://doi.org/10.1016/j.jde.2011.04.021>
- [13] Chang YK, N'Guérékata GM, Ponce R. *Bloch-Type Periodic Functions: Theory and Applications to Evolution Equations*. Series on Concrete and Applicable Mathematics. 22, Singapore, Singapore: World Scientific, 2022.
- [14] Chaouchi B, Kostić M, Velinov D. Metrical approximations of functions. <https://doi.org/10.48550/arXiv.2205.09031> (submitted).
- [15] Chávez A, Khalil A, Kostić M, Pinto M. Multi-dimensional almost periodic type functions and applications. <https://doi.org/10.48550/arXiv.2012.00543> (submitted).
- [16] Chávez A, Khalil K, Kostić M, Pinto M. Multi-dimensional almost automorphic type functions and applications. *Bulletin of the Brazilian Mathematical Society, New Series* 2022; 53: 801-851. <https://doi.org/10.1007/s00574-022-00284-x>
- [17] Cheban D. Levitan almost periodic and almost automorphic solutions of  $V$ -monotone differential equations. *Journal of Dynamics and Differential Equations* 2008; 20: 669-697. <https://doi.org/10.1007/s10884-008-9101-x>
- [18] Cheban D. Levitan almost periodic solutions of infinite-dimensional linear differential equations. *Buletinul Academiei de Stiinte a Republicii Moldova. Matematica* 2019; 90 (2): 56-78.
- [19] Cheban D, Shcherbakov BA. Poisson asymptotic stability of motions of dynamical systems and their comparability with regard to the recurrence property in the limit. *Differential Equations* 1977; 13 (5): 898-906.
- [20] Cheban D, Liu Z. Poisson stable motions of monotone nonautonomous dynamical systems. *Science China Mathematics* 2019; 62 (7): 1391-1418. <https://doi.org/10.1007/s11425-018-9407-8>



- [21] Dads EA, Es-sebbar B, Lhachimi L. On Massera and Bohr-Neugebauer type theorems for some almost automorphic differential equations. *Journal of Mathematical Analysis and Applications* 2023; 518 (2): 126761. <https://doi.org/10.1016/j.jmaa.2022.126761>
- [22] Danilov LI. The uniform approximation of recurrent functions and almost recurrent functions. *Vestnik Udmurtskogo Universiteta. Matematika. Mekhanika. Komp'yuternye Nauki* 2013; 4: 36-54.
- [23] de Vries J. *Elements of Topological Dynamics. Mathematics and Its Applications.* 257, Dordrecht, Netherlands: Springer-Science+Business Media B.V., 1993.
- [24] Diagana T. *Almost Automorphic Type and Almost Periodic Type Functions in Abstract Spaces.* New York, USA: Springer-Verlag, 2013.
- [25] Fečkan M, Khalladi MT, Kostić M, Rahmani A. Multi-dimensional  $\rho$ -almost periodic type functions and applications. *Applicable Analysis.* accepted. <https://doi.org/10.1080/00036811.2022.2103678>
- [26] Fink AM. *Almost Periodic Differential Equations.* Berlin, Germany: Springer-Verlag, 1974.
- [27] Haraux A, Souplet P. An example of uniformly recurrent function which is not almost periodic. *Journal of Fourier Analysis and Applications* 2004; 10 (2): 217-220. <https://doi.org/10.1007/s00041-004-8012-4>
- [28] Hartman S. Remark on interpolation by  $L$ -almost periodic functions. *Colloquium Mathematicum* 1974; 30 (1): 133-136. <https://doi.org/10.4064/cm-30-1-133-136>
- [29] Khalladi MT, Kostić M, Pinto M, Rahmani A, Velinov D.  $c$ -Almost periodic type functions and applications. *Nonautonomous Dynamical Systems* 2020; 7: 176-193. <https://doi.org/10.1515/msds-2020-0111>
- [30] Kochubei AN. Fractional-hyperbolic systems. *Fractional Calculus and Applied Analysis* 2013; 16 (4): 860-873. <https://doi.org/10.2478/s13540-013-0053-4>
- [31] Kostić M. *Almost Periodic and Almost Automorphic Type Solutions to Integro-Differential Equations.* Berlin, Germany: Walter de Gruyter, 2019.
- [32] Kostić M. *Abstract Degenerate Volterra Integro-Differential Equations.* Belgrade, Serbia: Mathematical Institute of Serbian Academy of Sciences and Arts, 2020.
- [33] Kostić M. *Selected Topics in Almost Periodicity.* Berlin, Germany: Walter de Gruyter, 2022.
- [34] Kostić M. Metrical almost periodicity and applications. *Annales Polonici Mathematici.* accepted. <https://doi.org/10.4064/ap220510-15-11>
- [35] Kostić M. Stepanov  $\rho$ -almost periodic functions in general metric. *Facta Universitatis-Series Mathematics And Informatics* 2022; 37 (2): 345-358. <https://doi.org/10.22190/FUMI211217024K>
- [36] Kostić M. Weyl  $\rho$ -almost periodic functions in general metric. *Mathematica Slovaca.* accepted. <https://www.researchgate.net/publication/357129349>
- [37] Kostić M. Multi-dimensional Besicovitch almost periodic type functions and applications. *Communications on Pure and Applied Analysis* 2022; 21 (12): 4215-4250. <https://doi.org/10.3934/cpaa.2022141>
- [38] Kunzinger M, Nigsch EA, Ortner N. Laplace transformation of vector-valued distributions and applications to Cauchy-Dirichlet problems. *Journal of Mathematical Analysis and Applications* 2019; 478 (2): 990-1004. <https://doi.org/10.1016/j.jmaa.2019.06.002>
- [39] Levin BY. On the almost periodic functions of Levitan. *Ukrainian Mathematical Journal* 1949; 1 (1): 49-100.
- [40] Levitan M. *Pochti peridicheskiye funkci.* Moscow, Russia: GITTL, 1953 (in Russian).
- [41] Levitan BM, Zhikov VV. *Almost Periodic Functions and Differential Equations.* Moscow, Russia: University Publishing House, Moscow, 1978 (English translation by Cambridge University Press, 1982).
- [42] Liu X, Liu ZX. Poisson stable solutions for stochastic differential equations with Lévy noise. *Acta Mathematica Sinica, English Series* 2022; 38: 22-54. <https://doi.org/10.1007/s10114-021-0107-1>

- [43] Lyubarskii MG. Rasprostraneniye teorii Favara na sluchay sistemy lineynykh differentsialnyye uravneniya s neogranichennymi pocti–periodicheskimi po levitanu koeffitsiyentami. Doklady Akademii Nauk SSSR 1972; 206 (4): 808–810 (in Russian).
- [44] Marchenko VA. Obobshchennyye pocti peridicheskiye funkicii. Doklady Akademii Nauk SSSR 1950; 74: 893–895 (in Russian).
- [45] Nawrocki A. On some applications of convolution to linear differential with Levitan almost periodic coefficients. Topological Methods in Nonlinear Analysis 2017; 50: 489–512.
- [46] N’Guérékata GM. Almost Automorphic and Almost Periodic Functions in Abstract Spaces. Dordrecht, Netherlands: Kluwer Academic Publishers, 2001.
- [47] Pankov AA. Bounded and Almost Periodic Solutions of Nonlinear Operator Differential Equations. Dordrecht, Netherlands: Kluwer Academic Publishers, 1990.
- [48] Pskhu AV. The fundamental solution of a diffusion-wave equation of fractional order. Izvestiya:Mathematics 2009; 73 (2): 351–392. <https://doi.org/10.1070/IM2009v073n02ABEH002450>
- [49] Reich A. Präkompakte gruppen und fastperiodizität. Mathematische Zeitschrift 1970; 114: 218–234 (in German).
- [50] Salsa S. Partial Differential Equations in Action: From Modelling to Theory. Milano, Italia: Springer-Verlag, 2008.
- [51] Shcherbakov BA. Topologicheskaya dinamika i ustoychivosti po Puassonu resheniy differencialnih uravneniya. Chisinau, Moldova: Stiinta 1972 (in Russian).
- [52] Shcherbakov BA. Ustoychivosti po Puassonu dvizheniy dinamicheskikh sistem i resheniy Differencialnyye uravneniya. Chisinau, Moldova: Stiinta, 1985 (in Russian).
- [53] Shcherbakov BA. Klassifikaciya ustojchivih po Puassonu dvizheniy. Pseudorekurrentnie dvizheniya. Doklady Akademii Nauk SSSR 1962; 146 (2): 322–324 (in Russian).
- [54] Shcherbakov BA. Rekurrentnie reshenija differencialnih uravneniy. Doklady Akademii Nauk SSSR 1966; 167 (5): 1004–1007 (in Russian).
- [55] Shcherbakov BA. Ob odnom klasse ustojchivih po Puassonu resheniy differencialnih uravneniy. Differencialnye Uravnenija 1968; 4 (2): 238–243 (in Russian).
- [56] Shcherbakov BA. O sravnimosti po harakteru vozvrashtaemosti dvizheniy dinamicheskikh sistem. Differencialnye Uravnenija. 1975; 11 (7): 1246–1255 (in Russian).
- [57] Sell G. Lectures on Topological Dynamics and Differential Equations. Van Nostrand Reinhold Mathematical Studies. 33, London, England: Van Nostrand Reinhold Company, 1971.
- [58] Shen W, Yi Y. Almost automorphic and almost periodic dynamics in skew-product semiflows. Memoirs of the American Mathematical Society 1998; 136: 647. <https://doi.org/http://dx.doi.org/10.1090/memo/0647>
- [59] Yuan R. On Favard’s theorems. Journal of Differential Equations 2010; 249 (8): 1884–1916. <https://doi.org/10.1016/j.jde.2010.07.014>
- [60] Zaidman S. Almost-Periodic Functions in Abstract Spaces. Pitman Research Notes in Mathematics. 126, Boston, USA: Pitman, 1985.