

9-25-2023

On positive periodic solutions to third-order integro-differential equations with distributed delays

MIMIA BENHADRI

TOMAS CARABALLO

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)



Recommended Citation

BENHADRI, MIMIA and CARABALLO, TOMAS (2023) "On positive periodic solutions to third-order integro-differential equations with distributed delays," *Turkish Journal of Mathematics*: Vol. 47: No. 6, Article 15. <https://doi.org/10.55730/1300-0098.3465>

Available at: <https://journals.tubitak.gov.tr/math/vol47/iss6/15>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

On positive periodic solutions to third-order integro-differential equations with distributed delays

Mimia BENHADRI¹ , Tomás CARABALLO^{2,3,*} 

¹Department of Mathematics, Faculty of Sciences, University 20 août 1955, Skikda, Algeria

²Departamento de Ecuaciones Diferenciales y Análisis Numérico, Facultad de Matemáticas,
Universidad de Sevilla, c/ Tarfia s/n, Sevilla, Spain

³Department of Mathematics, Wenzhou University, Wenzhou, Zhejiang Province, P. R. China

Received: 30.10.2022

Accepted/Published Online: 08.08.2023

Final Version: 25.09.2023

Abstract: In this paper, we investigate the existence of positive periodic solutions of a third-order nonlinear integro-differential equation with distributed delays, by using the Green function and the Krasnosel'skii fixed point theorem in cones of Banach spaces, providing new results on this field. Three examples are analyzed to illustrate the effectiveness of the abstract results.

Key words: Fixed point theorem, positive periodic solutions, third-order delay differential equations, distributed delays

1. Introduction

Delay differential equations form an important branch of functional differential equations that take into account the dependence on their history, which makes it possible to predict the future evolution of the system in a more reliable and efficient way. Such processes take place in the theories of optimal control, population dynamics, economics, mathematical ecological models, and other biotechnological systems, but they are characterized by specific properties which make their study difficult in both concepts and techniques. Note that many results concerning the theory of delay functional differential equations can be found in the monographs by Hale and Meyer [16], Hale and Lunel [17], Kuang [19] amongst others.

Third-order differential equations with and without delay have been studied by many authors, since they describe several models derived from natural phenomena, such as wave propagation in thermally relaxing viscous fluids or flexible space structures with internal damping, for example, a thin uniform rectangular panel and a spaceship with flexible attachments, and many others (see, e.g., [5]- [7], [12], [14], [15], [26], [27]). This means that it is interesting and meaningful to study the properties of the solutions of the differential equations of the third order with and without delay.

Recently, intensive scientific work has been carried out in various dynamical aspects of third-order delay differential equations, functional delay differential equations, and many results have been reported in the literature. For instance, uniqueness of periodic solutions and some other fundamental properties of solutions

*Correspondence: caraball@us.es

2010 *AMS Mathematics Subject Classification:* 34K20, 34K13, 92B20

of certain delay differential equations have been discussed in ([1]- [4], [10], [11], [18], [20]- [25], [29], [32], [33]). Some classical tools have been used to study delay differential equation in the literature, including the fixed point index theorem [25], Krasnoselskii's fixed point theorem ([2], [3]), the fixed point theorem in cones ([23], [24], [29], [33]), fixed point theorem of Leray-Schauder type [11], Mawhin's continuous theorem [4], and the continuation theorem of coincidence degree theory [1].

As far as we know, the existence of positive periodic solutions for third-order integro-differential equations with distributed delay that are considered in the present paper have not been investigated yet. For this reason, in this paper, we make a first attempt to fill this gap and obtain new sufficient conditions for the existence of one positive periodic solution thanks to the use of a fixed point theorem on cones. We are particularly motivated and inspired by the papers [2], [22], [23], [24], [25], [33] and the references therein.

In [28], the following third-order nonlinear delay differential equation with periodic coefficients is considered:

$$y'''(t) + p(t)y''(t) + q(t)y'(t) + r(t)y(t) = f(t, y(t)). \tag{1.1}$$

By applying a fixed point theorem index, the authors derived some verifiable sufficient conditions for the existence of a positive periodic solution to (1.1).

Very recently, Ardjouni and Djoudi [2] considered the following third-order nonlinear delay differential equation with periodic coefficients:

$$y'''(t) + p(t)y''(t) + q(t)y'(t) + r(t)y(t) = f(t, y(t - \tau(t))) + \frac{d}{dt}g(t, y(t - \tau(t))). \tag{1.2}$$

Thanks to the Green function and the Krasnoselskii fixed point theorem, they obtained a set of easily verifiable sufficient conditions for the existence of positive periodic solutions to equation (1.2).

In 2006, Li and Wang [22] studied the existence of positive periodic solutions of the following two kinds of nonlinear neutral differential equations

$$\frac{d}{dt}(y(t) - cy(t - \tau(t))) = -a(t)y(t) + g(t, y(t - \tau(t))), \tag{1.3}$$

and

$$\frac{d}{dt}\left(y(t) - c \int_{-\infty}^0 K(r)y(t+r) dr\right) = -a(t)y(t) + b(t) \int_{-\infty}^0 K(r)g(t, y(t+r)) dr. \tag{1.4}$$

By using the theory of fixed point index in cones, sufficient conditions are presented for the existence of positive periodic solutions of two kinds of neutral differential equations with periodic coefficients.

In [33], the authors analyzed the following impulsive equations

$$\begin{aligned} y'(t) &= -a(t)y(t) + \int_{-\infty}^0 K(r)g(t, y(t+r)) dr, t \neq t_k, \\ y(t_k^+) &= y(t_k^-) + I_k(y(t_k)), t = t_k, k \in \mathbb{Z}, \end{aligned} \tag{1.5}$$

$$\begin{aligned} y'(t) &= a(t)y(t) - \int_{-\infty}^0 K(r)g(t, y(t+r)) dr, t \neq t_k, \\ y(t_k^+) &= y(t_k^-) + I_k(y(t_k)), t = t_k, k \in \mathbb{Z}, \end{aligned} \tag{1.6}$$

where $y(t_k^+)$ and $y(t_k^-)$ represent the right and the left limit of $y(t_k)$, $a \in C(\mathbb{R}, \mathbb{R}^+)$, $g \in C(\mathbb{R} \times \mathbb{R}^+, \mathbb{R}^+)$, $I_k \in C(\mathbb{R}^+, \mathbb{R}^+)$, a and $g(t, y)$ are ω -periodic functions, where ω is a positive constant. Moreover, $K \in C(\mathbb{R}^-, \mathbb{R}^+)$

with $\int_{-\infty}^0 K(r) dr = 1$. There exists an integer $q > 0$ such that $t_{k+q} = t_k + \omega$, $I_{(k+q)}(y(t_{k+q})) = I_k(y(t_k))$, $k \in \mathbb{Z}$, where $0 < t_1 < t_2 < \dots < t_q < \omega$.

By using the theory of fixed point index in cones, existence theory for single and multiple positive periodic solutions to a kind of nonautonomous Volterra integro-differential equations with impulse effects (1.5), (1.6) has been investigated.

Motivated by the above statements, in this paper we study a third-order nonlinear integro-differential equations with distributed delays of the following form, by using a fixed point on a cone:

$$\begin{aligned}
 & y'''(t) + p(t)y''(t) + q(t)y'(t) + r(t)y(t) \\
 &= \frac{d}{dt} \int_{-\infty}^0 K(r) c(t) y(t+r) dr \\
 &+ d(t) \int_{-\infty}^0 K(r) f(t, y(t+r)) dr,
 \end{aligned}
 \tag{1.7}$$

where $d \in C(\mathbb{R}, \mathbb{R}^+ \setminus \{0\})$, $c, p, q, r \in C(\mathbb{R}, \mathbb{R}^+)$. The function $f : \mathbb{R} \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is continuous. Moreover, K is a continuous and integrable function on $]-\infty, 0]$ with $\int_{-\infty}^0 K(r) dr = 1$.

In order to prove our main results in the next section, we need the following definition and theorem.

Definition 1.1. (see [30]) *Let X be a Banach space and let K be a closed, nonempty subset of X . K is said to be a cone if*

- i) $\alpha x + \beta y \in K$ for all $x, y \in K$ and all $\alpha, \beta \geq 0$;*
- ii) $y, -y \in K$ imply $y = 0$.*

We now state the Krasnosel'skii fixed point theorem.

Theorem 1.1. (see [14], Theorem 2.3.3 on p. 93). *Let X be a Banach space, and let $K \subset X$ be a cone in X . Assume that Ω_1 and Ω_2 are open subsets of X with $0 \in \Omega_1$, $\bar{\Omega}_1 \subset \Omega_2$ and let*

$$\Phi : K \cap (\bar{\Omega}_2 \setminus \Omega_1) \rightarrow K,$$

be a completely continuous operator such that either

- a) $\|\Phi y\| \leq \|y\|$ for $y \in K \cap \partial\Omega_1$ and $\|\Phi y\| \geq \|y\|$ for $y \in K \cap \partial\Omega_2$; or*
- b) $\|\Phi y\| \geq \|y\|$ for $y \in K \cap \partial\Omega_1$ and $\|\Phi y\| \leq \|y\|$ for $y \in K \cap \partial\Omega_2$.*

Then Φ has a fixed point in $K \cap (\bar{\Omega}_2 \setminus \Omega_1)$.

This paper is devoted to studying the existence of positive periodic solutions of Equation (1.7) by using the Krasnosel'skii fixed point theorem and some mathematical analysis techniques. Equation (1.7) is a nonneutral third-order nonlinear differential equation, including the existing classical third-order differential equations, as well as the equations considered in ([27, 28]). Therefore, the results of this paper are more general and better applicable. The research method of this paper is different from the existing research methods (see, e.g., [2, 25, 28]). For more results about third-order differential equation with and without delays, see, e.g., ([1, 4, 8, 10, 25–27, 31]) and cited references.

The organization of this paper is as follows. In Section 2, we present the inversion of equation and state some preliminary results needed in later sections. We also describe the Green function of Eq. (1.7), which plays an important role in this paper. In Section 3, we establish our main results ensuring the existence of positive periodic solutions by applying the fixed point theorem in cones established previously. In Section 4, three examples are exhibited to illustrate that our results are feasible and more general than previous ones in the literature.

2. Green’s function and periodicity

In this section, we state and define the Green function for periodic solutions of third-order nonlinear integro-differential equations with distributed delays (1.7).

Let

$$C_\omega = \{y \in C(\mathbb{R}, \mathbb{R}), y(t + \omega) = y(t) \text{ for } t \in \mathbb{R}\},$$

with the norm

$$\|y\| = \max_{t \in [0, \omega]} |y(t)|.$$

It is easy to verify that $(C_\omega, \|\cdot\|)$ is a Banach space.

In this paper, we give the assumptions as follows that will be used in the main results.

(A1) There exist differentiable ω -periodic functions $a_1, a_2 \in C(\mathbb{R}, \mathbb{R}^+)$ and a positive real constant ρ such that

$$\begin{cases} a_1(t) + \rho = p(t), \\ a_1'(t) + a_2(t) + \rho a_1(t) = q(t), \\ a_2'(t) + \rho a_2(t) = r(t). \end{cases}$$

(A2) $p, q, r \in C(\mathbb{R}, \mathbb{R}^+)$ are ω -periodic functions and

$$\int_0^\omega q(u) du > 0, \int_0^\omega p(u) du > \rho\omega.$$

(A3) The function $f \in C(\mathbb{R} \times \mathbb{R}^+, \mathbb{R}^+)$ is a ω -periodic in t , and $c \in C(\mathbb{R}, \mathbb{R}^+)$, $d \in C(\mathbb{R}, \mathbb{R}^+ \setminus \{0\})$ are ω -periodic. For all $y \in \mathbb{R}^+$,

$$f(t, y) \geq \rho \frac{c(t)}{d(t)} y, \forall t \in [0, \omega].$$

We consider

$$y'''(t) + p(t)y''(t) + q(t)y'(t) + r(t)y(t) = h(t), \tag{2.1}$$

where $h \in C_\omega$. Obviously, by condition (A1), Eq. (2.1) can be transformed into

$$\begin{cases} z'(t) + \rho z(t) = h(t), \\ y''(t) + a_1(t)y'(t) + a_2(t)y(t) = z(t). \end{cases}$$

Lemma 2.1 (see [3]). *If $z, h \in C_\omega$, then z is a solution of equation*

$$z'(t) + \rho z(t) = h(t),$$

if and only if

$$z(t) = \int_t^{t+\omega} G_1(t, s) h(s) ds, \tag{2.2}$$

where

$$G_1(t, s) = \frac{\exp(\rho(s-t))}{\exp(\rho\omega) - 1}. \tag{2.3}$$

Corollary 2.1 (see [21]). *Green function G_1 satisfies the following properties*

$$\begin{aligned} G_1(t + \omega, s + \omega) &= G_1(t, s), G_1(t, t + \omega) = G_1(t, t) \exp(\rho\omega), \\ G_1(t + \omega, s) &= G_1(t, s) \exp(-\rho\omega), G_1(t, s + \omega) = G_1(t, s) \exp(\rho\omega), \\ \frac{\partial}{\partial t} G_1(t, s) &= -\rho G_1(t, s), \\ \frac{\partial}{\partial s} G_1(t, s) &= \rho G_1(t, s), \end{aligned}$$

and

$$m_1 \leq G_1(t, s) \leq M_1, \tag{2.4}$$

where

$$M_1 = \frac{1}{\exp(\rho\omega) - 1}, m_1 = \frac{\exp(\rho\omega)}{\exp(\rho\omega) - 1}. \tag{2.5}$$

Lemma 2.2 (see [32]). *Suppose that (A1) and (A2) hold and*

$$\frac{\bar{R}_1 [\exp(\int_0^\omega a_1(u) du) - 1]}{Q_1 \omega} \geq 1, \tag{2.6}$$

where

$$\begin{aligned} \bar{R}_1 &= \max_{t \in [0, \omega]} \left| \int_t^{t+\omega} \frac{\exp(\int_0^\omega a_1(u) du)}{\exp(\int_0^\omega a_1(u) du) - 1} a_2(s) ds \right|, \\ Q_1 &= \left(1 + \exp\left(\int_0^\omega a_1(u) du\right) \right)^2 \bar{R}_1^2. \end{aligned}$$

Then there are continuous ω - periodic functions a and b such that

$$b(t) > 0, \quad \int_0^\omega a(u) du > 0,$$

and

$$a(t) + b(t) = a_1(t), \quad b'(t) + a(t)b(t) = a_2(t), \quad \text{for } t \in \mathbb{R}.$$

Lemma 2.3 (see [21]). *Suppose the conditions of Lemma 2.2 hold and $y \in C_\omega$. Then, the equation*

$$y''(t) + a_1(t)y'(t) + a_1(t)y(t) = h(t),$$

possesses an ω - periodic solution. Moreover, the periodic solution can be expressed by

$$y(t) = \int_t^{t+T} G_2(t, s) h(s) ds, \tag{2.7}$$

where

$$G_2(t, s) = \frac{\int_t^s \exp \left[\int_t^u b(v) dv + \int_u^s a(v) dv \right] du}{\left[\exp \left(\int_0^\omega a(u) du \right) - 1 \right] \left[\exp \left(\int_0^\omega b(u) du \right) - 1 \right]} + \frac{\int_s^{t+\omega} \exp \left[\int_t^u b(v) dv + \int_u^{s+\omega} a(v) dv \right] du}{\left[\exp \left(\int_0^\omega a(u) du \right) - 1 \right] \left[\exp \left(\int_0^\omega b(u) du \right) - 1 \right]}. \tag{2.8}$$

Corollary 2.2 (see [32]). *The Green function G_2 satisfies the following properties:*

$$\begin{aligned} G_2(t + \omega, s + \omega) &= G_2(t, s), \quad G_2(t, t + \omega) = G_2(t, t), \\ G_2(t + \omega, s) &= \exp \left(- \int_0^\omega b(v) dv \right) \\ &\quad \times \left[G_2(t, s) + \int_t^{t+\omega} E(t, u) F(u, s) du \right], \\ \frac{\partial}{\partial s} G_2(t, s) &= a(s) G_2(t, s) - \frac{\exp \left(\int_t^s b(v) dv \right)}{\exp \left(\int_0^\omega b(v) dv \right) - 1}, \\ \frac{\partial}{\partial t} G_2(t, s) &= -b(t) G_2(t, s) + \frac{\exp \left(\int_t^s a(v) dv \right)}{\exp \left(\int_0^\omega a(v) dv \right) - 1}, \end{aligned}$$

where

$$E(t, s) = \frac{\exp \left(\int_t^s b(v) dv \right)}{\exp \left(\int_0^\omega b(v) dv \right) - 1}, \quad F(t, s) = \frac{\exp \left(\int_t^s a(v) dv \right)}{\exp \left(\int_0^\omega a(v) dv \right) - 1}. \tag{2.9}$$

Lemma 2.4 (see [21]). *Let $A = \int_0^\omega a_1(u) du$ and $B = \omega^2 \exp \left(\frac{1}{\omega} \int_0^\omega \ln(a_2(u)) du \right)$. If*

$$A^2 \geq 4B, \tag{2.10}$$

then

$$\begin{aligned} \min \left\{ \int_0^\omega a(u) du, \int_0^\omega b(u) du \right\} &\geq l, \\ \max \left\{ \int_0^\omega a(u) du, \int_0^\omega b(u) du \right\} &\leq L, \end{aligned}$$

where

$$l = \frac{1}{2} \left(A - \sqrt{A^2 - 4B} \right), \quad L = \frac{1}{2} \left(A + \sqrt{A^2 - 4B} \right). \tag{2.11}$$

Corollary 2.3 (see [32]). *Functions $G_2, E,$ and F satisfy*

$$m_2 \leq G_2(t, s) \leq M_2, \tag{2.12}$$

$$E(t, s) \leq \frac{e^L}{e^l - 1}, \tag{2.13}$$

$$F(t, s) \leq e^L, \tag{2.14}$$

where

$$m_2 = \frac{\omega}{(\exp(L) - 1)^2}, M_2 = \frac{\omega \left(\exp \int_0^\omega a_1(u) du \right)}{(\exp(l) - 1)^2}, \tag{2.15}$$

and

$$\exp(L) \neq 1, \exp(l) \neq 1.$$

Lemma 2.4 (see [8]). *Suppose the conditions of Lemma 2.2 hold and $h \in C_\omega$. Then, the equation*

$$y'''(t) + p(t)y''(t) + q(t)y'(t) + r(t)y(t) = h(t),$$

possesses an ω -periodic solution. Moreover, the periodic solution can be expressed by

$$y(t) = \int_t^{t+\omega} G(t, s) h(s) ds, \tag{2.16}$$

where

$$G(t, s) = \int_t^{t+\omega} G_2(t, \sigma) G_1(\sigma, s) d\sigma. \tag{2.17}$$

Corollary 2.4 (see [25]). *The Green function G satisfies the following properties*

$$G(t + \omega, s + \omega) = G(t, s), G(t, t + \omega) = G(t, t) \exp(\rho\omega),$$

$$\begin{aligned} \frac{\partial}{\partial t} G(t, s) &= (\exp(-\rho\omega) - 1) G_1(t, s) G_2(t, s) \\ &\quad - b(t) G(t, s) + \int_t^{t+\omega} F(t, \sigma) G_1(\sigma, s) d\sigma, \end{aligned}$$

$$\frac{\partial}{\partial s} G(t, s) = \rho G(t, s),$$

and

$$m \leq G(t, s) \leq M, \tag{2.18}$$

where

$$m = \frac{\omega^2}{(\exp(l) - 1)^2 (\exp(\rho\omega) - 1)}, M = \frac{\omega^2 \exp(\rho\omega + \int_0^\omega a(u) du)}{(\exp(l) - 1)^2 (\exp(\rho\omega) - 1)}, \tag{2.19}$$

and

$$\exp(\rho\omega) \neq 1, \exp(l) \neq 1.$$

Before stating the main result of this paper, we establish the equivalent integral formulation for the solution of equation (1.7).

Lemma 2.5. *Assume that (A1)-(A3) and (2.6) hold. The function $y(\cdot)$ is an ω -periodic solution of equation (1.7) if and only if $y(\cdot)$ is an ω -periodic solution of the following equation*

$$\begin{aligned}
 y(t) &= \int_t^{t+\omega} G(t, s) \left[d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right. \\
 &\quad \left. - \rho \int_{-\infty}^0 K(r) c(s) y(s+r) dr \right] ds \\
 &\quad + (\exp(\rho\omega) - 1) G(t, t) \int_{-\infty}^0 K(r) c(s) y(s+r) dr.
 \end{aligned} \tag{2.20}$$

Proof Let $y \in C_\omega$ be a solution of (1.7). From Lemma 2.4, we have

$$\begin{aligned}
 y(t) &= \int_t^{t+\omega} G(t, s) \left[d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right. \\
 &\quad \left. + \frac{\partial}{\partial s} \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\
 &= \int_t^{t+\omega} G(t, s) \left(d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right) ds \\
 &\quad + \int_t^{t+\omega} G(t, s) \frac{\partial}{\partial s} \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) ds.
 \end{aligned} \tag{2.21}$$

Performing an integration by parts, we obtain

$$\begin{aligned}
 &\int_t^{t+\omega} G(t, s) \frac{\partial}{\partial s} \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \\
 &= G(t, s) \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \Big|_t^{t+\omega} \\
 &\quad - \int_t^{t+\omega} \left[\frac{\partial}{\partial s} G(t, s) \right] \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) ds \\
 &= G(t, t+\omega) \int_{-\infty}^0 K(r) c(t+\omega) y(t+\omega+r) dr \\
 &\quad - G(t, t) \int_{-\infty}^0 K(r) c(t) y(t+r) dr - \int_t^{t+\omega} \left[\frac{\partial}{\partial s} G(t, s) \right] \int_{-\infty}^0 K(r) c(s) y(s+r) dr ds \\
 &= (G(t, t) \exp(\rho\omega)) \int_{-\infty}^0 K(r) c(t) y(t+r) dr - G(t, t) \int_{-\infty}^0 K(r) c(t) y(t+r) dr \\
 &\quad - \rho \int_t^{t+\omega} G(t, s) \int_{-\infty}^0 K(r) c(s) y(s+r) dr ds
 \end{aligned}$$

$$\begin{aligned}
 &= (\exp(\rho\omega) - 1) G(t, t) \int_{-\infty}^0 K(r) g(t, y(t+r)) dr \\
 &\quad - \rho \int_t^{t+\omega} G(t, s) \int_{-\infty}^0 K(r) c(s) y(s+r) dr ds.
 \end{aligned}
 \tag{2.22}$$

We obtain (2.20) by replacing (2.21) into (2.22). □

3. Existence of positive periodic solutions

This section is devoted to establish three theorems and two corollaries, which are the main results of our paper. In the analysis we will carry out, we use the idea and concept of Green’s function for third-order periodic solutions and transform equation (1.7) into an equivalent integral one. Then, by means of the Krasnosel’skii fixed theorem in cones of Banach spaces, we show the existence of at least one positive periodic solution of Eq. (1.7) in three different theorems. Some new and interesting sufficient conditions are obtained to guarantee the existence of such positive periodic solutions to (1.7).

Since Eq. (2.20) is equivalent to Eq. (1.7), we just have to study the existence of positive periodic solutions to (2.20). To this end, we will use Theorem 1.1 where we consider $(X, \|\cdot\|) = (C_\omega, \|\cdot\|)$.

Now, let

$$\delta = \frac{m}{M} = \frac{1}{\exp(\rho\omega + \int_0^\omega a(u)du)} \in (0, 1),$$

and define K as a cone in C_ω by

$$K = \{y(\cdot) \in C_\omega : y(t) \geq 0, \text{ and } y(t) \geq \delta \|y\|, t \in [0, \omega]\},$$

where m, M are given by (2.19). It is not difficult to check that K is a cone in C_ω .

Thanks to (2.20), we define operator $\Phi : C_\omega \rightarrow C_\omega$ by:

$$\begin{aligned}
 (\Phi y)(t) &= \int_t^{t+\omega} G(t, s) \left[\left(d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right) \right. \\
 &\quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\
 &\quad + (\exp(\rho\omega) - 1) G(t, t) \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right).
 \end{aligned}
 \tag{3.1}$$

To simplify our description, we introduce the following constants

$$\begin{aligned}
 \pi^* &= \exp(L), d^* = \max_{t \in [0, \omega]} |d(t)|, c^* = \max_{t \in [0, \omega]} |c(t)|, b^* = \max_{t \in [0, \omega]} |b(t)|, \\
 \hat{d} &= \min_{t \in [0, \omega]} |d(t)|, \hat{c} = \min_{t \in [0, \omega]} |c(t)|, \tilde{\theta} = \exp(\rho\omega) - 1.
 \end{aligned}
 \tag{3.2}$$

Thus, the existence of a positive periodic solution of equation (1.7) is equivalent to finding a fixed point of operator Φ .

Lemma 3.1. *Assume (A1)-(A3) hold. Then $\Phi: K \rightarrow K$ is well defined.*

Proof From (3.1), it is easy to verify that $(\Phi y)(t)$ is continuous in t . Moreover, for any $y \in K$,

$$\begin{aligned} & (\Phi y)(t + \omega) \\ &= \int_{t+\omega}^{t+2\omega} G(t + \omega, s) \left[d(s) \left(\int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right) \right. \\ & \quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\ & \quad + (\exp(\rho\omega) - 1) G(t + \omega, t + \omega) \times \\ & \quad \quad \times \left(\int_{-\infty}^0 K(r) c(t + \omega) y(t + \omega + r) dr \right) \\ &= \int_t^{t+\omega} G(t + \omega, s + \omega) \left[d(s + \omega) \left(\int_{-\infty}^0 K(r) f(s + \omega, y(s + \omega + r)) dr \right) \right. \\ & \quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s + \omega) y(s + \omega + r) dr \right) \right] ds \\ & \quad + (\exp(\rho\omega) - 1) G(t + \omega, t + \omega) \left(\int_{-\infty}^0 K(r) c(t + \omega) y(t + \omega + r) dr \right) \\ & \\ &= \int_t^{t+\omega} G(t, s) \left[d(s) \left(\int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right) \right. \\ & \quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\ & \quad + (\exp(\rho\omega) - 1) G(t, t) \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right) \\ &= (\Phi y)(t). \end{aligned}$$

Therefore, $(\Phi y) \in C_\omega$. By condition (A3), we have

$$(\Phi y)(t) \geq 0, \text{ for all } y \in K.$$

Also, for $y \in K$, by using (2.19) and (3.1), we deduce

$$\begin{aligned} |(\Phi y)| &\leq M \int_0^\omega \left[\left(d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right) \right. \\ & \quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\ & \quad + (\exp(\rho\omega) - 1) \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right). \end{aligned}$$

Noticing that

$$G(t, s) \left[d(t) \int_{-\infty}^0 K(r) f(t, y(t+r)) dr - \rho \int_{-\infty}^0 K(r) c(t)y(t+r)dr \right] \geq 0,$$

we obtain

$$\begin{aligned} (\Phi y)(t) &\geq m \int_0^\omega \left[\left(d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right) \right. \\ &\quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\ &\quad + (\exp(\rho\omega) - 1) \left(\int_{-\infty}^0 K(r) c(t)y(t+r)dr \right) \\ &\geq \frac{m}{M} \|\Phi y\| \geq \delta \|\Phi y\|. \end{aligned}$$

That is, ΦK is contained in K . The proof of Lemma 3.1 is complete. □

Lemma 3.2. *Assume that (A1)-(A3), (2.6), (2.10) hold. Then $\Phi : K \rightarrow K$ is completely continuous.*

Proof Now we have to show that Φ is continuous. Let $y_n, (n = 1, 2, \dots)$ be a sequence in K such that

$$\lim_{n \rightarrow \infty} \|y_n - y_0\| = 0.$$

Since K is closed, we have $y_0 \in K$. Since $y, f(t, y)$ are ω -periodic and continuous functions, $f(s, y_n(s+r)) \rightarrow f(s, y_0(s+r))$ uniformly, for $s \in [0, \omega]$.

$$\begin{aligned} &|(\Phi y_n)(t) - (\Phi y_0)(t)| \\ &\leq \int_t^{t+\omega} |G(t, s)| |d^*| \left| \int_{-\infty}^0 K(r) |f(t, y_n(t+r)) - f(t, y_0(t+r))| dr \right| ds \\ &\quad + \rho \int_t^{t+\omega} |G(t, s)| \int_{-\infty}^0 K(r) |f(t, y_n(t+r)) - f(t, y_0(t+r))| dr \\ &\quad + \tilde{\theta} c^* M \int_{-\infty}^0 K(r) |y_n(t+r) - y_0(t+r)| dr. \end{aligned}$$

By Lebesgue dominated convergence theorem and (3.1), we obtain

$$\|(\Phi y_n) - (\Phi y_0)\| \rightarrow 0.$$

This shows that Φ is a continuous on K .

Next, we show that Φ is completely continuous. Let λ be any positive constant and

$$S_\lambda = \{y \in K : \|y\| \leq \lambda\},$$

be a bounded set in K . Then for $y \in K$,

$$\begin{aligned}
 |(\Phi y)(t)| &\leq M \left[\int_0^\omega |d(s)| \left| \int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right| \right. \\
 &\quad \left. + \rho \left(\int_{-\infty}^0 K(r) |c(s)| |y(s+r)| dr \right) ds \right. \\
 &\quad \left. + (\exp(\rho\omega) - 1) \int_{-\infty}^0 K(r) |c(t)| |y(t+r)| dr \right] \\
 &\leq Md^* \int_0^\omega \int_{-\infty}^0 K(r) dr ds \max_{s \in [0, \omega], \|y\| \leq \lambda} |f(s, y)| \\
 &\quad + \rho Mc^* \int_0^\omega \int_{-\infty}^0 K(r) dr ds \max_{s \in [0, \omega]} |y(s)| \\
 &\quad + M\tilde{\theta}c^* \int_{-\infty}^0 K(r) dr \max_{s \in [0, \omega]} |y(s)| \\
 &\leq M\omega d^* \max_{s \in [0, \omega], \|y\| \leq \lambda} |f(s, y)| + \rho\omega Mc^* \lambda + M\lambda\tilde{\theta}c^* \\
 &= U.
 \end{aligned}$$

Therefore, for any $y \in S_\lambda$,

$$\|\Phi u\| \leq U,$$

which implies that $\Phi(S_\lambda)$ is a uniformly bounded set. By taking the derivative in (3.1),

$$\begin{aligned}
 &\frac{d(\Phi y)(t)}{dt} \\
 = &\int_t^{t+\omega} [(\exp(-\rho\omega) - 1) G_1(t, t) G_2(t, s) - b(t) G(t, s) \\
 &+ \int_t^{t+\omega} F(t, \mu) G_1(t, \mu) d\mu] \times \\
 &\times \left[d(s) \int_{-\infty}^s K(r) f(s, y(s+r)) dr - \rho \int_{-\infty}^0 K(r) c(s) y(s+r) dr \right] ds.
 \end{aligned}$$

Consequently, by invoking (2.5), (2.14), (2.15), (2.19), (3.2), we have

$$\begin{aligned} \left| \frac{d(\Phi y)(t)}{dt} \right| &= \int_t^{t+\omega} [|(\exp(-\rho\omega) - 1) G_1(t, t) G_2(t, s) - b(t) G(t, s)| \\ &\quad + \int_t^{t+\omega} |F(t, \mu) G_1(t, \mu)| d\mu] \times \\ &\quad \times \left[\left| d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right| \right. \\ &\quad \left. + \rho \left| \int_{-\infty}^0 K(r) c(s) y(s+r) dr \right| \right] ds \\ &\leq [\omega(\exp(-\rho\omega) - 1) M_1 M_2 + \omega b^* M + \omega M_1 \pi^*] \\ &\quad \times \left[d^* \max_{s \in [0, \omega], \|y\| \leq \lambda} |f(s, y)| + \rho \lambda \right] \\ &= Q, \end{aligned}$$

which implies that $\frac{d(\Phi y)(t)}{dt}$ is also uniformly bounded, for any $y \in S_\lambda$. Hence, $\{\Phi y : y \in K, \|y\| \leq \lambda\}$ is a family of uniformly bounded and equicontinuous functions on $[0, \omega]$. In view of the Arzelà-Ascoli Theorem (see, Royden [30]), operator Φ is completely continuous. The proof of Lemma 3.2 is complete. \square

We can now state and prove our main results.

Theorem 3.1. *In addition to conditions (A1)-(A3), (2.6), (2.10), let us further assume that the following assumptions hold:*

(A4)

$$\liminf_{y \rightarrow 0} \frac{f(t, y)}{y} = \alpha(t), \text{ and } \limsup_{y \rightarrow \infty} \frac{f(t, y)}{y} = \beta(t), \tag{3.3}$$

where α, β are continuous ω -periodic functions on \mathbb{R} .

(A5)

$$\int_0^\omega d(s) \alpha(s) ds \geq \rho \int_0^\omega c(s) ds + \frac{1}{m\delta} - \tilde{\theta} \hat{c}, \tag{3.4}$$

and

$$\int_0^\omega d(s) \beta(s) ds \leq \rho \int_0^\omega c(s) ds + \frac{1}{M} (1 - \tilde{\theta} c^*). \tag{3.5}$$

Then, equation (1.7) possesses at least one positive ω -periodic solution.

Proof We first construct two sets Ω_1 and Ω_2 in order to apply Theorem 1.1. Since $\liminf_{y \rightarrow 0} \frac{f(t, y)}{y} = \alpha(t)$, there exists $R_1 > 0$ such that

$$f(t, y) \geq \alpha(t) y, \text{ for } 0 < y \leq R_1.$$

Define now the open subset

$$\Omega_1 = \{y(\cdot) \in C_\omega : \|y\| < R_1\}.$$

Then, $\partial\Omega_1 = \{y(\cdot) \in C_\omega : \|y\| = R_1\}$, while $y \in K \cap \partial\Omega_1$, that is $\delta \|y\| \leq y \leq \|y\| = R_1$. Noticing that

$$G(t, s) \left(d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right) \geq 0,$$

by noticing inequality (3.4) and the definition of Φ , we obtain

$$\begin{aligned} (\Phi y)(t) &\geq m \int_t^{t+\omega} \left[d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\ &\quad + m\tilde{\theta} \int_{-\infty}^0 K(r) c(t) y(t+r) dr \tag{3.6} \\ &\geq m \int_0^\omega [d(s) \alpha(s) - \rho c(s)] y(s) ds + m\tilde{\theta} c(t) y(t) \\ &\geq m\delta R_1 \int_0^\omega [d(s) \alpha(s) - \rho c(s)] ds + m\delta R_1 \tilde{\theta} \hat{c} \\ &\geq \frac{m\delta R_1}{m\delta} - m\delta R_1 \tilde{\theta} \hat{c} + m\delta R_1 \tilde{\theta} \hat{c} \\ &> R_1 = \|y\|. \end{aligned}$$

Therefore, we conclude that

$$\|\Phi y\| > \|y\| \text{ for } y \in K \cap \partial\Omega_1. \tag{3.7}$$

Next we construct the set Ω_2 . Since $\limsup_{y \rightarrow \infty} \frac{f(t, y)}{y} = \beta(t)$, there exists N such that

$$g(t, y) \leq \beta(t) y, \text{ for } y \geq N.$$

Let

$$R_2 > \max \left\{ \frac{M\theta}{1 - c_2^* \tilde{\theta} - M \int_0^\omega (d(s) \beta(s) - \rho c_1(s)) ds}, N \right\} > R_1,$$

where

$$\theta = \omega (d^* \tilde{m}(f) + \rho c_2^* N), \quad \tilde{m}(f) = \max_{(t,u) \in [0,\omega] \times [0,N]} f(t, y).$$

Define now the open set $\Omega_2 = \{y(\cdot) \in C_\omega : \|y\| < R_2\}$, $E_1 = \{y(\cdot) \in C_\omega : \|y\| < N\}$, $E_2 = \{y(\cdot) \in C_\omega : \|y\| > N\}$,

obviously $\overline{\Omega}_1 \subset \Omega_2$. If $y \in K \cap \partial\Omega_2$, by (3.1), (3.5), (2.19), and (A3), we have

$$\begin{aligned}
 (\Phi y)(t) &\leq M \int_0^\omega \left[d(s) \left(\int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right) \right. \\
 &\quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\
 &\quad + M (\exp(\rho\omega) - 1) \int_{-\infty}^0 K(r) c(s) y(s+r) dr \\
 &\leq M \int_{E_1} \left[d(s) \left(\int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right) \right. \\
 &\quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\
 &\quad + M \int_{E_2} \left[d(s) \left(\int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right) \right. \\
 &\quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds + M\tilde{\theta} \int_{-\infty}^0 K(r) c(s) y(s+r) dr \\
 &\leq M\omega d^* \max_{(s,u) \in [0,\omega] \times [0,N]} f(s, y(s+r)) + M\omega \rho c^* N \\
 &\quad + M \int_{E_2} \left[d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr - \rho \int_{-\infty}^0 K(r) c(s) y(s+r) dr \right] ds \\
 &\quad + M\tilde{\theta} c^* R_2 \\
 &\leq M\omega (d^* \tilde{m}(f) + \rho c^* N) + MR_2 \int_0^\omega [d(s) \beta(s) - \rho c(s)] ds + M\tilde{\theta} c^* R_2 \\
 &\leq M\theta + MR_2 \int_0^\omega [d(s) \beta(s) - \rho c(s)] ds + M\tilde{\theta} c^* R_2 \\
 &< R_2,
 \end{aligned} \tag{3.8}$$

and therefore

$$\|\Phi y\| > \|y\|, \forall y \in K \cap \partial\Omega_2. \tag{3.9}$$

By Lemma 3.2, Φ is a completely continuous operator; from (3.7) and (3.9), condition (i) of Theorem 1.1 is fulfilled. Thus, Φ has at least one fixed point $y \in K \cap (\overline{\Omega}_2 \setminus \Omega_1)$. By Theorem 1.1, we can conclude that Φ has a fixed point $y \in K \cap (\overline{\Omega}_2 \setminus \Omega_1)$ and with $R_1 \leq \|y\| \leq R_2$, and $y \geq \delta \|y\| \geq \delta R_1 > 0$. In other words, equation (1.7) has at least one positive periodic solution. \square

Theorem 3.2. *Suppose that (A1)-(A3), (2.6), (2.10) hold, and the following conditions are fulfilled,*

$$A_0 = \lim_{\|y\| \rightarrow 0} \max_{t \in [0,\omega]} \frac{f(t, y) - \rho \frac{c(t)}{d(t)} y}{\|y\|} = 0, \tag{3.10}$$

$$A_\infty = \lim_{y \in K, \|y\| \rightarrow \infty} \min_{t \in [0,\omega]} \frac{f(t, y) - \rho \frac{c(t)}{d(t)} y}{\|y\|} = \infty. \tag{3.11}$$

Then equation (1.7) has at least one positive ω -periodic solution.

Proof i) In view of (3.10), there exist $\varepsilon > 0$ and corresponding $R_1 > 0$, such that

$$0 < \varepsilon < \frac{1 - \tilde{\theta}Mc^*}{Md^*}, \quad (0 \leq \tilde{\theta}Mc^* < 1),$$

and

$$f(t, y) - \rho \frac{c(t)}{d(t)}y \leq \varepsilon \|y\|, \quad \text{for } \|y\| \leq R_1, t \in [0, \omega].$$

Define the open subset $\Omega_1 = \{y(\cdot) \in C_\omega : \|y\| < R_1\}$, while $y \in K \cap \partial\Omega_1$, that is $\delta \|y\| \leq y \leq \|y\| = R_1$, we can have

$$\begin{aligned} (\Phi y)(t) &= \int_t^{t+\omega} G(t, s) \left[\left(d(s) \int_{-\infty}^0 K(r) f(s, y(s+r)) dr \right) \right. \\ &\quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\ &\quad + (\exp(\rho\omega) - 1) G(t, t) \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right) \\ &\leq M \int_t^{t+\omega} d(s) \int_{-\infty}^0 K(r) \left[f(s, y(s+r)) - \rho \frac{c(s)}{d(s)} y(s+r) \right] dr ds \\ &\quad + \tilde{\theta}M \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right) \\ &\leq Md^* \varepsilon \|y\| + \tilde{\theta}Mc^* \|y\| \\ &\leq \|y\|. \end{aligned}$$

Hence,

$$\|\Phi y\| \leq \|y\|,$$

namely

$$\|\Phi y\| \leq \|y\|, \quad \text{for each } y \in K \cap \partial\Omega_1. \tag{3.12}$$

ii) In view of (3.11), there exist D and corresponding $R_2 > R_1 (\delta)^{-1}$ such that

$$\delta Dm\hat{a} + \delta m\hat{c}\tilde{\theta} \geq 1, f(t, y) - \rho \frac{c(t)}{d(t)}y \geq D \|y\|, \quad \text{for } y \geq \delta \|y\|, \quad \text{and } \|y\| \geq \delta R_2.$$

Define $\Omega_2 = \{y(\cdot) \in C_\omega : \|y\| < R_2\}$, obviously, $\bar{\Omega}_1 \subset \Omega_2$. For $y \in K \cap \partial\Omega_2$, we can have

$$\begin{aligned}
 (\Phi y)(t) &= \int_t^{t+\omega} G(t,s) \left[\left(d(s) \int_{-\infty}^0 K(r) f(s,y(s+r)) dr \right) \right. \\
 &\quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\
 &\quad + (\exp(\rho\omega) - 1) G(t,t) \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right) \\
 &\geq m \int_t^{t+\omega} d(s) \int_{-\infty}^0 K(r) \left[f(s,y(s+r)) - \rho \frac{c(s)}{d(s)} y(s+r) \right] dr ds \\
 &\quad + m\tilde{\theta} \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right) \\
 &\geq m\hat{d}\delta DR_2 + m\tilde{c}\tilde{\theta}\delta R_2 \\
 &\geq (m\hat{d}D\delta + m\tilde{c}\tilde{\theta}\delta) R_2 \\
 &\geq R_2 = \|y\|,
 \end{aligned}$$

which leads to

$$(\Phi y)(t) \geq \|y\|, \text{ for each } y \in K \cap \partial\Omega_2. \tag{3.13}$$

By Lemma 3.2, Φ is a completely continuous operator; from (3.12) and (3.13), condition (i) of Theorem 1.1 is fulfilled. Thus, Φ has at least one fixed point $y \in K \cap (\bar{\Omega}_2 \setminus \Omega_1)$, and $y \geq \delta \|y\| \geq \delta R_1 > 0$ which means $y(\cdot)$ is a ω -periodic positive solution of (1.7). This completes the proof. \square

For convenience and simplicity in the following discussion, we use the following notations:

$$F_{R_2}^0 = \inf_{y \in K \cap \partial\Omega_2} \min_{t \in [0,\omega]} \left(\frac{f(t,y)}{y} - \rho \frac{c(t)}{d(t)} \right), \tag{3.14}$$

$$F_{R_1}^\infty = \sup_{y \in K \cap \partial\Omega_1} \max_{t \in [0,\omega]} \left(\frac{f(t,y)}{y} - \rho \frac{c(t)}{d(t)} \right). \tag{3.15}$$

Theorem 3.3. *Suppose that (A1)-(A3), (2.6), (2.10) hold, and there are positive constants R_1, R_2 with $R_1 < R_2$ such that:*

$$M \left(d^* F_{R_1}^\infty + \tilde{\theta} c^* \right) \leq 1, \tag{3.16}$$

and

$$m \left(\hat{d} F_{R_2}^0 + \tilde{\theta} \hat{c} \right) \geq 1. \tag{3.17}$$

Then, there exists an ω -periodic solution which is a fixed point of Φ and satisfies $R_1 \leq \|y\| \leq R_2$.

Proof Define two open sets Ω_1 and Ω_2 with $R_1 < R_2$. Let $\Omega_1 = \{y(\cdot) \in C_\omega : \|y\| < R_1\}$. Then for any $y \in K \cap \partial\Omega_1$, we have $\delta \|y\| \leq y \leq \|y\| = R_1$. From this, the definition of Φ and $F_{R_1}^\infty$, it follows that

$$\begin{aligned}
 (\Phi y)(t) &= \int_t^{t+\omega} G(t,s) \left[\left(d(s) \int_{-\infty}^0 K(r) f(s,y(s+r)) dr \right) \right. \\
 &\quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\
 &\quad + (\exp(\rho\omega) - 1) G(t,t) \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right) \\
 &\leq M \int_t^{t+\omega} d(s) \int_{-\infty}^0 K(r) \left[f(s,y(s+r)) - \rho \frac{c(s)}{d(s)} y(s+r) \right] dr ds \\
 &\quad + \tilde{\theta} M \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right) \\
 &\leq \left(Md^* F_{R_1}^\infty + \tilde{\theta} M c^* \right) \|y\|. \tag{3.18}
 \end{aligned}$$

Hence, in view of (3.16) and (3.18), we obtain

$$\|\Phi y\| \leq \|y\|, \text{ for each } y \in K \cap \partial\Omega_2. \tag{3.19}$$

On the other hand, let $\Omega_2 = \{y(\cdot) \in C_\omega : \|y\| < R_2\}$, obviously, $\bar{\Omega}_1 \subset \Omega_2$. For $y \in K \cap \partial\Omega_2$, then $\delta \|y\| \leq y \leq \|y\| = R_2$. From this, the definition of Φ and $F_{R_2}^0$, it follows that

$$\begin{aligned}
 (\Phi y)(t) &= \int_t^{t+\omega} G(t,s) \left[\left(d(s) \int_{-\infty}^0 K(r) f(s,y(s+r)) dr \right) \right. \\
 &\quad \left. - \rho \left(\int_{-\infty}^0 K(r) c(s) y(s+r) dr \right) \right] ds \\
 &\quad + (\exp(\rho\omega) - 1) G(t,t) \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right) \\
 &\geq m \int_t^{t+\omega} d(s) \int_{-\infty}^0 K(r) \left[f(s,y(s+r)) - \rho \frac{c(s)}{d(s)} y(s+r) \right] dr ds \\
 &\quad + m\tilde{\theta} \left(\int_{-\infty}^0 K(r) c(t) y(t+r) dr \right) \\
 &\geq \left(m\hat{d}F_{R_1}^0 + m\hat{c}\tilde{\theta} \right) \|y\|. \tag{3.20}
 \end{aligned}$$

Hence, from (3.17) and (3.20), we obtain

$$(\Phi y)(t) \geq \|y\|, \text{ for each } y \in K \cap \partial\Omega_1. \tag{3.21}$$

By Lemma 3.2, Φ is a completely continuous operator, from (3.19) and (3.21), condition (i) of Theorem 1.1 is fulfilled. Thus, Φ has at least one fixed point $y \in K \cap (\bar{\Omega}_2 \setminus \Omega_1)$. It follows that $y(\cdot)$ is an ω -periodic positive solution of (1.7) and $y \geq \delta \|y\| \geq \delta R_1 > 0$. This completes the proof. \square

As consequence of Theorem 3.3, we state a corollary whose proof is similar to the proof of Theorem 3.3 and hence we omit it.

Corollary 3.1. *Suppose that (A1)-(A3), (2.6), (2.10) hold, and there are positive constants R_1, R_2 with $R_1 < R_2$ such that:*

$$M \left(d^* F_{R_2}^0 + \tilde{\theta} c^* \right) \leq 1, \tag{3.22}$$

and

$$m \left(\widehat{d}F_{R_1}^\infty + \tilde{\theta} \widehat{c} \right) \geq 1. \tag{3.23}$$

Then, there exists an ω – periodic solution which is a fixed point of Φ and satisfies $R_1 \leq \|y\| \leq R_2$.

As a consequence of Theorem 3.3, we have the next Corollary.

Corollary 3.2. *Suppose that (A1)-(A3), (2.6), (2.10) hold, and there are positive constants R_1, R_2, \dots, R_{n+1} with $R_1 < R_2 < \dots < R_{n+1}$ such that:*

$$M \left(d^* F_{R_2}^\infty + \tilde{\theta} c^* \right) \leq 1,$$

$$m \left(\widehat{d}F_{R_1}^0 + \widehat{c} \tilde{\theta} \right) \geq 1,$$

$$M \left(d^* F_{R_4}^\infty + \tilde{\theta} c^* \right) \leq 1,$$

$$m \left(\widehat{d}F_{R_3}^0 + \widehat{c} \tilde{\theta} \right) \geq 1,$$

.....

$$M \left(d^* F_{R_{n+1}}^\infty + \tilde{\theta} c^* \right) \leq 1,$$

$$m \left(\widehat{d}F_{R_n}^0 + \widehat{c} \tilde{\theta} \right) \geq 1.$$

Then, Eq. (1.7) has n positive ω –periodic solutions y_1, y_2, \dots, y_n with $\|y_1\| \leq \|y_2\| \leq \dots \leq \|y_n\|$.

Remark 3.1. In our main results, condition (A3) is very important to guarantee that operator Φ maps K into itself.

4. Examples

Let us discuss three examples to illustrate our abstract theory.

Example 4.1. Let us consider the following third-order nonlinear integro-differential delay equation:

$$\begin{aligned} & y'''(t) + p(t)y''(t) + q(t)y'(t) + r(t)y(t) \\ &= \frac{d}{dt} \int_{-\infty}^0 K(r) c(t) y(t+r) dr \\ &+ d(t) \int_{-\infty}^0 K(r) f(t, y(t+r)) dr. \end{aligned} \tag{4.1}$$

Corresponding to equation (1.7), we let

$$\begin{aligned} p(t) &= 31 \times 10^{-2}, q(t) = 23 \times 10^{-3}, \omega = 2\pi, \\ c(t) &= 1 + 0.1 |\sin t|, r(t) = 2 \times 10^{-4}. \end{aligned}$$

Also assume that

$$f(t, y) = \frac{0.4}{d(t)} \frac{y}{1+y} + \rho \frac{c(t)}{d(t)} y,$$

where $d \in C(\mathbb{R}, \mathbb{R}^+ \setminus \{0\})$ is an arbitrary 2π -periodic function.

Moreover, $K \in C(\mathbb{R}^-, \mathbb{R}^+)$ is an arbitrary function satisfying

$$\int_{-\infty}^0 K(r) dr = 1.$$

Thanks to direct computations, we obtain

$$\begin{aligned} a(t) &= 0.2, b(t) = 0.01, \delta = \frac{m}{M} \simeq 0.151, \\ a_1(t) &= 21 \times 10^{-2}, a_2(t) = 2 \times 10^{-3}, \rho = 0.1, \\ c^* &= 1.1, \widehat{c} = 1. \end{aligned}$$

It is clear that, $\forall y \in \mathbb{R}^+, t \in [0, 2\pi]$, we have

$$f(t, y) - \rho \frac{c(t)}{d(t)} y = \frac{0.4}{d(t)} \frac{y}{1+y} \geq 0.$$

Again by straightforward computations, we have

$$\liminf_{y \rightarrow 0} \frac{f(t, y)}{y} = \frac{0.4}{d(t)} + \rho \frac{c(t)}{d(t)} = \alpha(t),$$

and

$$\limsup_{y \rightarrow \infty} \frac{f(t, y)}{y} = \rho \frac{c(t)}{d(t)} = \beta(t).$$

It is easy to check α, β are continuous 2π -periodic functions on \mathbb{R} .

From the above parameters, it follows that

$$\begin{aligned} &\int_0^\omega d(s) \alpha(s) ds - \int_0^\omega \rho c(s) ds - \frac{1}{m\delta} + (\exp(\rho\omega) - 1) \widehat{c} \\ &= 0.4\omega - \frac{1}{m\delta} + (\exp(\rho\omega) - 1) \widehat{c} \\ &\simeq 2.6 \geq 0. \end{aligned}$$

and

$$\begin{aligned} &\int_0^\omega d(s) \beta(s) ds - \rho \int_0^\omega c(s) ds - \frac{1}{M} (1 - (\exp(\rho\omega) - 1) c^*) \\ &= -\frac{1}{M} (1 - (\exp(\rho\omega) - 1) c^*) \\ &\simeq -54 \times 10^{-8} \leq 0. \end{aligned}$$

Consequently, all conditions of Theorem 3.1 are fulfilled. Hence, we conclude that Equation (4.1) possesses at least one positive 2π -periodic solution $y \in K \cap (\overline{\Omega}_2 \setminus \Omega_1)$.

Example 4.2. Let us consider the following third-order nonlinear integro-differential delay equation:

$$\begin{aligned} & y'''(t) + p(t)y''(t) + q(t)y'(t) + r(t)y(t) \\ &= \frac{d}{dt} \int_{-\infty}^0 K(r)c(t)y(t+r)dr \\ &+ d(t) \int_{-\infty}^0 K(r)f(t,y(t+r))dr. \end{aligned} \tag{4.2}$$

Corresponding to equation (1.7), we let

$$\begin{aligned} p(t) &= 31 \times 10^{-2}, q(t) = 23 \times 10^{-3}, \omega = 2\pi, \\ c(t) &= 4|\cos t| + 2, r(t) = 2 \times 10^{-4}, d(t) = 0.2|\cos t| + 0.1. \end{aligned}$$

Also assume that

$$f(t, y) = 0.4y^2 + 2y.$$

Moreover, $K \in C(\mathbb{R}^-, \mathbb{R}^+)$ is an arbitrary function satisfying

$$\int_{-\infty}^0 K(r)dr = 1.$$

Thanks to direct computations, we obtain

$$\begin{aligned} a(t) &= 0.2, b(t) = 0.01, \delta = \frac{m}{M} \simeq 0.151, \\ a_1(t) &= 21 \times 10^{-2}, a_2(t) = 2 \times 10^{-3}, \rho = 0.1, \\ c^* &= 6, \hat{c} = 2. \end{aligned}$$

It is clear that, $\forall y \in \mathbb{R}^+, t \in [0, 2\pi]$, we have

$$f(t, y) - \rho \frac{c(t)}{d(t)}y = 0.4y^2 \geq 0.$$

It is easy to verify that

$$\begin{aligned} A_0 &= \lim_{\|y\| \rightarrow 0} \max_{t \in [0, \omega]} \frac{f(t, y) - \rho \frac{c(t)}{d(t)}y}{\|y\|} \\ &= \lim_{\|y\| \rightarrow 0} \max_{t \in [0, \omega]} \frac{0.4y^2}{\|y\|} = 0, \end{aligned}$$

and

$$\begin{aligned} A_\infty &= \lim_{y \in K, \|y\| \rightarrow \infty} \min_{t \in [0, \omega]} \frac{f(t, y) - \rho \frac{c(t)}{d(t)}y}{\|y\|} \\ &= \lim_{y \in K, \|y\| \rightarrow \infty} \min_{t \in [0, \omega]} \frac{0.4y^2}{\|y\|} = \infty. \end{aligned}$$

It is straightforward to show that all conditions of Theorem 3.2 are fulfilled. Hence, we conclude that equation (4.2) possesses at least one positive 2π -periodic solution $y \in K \cap (\bar{\Omega}_2 \setminus \Omega_1)$.

Example 4.3. Let us consider the following equation:

$$\begin{aligned} & y'''(t) + p(t)y''(t) + q(t)y'(t) + r(t)y(t) \\ &= \frac{d}{dt} \int_{-\infty}^0 K(r) c(t) y(t+r) dr \\ &+ d(t) \int_{-\infty}^0 K(r) f(t, y(t+r)) dr. \end{aligned} \tag{4.3}$$

Corresponding to equation (1.7), we let

$$\begin{aligned} p(t) &= 31 \times 10^{-2}, q(t) = 23 \times 10^{-3}, r(t) = 2 \times 10^{-4}, \omega = 2\pi, \\ d(t) &= 10^{-4} (0.2 + 0.2 |\sin t|), c(t) = 10^{-5} (1 + 0.5 |\cos t|). \end{aligned}$$

Also assume that

$$f(t, y) = \frac{0.1 (1 + 10^2 \times 0.151^2 |\sin t|)}{0.001 + y^2} y + \frac{1}{100} \times \frac{1 + 0.5 |\cos t|}{0.2 + 0.2 |\sin t|} y,$$

and $K \in C(\mathbb{R}^-, \mathbb{R}^+)$ satisfying

$$\int_{-\infty}^0 K(r) dr = 1.$$

Then we can check that

$$\begin{aligned} a(t) &= 0.2, b(t) = 0.01, \rho = 0.1, \\ a_1(t) &= 21 \times 10^{-2}, a_2(t) = 2 \times 10^{-3}, \\ m &\simeq 10735.730, M \simeq 70706.185, \\ \delta &= \frac{m}{M} \simeq 0.151. \end{aligned}$$

It is clear that, $\forall y \in \mathbb{R}^+, t \in [0, 2\pi]$, we have

$$f(t, y) - \rho \frac{c(t)}{d(t)} y = \frac{0.1 (1 + 10^2 \times 0.151^2 |\sin t|) y}{0.001 + y^2} \geq 0.$$

There are positive constants $R_1 = 0.02, R_2 = 10$, such that

$$\begin{aligned} F_{R_1}^0 &= \inf_{y \in K \cap \delta\Omega_1} \min_{t \in [0, \omega]} \left(\frac{f(t, y)}{y} - \rho \frac{c(t)}{d(t)} \right) \\ &= \inf_{y \in K \cap \delta\Omega_1} \min_{t \in [0, \omega]} \left(\frac{0.1 (1 + 10^2 \delta^2 |\sin t|)}{0.001 + y^2} \right) \\ &= \inf_{y \in K \cap \delta\Omega_1} \left(\frac{0.1}{0.001 + \|y\|^2} \right) \\ &= \frac{0.1}{0.001 + 0.0004} \simeq 71.428. \end{aligned}$$

$$\begin{aligned}
 F_{R_2}^\infty &= \sup_{y \in K \cap \delta\Omega_2} \max_{t \in [0, \omega]} \left(\frac{f(t, y)}{y} - \rho \frac{c(t)}{d(t)} \right) \\
 &= \sup_{y \in K \cap \delta\Omega_2} \max_{t \in [0, \omega]} \left(\frac{0.1 (1 + 10^2 \delta^2 |\sin t|)}{0.001 + y^2} \right) \\
 &= \sup_{y \in K \cap \delta\Omega_2} \frac{0.1 (1 + 10^2 \delta^2)}{0.001 + y^2} \\
 &= \frac{0.1 (1 + 10^2 \delta^2)}{0.001 + \delta^2 R_2^2} \simeq 0.143.
 \end{aligned}$$

Now, some simple calculations show that $\forall (t, s) \in \mathbb{R} \times \mathbb{R}$,

$$10735.730 \simeq m \leq G(t, s) \leq M \simeq 70706.185.$$

For the above parameters, it is easy to verify that

$$M \left(d^* F_{R_2}^\infty + \tilde{\theta} c^* \right) \simeq 0.713 \leq 1,$$

and

$$m \left(\widehat{d} F_{R_1}^0 + \widehat{c} \tilde{\theta} \right) \simeq 15.430 \geq 1,$$

where

$$\begin{aligned}
 d^* &= \max_{t \in [0, \omega]} d(t) = 0.4 \times 10^{-4}, \widehat{d} = \min_{t \in [0, \omega]} d(t) = 0.2 \times 10^{-4}, \\
 c^* &= \max_{t \in [0, \omega]} c(t) = 0.5 \times 10^{-5}, \widehat{c} = \min_{t \in [0, \omega]} c(t) = 10^{-5}, \\
 \tilde{\theta} &= \exp(\rho\omega) - 1 \simeq 0.874.
 \end{aligned}$$

All hypotheses of Theorem 3.3 are fulfilled and, therefore, Eq. (4.3) has at least one positive 2π -periodic solution $y \in K \cap (\overline{\Omega_2} \setminus \Omega_1)$ satisfying

$$0.2 = R_1 \leq \|y\| \leq R_2 = 10.$$

Remark. 4.2. Observe that Example 4.1 and Example 4.2 cannot be analyzed by applying Theorem 3.1 in [25] (see also Theorem 3.1 in [2]). Indeed, in order to apply Theorem 3.1 in [2] or [25], we need to construct two mappings Φ_1 and Φ_2 , Φ_1 is a contraction and Φ_2 is compact. Therefore, we express (4.1) or (4.2) as

$$(\Phi x)(t) = (\Phi_1 x)(t) + (\Phi_2 x)(t),$$

where $\Phi_1, \Phi_2 : C_\omega \rightarrow C_\omega$ are given by

$$(\Phi_1 x)(t) = (\exp(\rho\omega) - 1) G(t, t) \int_{-\infty}^0 K(r) c(s) x(s+r) dr,$$

$$\begin{aligned}
(\Phi_2 x)(t) &= \int_t^{t+\omega} G(t, s) \left[d(s) \int_{-\infty}^0 K(r) f(s, x(s+r)) dr \right. \\
&\quad \left. - \rho \int_{-\infty}^0 K(r) c(s) x(s+r) dr \right] ds.
\end{aligned}$$

However, notice that

$$\exists \gamma > 1, \forall x, y \in C_\omega, \|\Phi_1 x - \Phi_1 y\| \leq \gamma \|x - y\|,$$

where

$$\gamma = M (\exp(\rho\omega) - 1) c^*.$$

This implies that Φ_1 is not contraction. Thus, Theorem 3.1 in [2] and [25] cannot be applied to equations (4.1) and (4.2). Therefore, the results in [2] and [25] are not applicable. However, the results obtained in our work are quite significant compared to the ones in the aforementioned papers [2, 25].

Acknowledgments

The research has been partially supported by the Spanish Ministry of Science and Innovation (MCI), State Investigation Agency (AEI) and European Regional Development Fund (FEDER) under the project PID2021-122991NB-C21, and by Regional Government of Andalusia (Department of Economy and Knowledge) and FEDER under project P18-FR-4509.

Conflict of interests

The authors declare that they do not have any conflict of interest.

References

- [1] Abou-El-Ela AMA, Sadek AI, Mahmoud AM. Periodic solutions for a kind of third-order delay differential equations with a deviating argument. *Journal of Mathematical Sciences, The University of Tokyo*. 2011; 18 (1): 35-49.
- [2] Ardjouni A, Djoudi A. Existence of positive periodic solutions for third-order nonlinear delay differential equations with variable coefficients. *Mathematica Moravica* 2019; 23 (2): 17-28. <https://doi.org/10.5937/MatMor1902017A>
- [3] Ardjouni A, Djoudi A. Existence of positive periodic solutions for a nonlinear neutral differential equations with variable delay. *Applied Mathematics E-Note* 2012; 12: 94-101.
- [4] Balamuralithran S. Periodic solutions for third-order nonlinear delay equation impulses with Fredholm operator of index operator zero. *Konuralp Journal of Mathematics* 2016; 4 (2): 158-168.
- [5] Bose SK, Gorain GC. Exact controllability and boundary stabilization of torsional vibrations of an internally damped flexible space structure. *Journal of Optimization Theory and Applications* 1998; 99 (2): 423-442. <https://doi.org/10.1023/A:1021778428222>
- [6] Bose SK, Gorain GC. Exact controllability and boundary stabilization of flexural vibrations of an internally damped flexible space structure. *Applied Mathematics and Computation* 2002; 126 (2-3): 341-360. [https://doi.org/10.1016/S0096-3003\(00\)00112-0](https://doi.org/10.1016/S0096-3003(00)00112-0)
- [7] Bose SK, Gorain GC. Uniform stability of damped nonlinear vibrations of an elastic string. In: *Mathematical Sciences-Proceedings of the Indian Academy of Sciences; India; 2003*. pp. 443-449. <https://doi.org/10.48550/arXiv.math/0311527>

- [8] Cheng Z, Ren J. Existence of positive periodic solution for variable coefficient third-order differential equation with singularity. *Mathematical Methods in the Applied Sciences* 2014; 37 (15): 2281-2289. <https://doi.org/10.1002/mma.2975>
- [9] Cheng Z, Xin Y. Multiplicity results for variable-coefficient singular third-order differential equation with a parameter. *Abstract and Applied Analysis* 2014; 2014 (1): 1-10. <https://doi.org/10.1155/2014/527162>
- [10] Cai G, Bu S. Periodic solutions of third-order integro-differential equations in vector-valued functional spaces. *Journal of Evolution Equations* 2017; 17 (2): 749-780. <https://doi.org/10.1007/s00028-016-0335-5>
- [11] Cheng Z, Lv L, Liu J. Positive periodic solution of first-order neutral differential equation with infinite distributed delay and applications. *AIMS Mathematics* 2020; 5 (6): 7372-7386. <https://doi.org/10.3934/math.2020472>
- [12] Gregus M. *Third Order Linear Differential Equations. Series, Mathematics and its applications*, Dordrecht, Boston, Lancaster, Tokyo: Reidel Publishing Company, 1987. <https://doi.org/10.1007/978-94-009-3715-4>
- [13] Guo D, Lakshmikantham V. *Nonlinear Problems in Abstract Cones*. Academic Press, New York, USA: Elsevier, 1988.
- [14] Gorain GC. Exponential energy decay estimate for the solutions of internally damped wave equation in a bounded domain. *Journal of Mathematical Analysis and Applications* 1997; 216 (2): 510-520. <https://doi.org/10.1006/jmaa.1997.5678>
- [15] Gorain GC. Boundary stabilization of nonlinear vibrations of a flexible structure in a bounded domain in \mathbb{R}^n . *Journal of Mathematical Analysis and Applications* 2006; 319 (2): 635-650. <https://doi.org/10.1016/j.jmaa.2005.06.031>
- [16] Hale JK, Meyer KR. A class of functional equations of neutral type. *Memoirs of the American Mathematical Society* 1967; 76: 1-65. <https://doi.org/http://dx.doi.org/10.1090/memo/0076>
- [17] Hale JK, Lunel SMV. *Introduction to Functional Differential Equations*. Applied Mathematical Sciences, New York, USA: Springer, 1993. <https://doi.org/10.1007/978-1-4612-4342-7>
- [18] Jiang D, Wei J, Zhang B. Positive periodic solutions of functional differential equations and population models. *Electronic Journal of Differential equations* 2002; 2002 (71): 1-13.
- [19] Kuang Y. *Delay Differential Equations with Application in Population Dynamics*. Boston, New York, USA: Academic Press, 1993.
- [20] Liu B, Huang L. Existence and uniqueness of periodic solution for a kind of first order neutral functional differential equations. *Journal of Mathematical Analysis and Applications* 2006; 322 (1): 121-132. <https://doi.org/10.1016/j.jmaa.2005.08.069>
- [21] Liu Y, Ge W. On the positive periodic solutions of nonlinear Duffing equations with delay and variable coefficients. *Tamsui Oxford Journal of Information and Mathematical Sciences* 2004; 20 (2): 235-255. <https://doi.org/10.36045/bbms/1093351383>
- [22] Li Z, Wang X. Existence of positive periodic solutions for neutral functional differential equations. *Electronic Journal of Differential equations* 2006; 2006 (34): 1-8.
- [23] Li J, Shen J. On positive periodic solutions to impulsive differential equations with delays. *Results in Mathematics* 2004; 45: 67-78. <https://doi.org/10.1007/BF03322998>
- [24] Meng Q, Yan J. Existence and n - multiplicity of positive periodic solutions for impulsive functional differential equations with two parameters. *Boundary Value Problems* 2015; 2015 (212): 1-10. <https://doi.org/10.1186/s13661-015-0478-2>
- [25] Nouioua F, Ardjouni A, Djoudi A. Periodic solutions for a third-order delay differential Equation. *Applied Mathematics E-Notes* 2016; 2016 (16): 210-221.
- [26] Poblete V, Pozo JC. Periodic solutions of an abstract third-order differential equation. *Studia Mathematica* 2013; 215 (3): 195-219. <http://doi.org/10.4064/sm215-3-1>

- [27] Padhi S, Pati S. Theory of third-order differential equations. New York Dordrecht London: Springer New Delhi Heidelberg, 2014. <https://doi.org/10.1007/978-81-322-1614-8>
- [28] Ren J, Siegmund S, Chen Y. Positive periodic solutions for third-order nonlinear differential equations. *Electronic Journal of Differential Equations* 2011; 2011 (66): 1-19.
- [29] Raffoul YN, Tisdell CC. Positive periodic solutions of functional discrete systems and population models. *Advances in Difference Equations* 2005; 2005 (3): 369-380. <https://doi.org/10.1155/ADE.2005.369>
- [30] Royden HL. *Real Analysis*. Stanford University, New York, USA: MacMillan Publishing Company, 1998.
- [31] Tiriyaki A, Aktas MF. Oscillation criteria of a certain class of third order nonlinear delay differential equations with damping. *Journal of Mathematical Analysis and Applications* 2007; 325 (1): 54-68. <https://doi.org/10.1016/j.jmaa.2006.01.001>
- [32] Wang Y, Lian H, Ge W. Periodic solutions for a second order nonlinear functional differential equation. *Applied Mathematics Letters* 2007; 20 (1): 110-115. <https://doi.org/10.1016/j.aml.2006.02.028>
- [33] Zhang X, Jian D, Li X, Wang K. A new existence theory for single and multiple positive periodic solutions to integro-differential equations with impulse effects. *Computers and Mathematics with Applications* 2006; 51 (1): 17-32. <https://doi.org/10.1016/j.camwa.2005.09.002>