

1-1-2008

The Effect of Erythropoietin on Neurotrophic Factors in N9 Murine Microglial Cells

FİLİZ KURALAY

BAŞAK BİNGÖL ÇAKIRLI

ŞERMİN GENÇ

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

KURALAY, FİLİZ; ÇAKIRLI, BAŞAK BİNGÖL; and GENÇ, ŞERMİN (2008) "The Effect of Erythropoietin on Neurotrophic Factors in N9 Murine Microglial Cells," *Turkish Journal of Medical Sciences*: Vol. 38: No. 6, Article 3. Available at: <https://journals.tubitak.gov.tr/medical/vol38/iss6/3>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

The Effect of Erythropoietin on Neurotrophic Factors in N9 Murine Microglial Cells*

Filiz KURALAY¹

Başak BİNGÖL ÇAKIRLI¹

Şermin GENÇ²

Aim: In this study, we investigated whether interferon gamma (IFN γ), lipopolysaccharides (LPS) and amyloid beta (AMY β), as toxic stimulator agents, and erythropoietin (EPO), as a neurotrophic agent, have an effect on the production of the following neurotrophic factors in the N9 murine microglia cell line: neurotrophin 3 (NT3), neurotrophin 4 (NT4), and brain-derived neurotrophic factor (BDNF).

Materials and Methods: Microglial cells were incubated with 50 μ g/ml AMY β , or 1 μ g/ml of LPS plus 100 U/ml recombinant murine IFN γ , and/or one of three different concentrations (0.1, 1.0, and 5.0 U/ml) of recombinant mouse EPO for 24 h.

Results: EPO 0.1 U/ml dose significantly increased NT4 levels compared to EPO 5.0 U/ml dose ($P < 0.05$). EPO, in all doses, and AMY β significantly induced NT4 secretion in microglia, while BDNF and NT3 were not changed by AMY β or EPO. LPS + IFN γ alone did not change neurotrophic factor levels in any group. However, EPO with LPS and IFN γ induced NT4 secretion, especially the 5.0 U/ml dose of EPO.

Conclusions: NT4 secretion, which was markedly induced by exposure to both AMY β and EPO in N9 murine microglia, may be an important result for neuronal survival. These results suggest that inflammatory mechanisms in microglia may also involve the neuroprotective response of these cells; this may be a promising area of study of neurodegenerative processes.

Key Words: Microglia, N9 cell line, murine, erythropoietin, neurotrophic factors

¹ Department of Biochemistry,
Faculty of Medicine,
Dokuz Eylül University,
Balçova, İzmir - TURKEY

² Learning Resources Center Research
Laboratory, Faculty of Medicine,
Dokuz Eylül University,
Balçova, İzmir - TURKEY

N9 Fare Mikroglial Hücrelerinde Nörotrofik Faktörler Üzerine Eritropoetin Etkisi

Amaç: Bu çalışmada N9 murin mikroglia hücre hattında toksik uyarıcı ajanlar olarak interferon gama (IFN γ), lipopolisakkarid (LPS), amiloid beta (AMY β) ve nörotrofik ajan olarak eritropoetin (EPO)'ün nörotrofik faktörler olan nörotrofin 3 (NT3), nörotrofin 4 (NT4) ve beyin kökenli nörotrofik faktör (BDNF) üretimi üzerine etkisi olup olmadığını araştırdık.

Yöntem ve Gereç: Mikroglial hücreler 50 μ g/ml AMY β veya 1 μ g/ml of LPS + 100 U/ml IFN γ ve/veya EPO'nun üç ayrı konsantrasyonu (0.1, 1.0 ve 5.0 U/ml) ile 24 saat inkübe edildi.

Bulgular: EPO'nun 0.1 U/ml dozu, EPO 5.0 U/ml dozuna göre, NT4 düzeylerini daha anlamlı olarak arttırmıştır ($P < 0.05$) EPO'nun her üç dozu ve AMY β mikroglial NT4 sekresyonunu anlamlı olarak indüklerken; BDNF ve NT3 düzeyleri AMY β veya EPO ile değişmedi. LPS + IFN γ uygulaması tek başına herhangi bir grupta nörotrofik faktör düzeylerini değiştirmedi. Fakat, EPO ile birlikte LPS + IFN γ uygulaması, NT4 salınımını, özellikle EPO'nun 5.0 U/ml dozunda arttırdı.

Sonuç: N9 murin mikroglia hücre hattında AMY β veya EPO uygulamasının NT4 salınımını indüklemesi, nöronal hayatta kalım için önemli bir sonuç olabilir. Bu bulgular, mikroglia'da inflamatuvar mekanizmaların bu hücrelerin nöroprotektif cevabını da kapsayan; belki de nörodejeneratif olaylarla ilgili çalışmalara da ışık tutacak bir alan olabileceği yönündedir.

Anahtar Sözcükler: Mikroglia, N9 hücre hattı, murin, eritropoetin, nörotrofik faktörler

Received: August 20, 2007
Accepted: August 27, 2008

Correspondence

Filiz KURALAY

Department of Biochemistry,
Faculty of Medicine,
Dokuz Eylül University,
35340 Balçova, İzmir - TURKEY

filiz.kuralay@deu.edu.tr

* This study was supported by a Research Foundation Grant (Project no: 923 20 01 09) from Dokuz Eylül University.

Introduction

A glia-mediated, inflammatory immune response is an important component of the neuropathophysiology of neurodegenerative diseases such as Alzheimer's disease (AD) and Parkinson's disease (PD). Activated microglia can mediate neuronal cell death and the repair processes of central nervous system (CNS) injuries. Recent studies suggest that the activation of microglia in response to injury, illness, aging, or other causes begins a cascade of events that can best be characterized as an inflammatory process. This cascade is mediated at first by the proinflammatory cytokine interleukin 1, which is overexpressed by the activated microglia. Microglia produce and release certain growth factors that are essential for neuron survival, such as erythropoietin (EPO) and interferon gamma ($IFN\gamma$) (1,2). Amyloid deposition, inflammation and regenerative mechanisms are also early pathogenic events in experimental models (3,4).

Erythropoietin (EPO) was first characterized as a hematopoietic growth factor and has been in clinical use for over a decade for the treatment of anemia. EPO is currently named "NeoRecormon", and since its receptor was found in glial cells, it was assumed to have beneficial effects on neuronal cells (5-7). The neuroprotective effects of EPO have been demonstrated *in vitro* and *in vivo* in a variety of CNS injury models, such as focal cerebral ischemia and chemical-induced neurotoxicity (8-15). In CNS cell culture studies, EPO was shown to be involved in the regulation of intracellular calcium and signal mechanisms; the enhancement of calcium-dependent or nondependent neurotransmitter release may explain its neuroprotective efficacy. EPO has also been proposed to be an antioxidant, anti-inflammatory and trophic agent in the CNS due to its stimulation of neuronal supportive cells and microglia and it increases their viability (12-14,16). Another possibility is that EPO combined with another cytokine such as $IFN\gamma$ and lipopolysaccharides (LPS) or various neuroactive drugs stimulate glial cells maintaining support for neurons by affecting neurotrophic factors and nitric oxide (NO) production (8,9,15). In our previous studies, we demonstrated that EPO increased antioxidant enzyme activity and NO production in the substantia nigra in a model of methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced parkinsonism in C57BL mice (10,11). Neurotrophic factors, such as nerve growth factor (NGF),

neurotrophin 3 (NT3), neurotrophin 4 (NT4), and brain-derived neurotrophic factor (BDNF), in addition to their neurotrophic actions, may also regulate microglial dynamics, thereby influencing the surrounding milieu during neuronal inflammation and regeneration. In this study, we investigated whether microglial activation with $IFN\gamma$ + LPS or amyloid beta ($AMY\beta$) as toxic stimulator agents, and with EPO as a neurotrophic agent, has an effect on the production of neurotrophic factors (BDNF, NT3 and NT4) in an N9 murine microglial cell line.

Materials and Methods

Microglial Cultures

The N9 murine microglia cell line was kindly provided by Paola Ricciardi-Castagnoli (Cellular Pharmacology Center, Milan, Italy) and maintained with Roswell Park Memorial Institute (RPMI) medium supplemented with 5% fetal calf serum, 100 U/ml penicillin and 100 mg/ml streptomycin. The purity of the microglia cultures was assessed by immunostaining with monoclonal antibody against CD11b/c (Invitrogen, Caltag Laboratories, California, USA), and more than 90% of the cells were positively stained.

Microglial cells were incubated with 50 mg/ml $AMY\beta$ (Sigma-Aldrich, Taufkirchen, Germany), or 1 mg/ml LPS (from *Escherichia coli*; Sigma-Aldrich, Taufkirchen, Germany) plus 100 U/ml recombinant murine $IFN\gamma$ (Roche Diagnostics, Basel, Switzerland) and/or one of three different concentrations (0.1, 1.0, and 5.0 U/ml) of recombinant mouse EPO (Roche Diagnostics, Basel, Switzerland) for 24 hours (h). None of the cytokines was added to the control culture. In summary, the nine treatment groups (three cell lines in each group) were as follows:

- 1) Control
- 2) LPS (1 mg/ml) + $IFN\gamma$ (100 U/ml)
- 3) $AMY\beta$ (50 mg/ml)
- 4) EPO (0.1 U/ml)
- 5) EPO (1.0 U/ml)
- 6) EPO (5.0 U/ml)
- 7) LPS (1 mg/ml) + $IFN\gamma$ (100 U/ml) + EPO (0.1 U/ml)
- 8) LPS (1 mg/ml) + $IFN\gamma$ (100 U/ml) + EPO (1.0 U/ml)
- 9) LPS (1 mg/ml) + $IFN\gamma$ (100 U/ml) + EPO (5.0 U/ml)

Neurotrophic Factor Determinations

After incubation for 24 h, BDNF, NT3 and NT4 levels were duplicatedly measured by ELISA methods (Promega Inc, Madison, WI, USA). Microplates were coated with polyclonal antibodies specific to the neurotrophic factor to be analyzed, which were diluted in carbonate coating buffer (0.025 M sodium bicarbonate, 0.025 M sodium carbonate, pH = 9.7) by incubating for 18 h at 4°C. Nonspecific binding was prevented by adding a blocking solution containing 1M phosphoric acid. The captured antigens were bound by a second specific monoclonal antibody (mAb). After washing, the amount of specifically bound mAb was then detected using anti-rat IgG conjugated to horse radish peroxidase as a tertiary reactant. The unbound conjugate was removed and following incubation with chromogenic substrate 3,3',5,5'-tetramethylbenzidine, the color change at 450 nm was recorded, as this indicated the amount of neurotrophic factors in each well.

Statistical Analysis

Comparisons were made using Mann-Whitney U test and one-way analysis of variance (ANOVA), and Spearman correlation coefficient by SPSS 10.0® (SPSS, Inc., Chicago, IL, USA). P values <0.05 were considered statistically significant.

Results

BDNF: AMYβ did not significantly change BDNF levels compared to controls (P = 0.075). Similarly, EPO, LPS + IFN γ and LPS + IFN γ + EPO did not change BDNF levels a significant amount (P > 0.05) (Table 1). A significant positive correlation was found between BDNF and NT3 (P < 0.0001; r = 0.667).

NT3: AMYβ did not significantly change NT3 levels compared to the controls (P = 0.067). EPO, LPS + IFN γ , and LPS + IFN γ + EPO did not change NT3 levels significantly (P > 0.05) (Table 1).

NT4: The lowest NT4 levels were found in the control group, while the highest levels were observed in the EPO 0.1 U/ml group. EPO at 0.1, 1.0 and 5.0 U/ml doses increased NT4 levels significantly compared to controls (P < 0.000 for all doses) and the levels of NT4 of the EPO 0.1 group were significantly higher than those of EPO 5.0 U/ml group (P < 0.05). AMYβ significantly increased NT4 levels compared to the control group (P < 0.05 or less for each group). Although levels of NT4 after LPS + IFN γ treatment were not significantly different than that of the controls, microglial activation by LPS + IFN γ + EPO (0.1, 1.0 and 5.0 U/ml) significantly increased NT4 levels compared to controls (P < 0.05, P < 0.05 and P < 0.0001, respectively). Of these groups, LPS + IFN γ +

Table 1. Levels of microglial brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT3) and neurotrophin-4 (NT4) levels (pg/ml) after exposure to amyloid beta (AMYβ), lipopolysaccharides (LPS) + interferon gamma (IFN γ), and erythropoietin (EPO). BDNF and NT3 results are reported as mean \pm SD while NT4 results are median.

Groups	BDNF	NT3	NT4
Control	3933 \pm 388	1000 \pm 98	501 ^c
LPS + IFN γ	3583 \pm 782	725 \pm 240	504
AMYβ	4433 \pm 759	1170 \pm 166	1164 ^d
EPO (0.1 U/ml)	3167 \pm 580	952 \pm 65	1859 ^{a,b}
EPO (1.0 U/ml)	2817 \pm 1327	1034 \pm 192	1656 ^a
EPO (5.0 U/ml)	3583 \pm 586	996 \pm 51	1439 ^a
LPS + IFN γ + EPO (0.1 U/ml)	2817 \pm 617	803 \pm 82	1400 ^d
LPS + IFN γ + EPO (1.0 U/ml)	3383 \pm 858	956 \pm 172	1223 ^d
LPS + IFN γ + EPO (5.0 U/ml)	3167 \pm 231	1394 \pm 537	1658 ^{a,e}

^a P < 0.0001, significantly higher than the control and LPS + IFN groups.

^b P < 0.05, significantly higher than the EPO 5.0 group.

^c P < 0.05, significantly lower than the other groups except LPS + IFN γ group.

^d P < 0.05, significantly higher than the control and LPS + IFN γ groups.

^e P < 0.01 significantly higher than the EPO 5.0 group.

EPO 5.0 U/ml had a greater effect than the other only EPO groups, and NT4 levels in the LPS + IFN γ + EPO 5.0 U/ml group were significantly higher than those of the only EPO 5.0 U/ml group ($P < 0.01$) (Table 1).

Discussion

Neurotrophic factor expressions were investigated in the pathogenesis of neurodegenerative diseases and were found useful for both neuroactive drug screening *in vitro* and possibly cell therapy for neurodegenerative diseases (17,18). Cortical BDNF levels were upregulated, showing an over 10-fold increase compared with age-matched controls in AD. BDNF upregulation was significantly correlated with the AMY β load, and thus the authors implicated this NT as playing an important role in the regulation of neuronal survival, axonal regeneration processes and neurological disorders, including epilepsy (19-24). In PD, glia may upregulate NTs like BDNF, NT3, NT4 and the tyrosine kinase receptors, TRKB and TRKC, in response to signals released mainly from activated microglial cells, failing nigral neurons and, to a lesser extent, from reactive astrocytes (25). This glial response may be the source of neurotrophic factors and can protect against reactive oxygen species including NO and glutamate (26). Alternatively, this glial response can also cause release of deleterious pro-oxidant reactive species and pro-inflammatory prostaglandins, LPS and IFN γ (25). These are also powerful inducers of NGF and BDNF production in mouse astrocyte cultures and thus play a neuroprotective role in the CNS (27). Likewise, in brain microglial cells, LPS modulate the expression of both NT3 and TRKC, which play pivotal roles in inflammatory and degenerative disorders (15). Some studies indicated that LPS induce microglial NGF expression and also analyzed the effects of the inflammatory agent on TRKA, the high-affinity receptor for NGF (28,29). In particular, it was reported that microglial NGF induces death of retinal ganglion cells (30). Therefore, it remains to be determined whether an induction in microglial NT expression during inflammation is beneficial or detrimental for neuronal or non-neuronal cells. Miwa et al. (31) demonstrated that microglia express mRNAs of BDNF and NT4 but not NT3 by LPS induction. Their results for NT3 and NT4 are inconsistent with those of Elkabes et al. (32), who showed that LPS induced microglial NT3 expression, while BDNF and NT4 expression were unaltered by LPS.

Proinflammatory cytokines were found to regulate expression of EPO and EPO receptor in human neurons, astrocytes, and microglia and further facilitate interactions among different cell types in the human CNS (18). Treatment with EPO inhibited the expression of iNOS mRNA and nitrite production resulting from proinflammatory stimulation by IFN γ and LPS in rat oligodendrocytes (8). Park et al. (9) revealed that EPO promoted neuronal cell differentiation through increased release of NGF from astrocytes, and this effect may be associated with signals of an extracellular signal-regulated kinase pathway. Neuronal differentiation was even further promoted when the neuronal stem cells were cultured with an astrocyte culture medium treated with a 10 U/ml dose of EPO. Compared to untreated astrocytes, EPO-treated astrocytes increased about two-fold in beta-NGF, but did not raise BDNF and NT3 levels. Similar to the study of Park et al., we also found in our study that BDNF and NT3 levels were unchanged and positively correlated with each other in microglia.

Wang et al. (33) showed that EPO treatment significantly increased brain levels of BDNF, improved functional recovery, and increased the density of cerebral microvessels at the site of stroke. Zhang et al. (34) found that EPO improved functional recovery after experimental allergic encephalomyelitis in mice, possibly via stimulating oligodendroglial BDNF expression. EPO also affects the ability of oligodendrocytes to promote myelin repair in the normal and damaged adult CNS.

It was reported by Viviani et al. (35) that exposure of primary hippocampal neurons to EPO significantly increased BDNF after 1 h. BDNF mRNA levels further increased up to 4 h, to return to control levels after 18 h of EPO treatment. Miwa et al. (15) found an increased BDNF and NT4 mRNA expression in primary microglia after LPS treatment for 12 h. Olivieri et al. (36) reported increased BDNF and NT4 levels in SHSY5Y cell lines after 12-h AMY β treatment. Mizuno et al. (37) found decreased BDNF mRNA levels after 24 h incubation of primary microglia with nicergoline. Elkabes et al. (32) reported an increase in NT3 and NGF with LPS continuing up to 24 h, while no change was noted in BDNF and NT4 levels. For the incubation period (24 h) selected in our study, the studies of Mizuno (37) and Elkabes (32) were taken as reference.

Both "EPO alone" groups and EPO + LPS + IFN γ group activated microglia and significantly increased NT4

levels compared to the control. However, the latter's significance was less than that of the former. We also found that increases in NT4 were more evident at the 0.1 U/ml and 1.0 U/ml doses of EPO, which is similar to the findings of Wen et al. (38). The enhanced NT4 secretion in microglial cells after exposure to AMY β may be an inflammatory reaction against this toxic agent. That is, microglia produce an immunological response to inflammation caused by AMY β , which is a molecule responsible for neuronal injury in neurodegeneration. EPO is more neuroprotective at lower doses and may also be a useful agent for exploring the regulatory mechanism of NT4 synthesis in murine microglia.

AMY β is a toxic and EPO is a protective agent; both of them significantly increased NT4 levels compared to the control. Ending with similar results can be speculated to arise by the following mechanisms: as a defense reaction against AMY β , microglial cells could have secreted neurotrophic factor; on the other hand, neurotrophic factor release could be the result of one of EPO's protective effects. Chong et al. (39) showed that EPO treatment protected the primary hippocampal neurons from apoptotic tissue injury in a rat model of AMY β toxicity. In their study, EPO in a concentration range of 0.001 ng/ml to 1000 ng/ml was not toxic to neurons. 1-50 ng/ml concentrations of EPO enhanced neuronal survival. However, EPO concentration <1 ng/ml or >50

ng/ml did have any positive effect on neuronal survival during AMY β administration. Chong et al. (39) showed that in a concentration-specific manner, EPO was able to protect DNA integrity and cellular membrane asymmetry during AMY β exposure. Additional studies in a number of cell models also illustrate a tight therapeutic concentration range for EPO (14,40-42). In the present study, AMY β significantly increased NT4 levels compared to controls. Although it also caused increases in BDNF and NT3, these were not significant. These results may indicate a reactive inflammatory response against damage induced by AMY β , and having significant changes only in NT4 could be the consequence of differing receptor sensitivities and receptor densities of different glial cell types.

In summary, our study results reveal that both AMY β and EPO stimulate murine microglial cells, maintaining support for neurons by increasing NT4 production, which is known to facilitate neuronal survival. Targeting microglia as mediators for neuroprotective drugs may be a promising strategy in the treatment of neurodegenerative processes.

Acknowledgements

The authors thank Dr. Paola Ricciardi-Castagnoli (Cellular Pharmacology Center, Milan, Italy) for providing the N9 murine microglial cell line.

References

- Mrak RE, Griffin WS. Glia and their cytokines in progression of neurodegeneration. *Neurobiol Aging* 2005; 26: 349-54.
- Walker DG, Lue LF. Investigations with cultured human microglia on pathogenic mechanisms of Alzheimer's disease and other neurodegenerative diseases. *J Neurosci Res* 2005; 81: 412-25.
- Dudal S, Krzywkowski P, Paquette J, Morissette C, Lacombe D, Tremblay P et al. Inflammation occurs early during the Abeta deposition process in TgCRND8 mice. *Neurobiol Aging* 2004; 25: 861-71.
- Rozemuller AJ, Van Gool WA, Eikelenboom P. The neuroinflammatory response in plaques and amyloid angiopathy in Alzheimer's disease: therapeutic implications. *Curr Drug Targets CNS Neurol Disord* 2005; 4: 223-33.
- Sheng JG, Mrak RE, Griffin WS. Glial-neuronal interactions in Alzheimer disease: progressive association of IL-1 alpha + microglia and S100beta + astrocytes with neurofibrillary tangle stages. *J Neuropathol Exp Neurol* 1997; 56: 285-90.
- Koshimura K, Murakami Y, Sohmiya M, Tanaka J, Kato Y. Effects of erythropoietin on neuronal activity. *J Neurochem* 1999; 72: 2565-72.
- Griffin WS. Inflammation and neurodegenerative diseases. *Am J Clin Nutr* 2006; 83: 470S-474S.
- Genc K, Genc S, Baskın H, Semin I. Erythropoietin decreases cytotoxicity and nitric oxide formation induced by inflammatory stimuli in rat oligodendrocytes. *Physiol Res* 2006; 55: 33-8.
- Park MH, Lee SM, Lee JW, Son DJ, Moon DC, Yoon DY et al. ERK-mediated production of neurotrophic factors by astrocytes promotes neuronal stem cell differentiation by erythropoietin. *Biochem Biophys Res Commun* 2006; 339: 1021-8.
- Genc S, Akhisaroglu M, Kuralay F, Genc K. Erythropoietin restores glutathione peroxidase activity in 1-methyl-4-phenyl-1,2,5,6-tetrahydropyridine-induced neurotoxicity in C-57BL mice and stimulates murine astroglial glutathione peroxidase production in vitro. *Neurosci Lett* 2002; 321: 73-6.
- Genc S, Kuralay F, Genc K, Akhisaroglu M, Fadiloglu S, Yorukoglu K et al. Erythropoietin exerts neuroprotection in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-treated C57BL mice via increasing nitric oxide production. *Neurosci Lett* 2001; 298: 139-41.

12. Siren AI, Fratelli M, Brines M, Geomans C, Casagrande S, Lewczuk P et al. Erythropoietin prevents neuronal apoptosis after cerebral ischemia and metabolic stress. *Proc Natl Acad Sci* 2001; 98: 4044-9.
13. Bernaudin M, Bellail A, Marti HH, Yvon A, Vivien D, Duchatelle I et al. Neurons and astrocytes express EPO mRNA: oxygen-sensing mechanisms that involve the redox-state of the brain. *Glia* 2000; 30: 271-8.
14. Bernaudin M, Marti HH, Roussel S, Divoux D, Nouvelot A, Mackenzie ET. A potential role for erythropoietin in focal permanent cerebral ischemia in mice. *J Cereb Blood Flow Metab* 1999; 19: 643-51.
15. Miwa J, Furukawa S, Nakajima K, Furukawa Y, Kohsaka S. Lipopolysaccharide enhances synthesis of brain-derived neurotrophic factor in cultured rat microglia. *J Neurosci Res* 1997; 50: 1023-9.
16. Tanaka J, Koshimura K, Sohmiya M, Murakami Y, Kato Y. Involvement of tetrahydrobiopterin in trophic effect of erythropoietin on PC12 cells. *Biochem Biophys Res Commun* 2001; 30: 358-62.
17. Caldwell MA, He X, Wilkie N, Pollack S, Marshall G, Wafford KA et al. Growth factors regulate the survival and fate of cells derived from human neurospheres. *Nat Biotechnol* 2001; 19: 475-9.
18. Darlington CL. Astrocytes as targets for neuroprotective drugs. *Curr Opin Investig Drugs* 2005; 6: 700.
19. Lahtinen S, Pitkanen A, Knuutila J, Toronen P, Castren E. Brain-derived neurotrophic factor signaling modifies hippocampal gene expression during epileptogenesis in transgenic mice. *Eur J Neurosci* 2004; 19: 3245-54.
20. Burbach GJ, Hellweg R, Haas CA, Del Turco D, Deicke U, Abramowski D et al. Induction of brain-derived neurotrophic factor in plaque-associated glial cells of aged APP23 transgenic mice. *J Neurosci* 2004; 24: 2421-30.
21. Duman RS. Role of neurotrophic factors in the etiology and treatment of mood disorders. *Neuromolecular Med* 2004; 5: 11-25.
22. Fraher J. Axons and glial interfaces: ultrastructural studies. *Anat* 2002; 200: 415-30.
23. Chan JR, Cosgaya JM, Wu YJ, Shooter EM. Neurotrophins are key mediators of the myelination program in the peripheral nervous system. *Proc Natl Acad Sci U S A* 2001; 98: 14661-8.
24. Dougherty KD, Dreyfus CF, Black IB. Brain-derived neurotrophic factor in astrocytes, oligodendrocytes, and microglia/macrophages after spinal cord injury. *Neurobiol Dis* 2000; 7: 574-85.
25. Teismann P, Tieu K, Cohen O, Choi DK, Wu DC, Marks D et al. Pathogenic role of glial cells in Parkinson's disease. *Mov Disord* 2003; 18: 121-9.
26. Zhang J, Geula C, Lu C, Koziel H, Hatcher LM, Roien FJ. Neurotrophins regulate proliferation and survival of two microglial cell lines in vitro. *Exp Neurol* 2003; 183: 469-81.
27. Toyomoto M, Ohta M, Okumura K, Yano H, Matsumoto K, Inoue S et al. Prostaglandins are powerful inducers of NGF and BDNF production in mouse astrocyte cultures. *FEBS Lett* 2004; 562: 211-5.
28. Mallat M, Houlgatte R, Brachet P, Prochiantz A. Lipopolysaccharide-stimulated rat brain macrophages release NGF in vitro. *Dev Biol* 1989; 133: 309-11.
29. Heese K, Hock C, Otten U. Inflammatory signals induce neurotrophin expression in human microglial cells. *J Neurochem* 1998; 70: 699-707.
30. Frade JM, Barde YA. Nerve growth factor: two receptors, multiple functions. *Bioessays* 1998; 20: 137-45.
31. Miwa T, Furukawa S, Nakajima K, Furukawa Y, Kohsaka S. Lipopolysaccharide enhances synthesis of brain-derived neurotrophic factor in cultured rat microglia. *J Neurosci Res* 1997; 50: 1023-9.
32. Elkabes S, Peng L, Black IB. Lipopolysaccharide differentially regulates microglial trk receptor and neurotrophin expression. *J Neurosci Res* 1998; 54: 117-22.
33. Wang L, Zhang Z, Wang Y, Zhang R, Chopp M. Treatment of stroke with erythropoietin enhances neurogenesis and angiogenesis and improves neurological function in rats. *Stroke* 2004; 35: 1732-7.
34. Zhang J, Li Y, Cui Y, Chen J, Lu M, Elias SB et al. Erythropoietin treatment improves neurological functional recovery in EAE mice. *Brain Res* 2005; 1034: 34-9.
35. Viviani B, Bartesaghi S, Corsini E, Villa P, Ghezzi P, Garau A et al. Erythropoietin protects primary hippocampal neurons increasing the expression of brain-derived neurotrophic factor. *J Neurochem* 2005; 93: 412-21.
36. Olivieri G, Otten U, Meier F, Baysang G, Dimitriades-Schmutz B, Müller-Spahn F et al. Beta-amyloid modulates tyrosine kinase B receptor expression in SHSY5Y neuroblastoma cells: influence of the antioxidant melatonin. *Neurosci* 2003; 120: 659-65.
37. Mizuno T, Kuno R, Nitta A, Nabeshima T, Zhang G, Kawanokuchi J et al. Protective effects of nicergoline against neuronal cell death induced by activated microglia and astrocytes. *Brain Res* 2005; 1066: 78-85.
38. Wen TC, Sadamoto Y, Tanaka J, Zhu PX, Nakata K, Ma YJ et al. Erythropoietin protects neurons against chemical hypoxia and cerebral ischemic injury by up-regulating Bcl-xl expression. *J Neurosci Res* 2002; 67: 795-803.
39. Chong ZZ, Li F, Maiese K. Erythropoietin requires NF-kappaB and its nuclear translocation to prevent early and late apoptotic neuronal injury during beta-amyloid toxicity. *Curr Neurovasc Res* 2005; 2: 387-99.
40. Chong ZZ, Kang JQ, Maiese K. Erythropoietin is a novel vascular protectant through activation of Akt1 and mitochondrial modulation of cysteine proteases. *Circulation* 2002; 106: 2973-9.
41. Chong ZZ, Kang JQ, Maiese K. Erythropoietin fosters both intrinsic and extrinsic neuronal protection through modulation of microglia, Akt1, Bad, and caspase-mediated pathways. *Br J Pharmacol* 2003; 138: 1107-18.
42. Kawakami M, Sekiguchi M, Sato K, Kozaki S, Takahashi M. Erythropoietin receptor-mediated inhibition of exocytotic glutamate release confers neuroprotection during chemical ischemia. *J Biol Chem* 2001; 276: 39469-75.