

1-1-2007

Cryptosporidiosis Among Children Attending Al-Nasser Pediatric Hospital, Gaza, Palestine

ADNAN IBRAHIM AL-HINDI

ABDELRAOUF A. ELMANAMA

KAMAL JAD ALLAH ELNABRIS

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

AL-HINDI, ADNAN IBRAHIM; ELMANAMA, ABDELRAOUF A.; and ELNABRIS, KAMAL JAD ALLAH (2007) "Cryptosporidiosis Among Children Attending Al-Nasser Pediatric Hospital, Gaza, Palestine," *Turkish Journal of Medical Sciences*: Vol. 37: No. 6, Article 7. Available at: <https://journals.tubitak.gov.tr/medical/vol37/iss6/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Cryptosporidiosis Among Children Attending Al-Nasser Pediatric Hospital, Gaza, Palestine

Adnan Ibrahim AL-HINDI¹
Abdelraouf A. ELMANAMA²
Kamal Jad Allah ELNABRIS¹

Aim: The present study was undertaken to determine the prevalence of cryptosporidiosis in patients with diarrhea who attended Al-Nasser Pediatric Hospital in Gaza.

Materials and Methods: From January to May 2005, single stool specimens from each of 416 children attending Al-Nasser Hospital were examined by Ziehl Neelsen (ZN) stain and ELISA for the presence of *Cryptosporidium* oocysts.

Results: *Cryptosporidium* oocysts were detected in 62 (14.9%) of the tested specimens by acid-fast staining technique and in 68 (16.3%) using ELISA kit. The number of infected females was found to be significantly higher than males ($P = 0.03$). The highest number of infected children was observed in March. *Entamoeba histolytica/dispar*, *Giardia lamblia*, *Ascaris lumbricoides* and *Hymenolepis nana* were the most common parasites detected (14.4%, 7.2%, 0.7% and 0.7%, respectively).

Conclusions: It is concluded that cryptosporidiosis still exists among children in Gaza. In addition, the diagnosis of protozoal parasites such as *Cryptosporidium* should draw more attention and should not be neglected due to its clinical importance.

Key Words: Antigen, children, *Cryptosporidium*, diarrhea, prevalence

¹ Department of Biology,
Faculty of Science,
The Islamic University of Gaza -
PALESTINE

² Department of Medical Technology,
Faculty of Science,
The Islamic University of Gaza -
PALESTINE

Al-Nasser Çocuk Hastanesine Başvuran Çocuklar Arasında Cryptosporidiosis Sıklığı

Amaç: Bu çalışma, Gazadaki Al-Nasser çocuk hastanesine ishal yakınması ile başvuran çocuklardaki Cryptosporidiosis sıklığını araştırmak amacıyla yapıldı.

Yöntem ve Gereç: 2005 yılı ocak ve mayıs ayları arasında başvuran 416 çocuktan alınan tek bir dışkı örneğinde direk boyama ve ELISA yöntemleri kullanılarak *Cryptosporidium* oocistleri araştırıldı.

Bulgular: Test edilen örneklerin 62'sinde (%14.9) boyama yöntemi ile ve 68'inde (%16.3) ELISA kiti kullanılarak *Cryptosporidium* oocistleri tespit edildi. Enfekte kız çocuklarının sayısı erkeklere göre daha yüksekti ($P = 0.03$). En fazla sayıda enfekte çocuk mart ayında saptandı. Diğer en sık saptanan parazitler sırasıyla *Entamoeba histolytica/dispar* (%14.4), *Giardia lamblia* (%7.2), *Ascaris lumbricoides* (%0.7) and *Hymenolepis nana* (%0.7) idi.

Sonuç: Gaza'daki çocuklarda Cryptosporidiosis halen mevcuttur ve klinik önemi nedeniyle bu durum göz ardı edilmeden tanı için daha dikkatli davranılmalıdır.

Anahtar Sözcükler: Çocuk, ishal, antijen, prevalans, cryptosporidium

Received: January 30, 2007
Accepted: November 15, 2007

Introduction

Cryptosporidium parvum, an intestinal protozoan parasite, is considered as one of the major diarrheagenic pathogens in humans throughout the world (1-4). There are several reports of hospital-based studies from different countries on the prevalence of *Cryptosporidium* in children with diarrhea. However, little information is available on asymptomatic carriage of *Cryptosporidium* in an apparently healthy population (5), and community-based studies carried out in depth are insufficient in the case of developing countries. Acute *Cryptosporidium* infection in children is usually associated with both acute and persistent diarrhea and various other gastrointestinal symptoms in immunocompetent persons and life-threatening illness in immunocompromised persons,

Correspondence

Adnan AL-HINDI
Department of Biology
Faculty of Science
The Islamic University of Gaza
P.O.Box 108
Gaza - PALESTINE

ahindi@iugaza.edu.ps

e.g. in acquired immunodeficiency syndrome (AIDS) patients (6). Asymptomatic cryptosporidiosis, which represents potential reservoirs of unrecognized, infected individuals who are capable of transmitting the infection to other individuals, was also documented (7-9). Asymptomatic individuals, livestock such as cattle and drinking water are known to be important vehicles in the epidemiology of the pathogen.

Cryptosporidium parvum appears to have become a threat to public health as it is ubiquitous and highly resistant to disinfectants and there is no effective therapy. The oocysts are shed in large quantities with the feces of infected individuals and are spread into the environment contaminating surface and ground waters. In a watery milieu, the parasite can survive for several months. It is completely resistant to the drinking water disinfection with chlorine. Cryptosporidiosis is endemic in developing and neighboring countries, such as in Fayoum (15%) (10) and among young children of the Nile River Delta in Egypt (17%) (11); in Israel (3.4-7.4%) (12); and among children in Jordan (37.3%) using direct immunofluorescence (13) as a result of poor sanitation and crowded living conditions (14). Gaza Strip dwellers have potentially high rates of parasitic infestation due to poor sanitation, groundwater contamination, infiltration of uncontrolled discharge of untreated sewage into the ground, contaminated water supply, and high population density (15).

In the Gaza Strip, a few unpublished reports mentioned the infection status of *Cryptosporidium*. The data resulting from this work would be of great importance in revealing the baseline prevalence of this parasite in Gaza and Palestine, and may help in undertaking any further interventions for its prevention and management.

Materials and Methods

The study population: The population included children attending Al- Nasser Pediatric Hospital in Gaza city from different ages and sex in the period from January to May 2005.

Description of study area: Al Nasser Hospital has four divisions in addition to the blood disease department (leukemia, hemophilia and thalassemia), intensive care unit, immature infants department, reception pharmacy, laboratory, and X-ray department. The total number of

inpatient beds is 151, with 33 beds for daily care (reception and emergency). The reception department receives 3,500–4,000 patients monthly. The general department receives 950-1,000 patients monthly, and inpatients normally stay for three days and are then discharged.

Sample size: A total of 416 stool samples (one sample for each child) were randomly collected from children attending Al Nasser Pediatric Hospital in the period of 15 January to 7 May 2005. The collection of stool samples was performed twice per week from 8 a.m. - 12 p.m.

Ethical considerations: Permission was obtained from the director of Al-Nasser Pediatric Hospital, the Director of the hospital's laboratory and the Helsinki ethical committee to facilitate the collection of stool samples from children attending the hospital. Families of all children were informed verbally about the purpose of the study.

Questionnaire: All children or their guardians were interviewed. The designed questionnaire was reviewed by a consultant in public health and was piloted before starting the study. Age, sex, residence, clinical information, and family information data were recorded.

Parasitological Methods

Direct smear microscopy: Each stool sample was subjected to a systematic scanning after being processed onto a clean slide, where the stool sample was diluted with a drop of normal saline, and a cover slip was placed on this preparation. In the present study, one stool sample was used due to difficulty in collection of three samples.

Formal ether concentration technique: The present study employed just one method of concentration, formal ether sedimentation technique, to detect the parasite or oocyst.

Modified Ziehl-Neelsen (ZN) technique: This method was used in the present study. After spreading the sediment onto a slide, it was stained with Ziehl-Neelsen (ZN) and examined under light microscope.

Enzyme Linked Immunosorbent Assay (ELISA): Stool samples were extracted and processed according to the manufacturer's recommendations (International Immuno-Diagnostics, 2003). Absorbance was read at "wavelength 450nm" using StatFax ELISA reader.

Statistical Analysis

Data generated from the sampling program along with those from the questionnaire were uploaded to Statistical Package for Social Sciences SPSS (version 11). Frequency and cross-tabulation were performed.

Results

During the study period from January to May 2005, stool specimens from 416 children were examined; 122 (29.3%) children had diarrhea. Sixty-two (14.9%) were confirmed as true positive for *Cryptosporidium* (Table 1). Two protozoan types (*Entamoeba histolytica/dispar* and *Giardia lamblia*) but only one type each of round worms and tapeworms (*Ascaris lumbricoides* and *Hymenolepis nana*, respectively) were observed during the present study, as shown in Table 2.

It was found that the rate of infection by both types of detected protozoa was higher in boys than in girls, while the rate of infection by both helminths was higher in girls than in boys, as indicated in Table 2, but there was no statistical difference.

The clinical features and other variables with relation to *Cryptosporidium* are summarized in Table 3. The age group with the highest rate of positivity for cryptosporidiosis was 1-4 years. Twenty-eight of cases (20.3%) positive for *Cryptosporidium* were girls, while

34 cases (12.2%) were boys. There was a significant difference in the distribution of cases between boys and girls. Prevalence of *Cryptosporidium* infection was similar in March and April.

Discussion

Cryptosporidium parvum cause a persistent diarrhea in developing countries (16). In the present study, *Cryptosporidium* oocysts were detected by modified ZN in 62 (14.9%) out of 416 examined stool specimens. Slightly higher prevalence (19%) was reported by Sallon et al. (17) among children attending the same institution as in the present study, Al-Nasser Pediatrics Hospital. Osman et al. (18) reported a similar prevalence (14.19%) among Egyptian children and attributed the high prevalence to lack of specific treatment and public health awareness of cryptosporidiosis. Our poor hygiene practices and lack of public health awareness among the community also contributed to such a prevalence. This has been only the second study since 1990 to investigate the prevalence of cryptosporidiosis in Gaza, and the prevalence is still high. We found apparent significant difference among females (20.3%) and males (12.2%) positive for *Cryptosporidium* ($P = 0.03$), which may be explained by the fact that girls are more exposed to the source of infection than boys.

Table 1. Diagnostic techniques for *Cryptosporidium* by staining and antigen detection.

Examination technique	<i>Cryptosporidium</i> positive n = 416	Rates of detection
Ziehl-Neelsen staining	62	14.9
Antigen detection	68	16.3

Table 2. Parasite distribution according to sex.

Parasite	Infected boys n = 276		Infected girls = 138	
	No.	%	No.	%
<i>Entamoeba histolytica/dispar</i>	40	(14.5)	12	(8.7)
<i>Giardia lamblia</i>	20	(7.2)	4	(2.9)
<i>Ascaris lumbricoides</i>	2	(0.7)	4	(2.9)
<i>Hymenolepis nana</i>	2	(0.7)	4	(2.9)

Table 3. Different variables associated with *Cryptosporidium* infection (n = 416).

Variables		Positive for <i>Cryptosporidium</i>		Negative for <i>Cryptosporidium</i>		P-value
		No.	%	No.	%	
Age	< 1 year	42	13.2	276	86.8	>0.05
	1-4 years	12	22.2	42	77.8	
	>5 years	8	18.2	36	81.8	
Sex	Boys	34	12.2	244	87.8	0.030
	Girls	28	20.3	110	79.7	
Patient	In-patient	32	12.4	226	87.6	0.067
	Out-patient	30	19.0	128	81.0	
Constipation	yes	8	30.8	18	69.2	0.019
	no	54	13.8	336	86.2	
Abdominal pain	yes	12	19.4	122	34.5	0.012
	no	50	80.6	232	65.5	
Loss of appetite	yes	8	7.1	104	92.9	0.007
	no	54	17.8	250	82.2	
Diarrhea	yes	6	4.4	130	95.6	0.001
	no	56	20.0	224	80.0	
Seasonal variation	Jan	12	9.7	112	90.3	0.017
	Feb	14	11.1	112	88.9	
	March	22	21.2	82	78.8	
	April	14	22.6	48	77.4	
Years of education (father)	<12 y	26	13.3	170	86.7	0.043
	12-18 y	2	4.8	40	95.2	
	>18 y	34	19.1	144	80.9	
Parents' occupation	Laborer	48	15.7	258	84.3	0.627
	Employed	10	11.6	76	88.4	
	Unemployed	4	16.7	20	83.3	

P < 0.05: significant.

P > 0.05: not significant.

When coproantigen was used for the same stained specimens by ZN in the present study, six specimens were found to be positive for *Cryptosporidium*. Regarding coproantigen and acid-fast staining, similar numbers and similar percentages were shown using both techniques. It seems there was no statistical or observational difference. Staining of stool specimens for *Cryptosporidium* and other protozoa is not performed in the local laboratories of private and/or governmental hospital clinics and

hospitals in Gaza Strip. Unfortunately, the dependence on direct microscopy using wet mount saline is the only method, which may affect the estimation of the prevalence of detected protozoa and/or epidemiological studies based on the Ministry of Health records. Most studies recommend that up to three separate, sequentially collected stool specimens should be examined in the laboratory to accurately diagnose enteric parasitic infection (19,20). In contrast, other studies have stated

that immunological-based detection methods are not significantly more sensitive than conventional microscopy (21,22).

In the present study, 22.2% of children positive for cryptosporidiosis were aged 1-4 years. Most studies have indicated that in developing countries, children may be most susceptible and those less than 2 years old may demonstrate the greatest prevalence (23,24). In another study, *Cryptosporidium* oocyst was found more among children aged below 2 years. MacPherson and Stephenson (25) reported that 30% of infected cases were aged 1-4 years, while Adegbola et al. (26) and Chacin-Bonilla et al. (27) reported that *Cryptosporidium* was frequent in children below 2 years. This high prevalence in children may be attributed to immature immunity and malnutrition (28) and the overall pattern suggests a decrease in prevalence after 12 months of age, which is most probably related to the development of immunity (29,30).

Despite age associations with *Cryptosporidium*, there was no clear trend in the prevalence of cryptosporidiosis with age in this study. In addition, it is known that children <5 years attend nurseries and kindergartens, where there is a little inspection for public health. The present finding indicated that 16.7% of infected children lived in houses with open sewage system versus 10.5% in houses with closed sewage system, with the higher rate due to more exposure to infection and contamination. The present study showed that diarrhea (4.4%) and other clinical features like constipation (30.8%), abdominal pain (9.0%), and loss of appetite (7.1%) were associated significantly with positive *Cryptosporidium* infection.

According to Al-Braiken et al. (31), the prevalence of mono-infection with *Cryptosporidium* among children with diarrhea (who presented to pediatric outpatient clinics) was 32% in Saudi Arabia. In Gaza, diarrheal disease is one of the greatest concerns among children. Another study carried out by Sallon et al. (17) showed that 19% of children with diarrhea were found to excrete *Cryptosporidium* oocyst. Inability of laborers and the unemployed to purchase medication may explain the high prevalence of *Cryptosporidium* among their children (15.7% and 16.7%, respectively) when compared to children of those employed (11.6%), which may be associated with the difficult socioeconomic conditions in which they live. Level of education was found to be associated with infection with *Cryptosporidium*. In our study, prevalence of cryptosporidiosis was observed in both March and April (21.2% and 22.6%, respectively), when there is transition from the cooler season to the start of the hot season. According to Sallon et al. [17], there was a striking difference in the prevalence of cryptosporidiosis between September 1989 (28%) and November 1989 (9.0%). Regarding seasonal activity, the hot season is more suitable for protozoal activity. It is concluded that cryptosporidiosis still constitutes a concern in Gaza and the diagnosis of *Cryptosporidium* should be improved in the governmental and private sector.

Acknowledgement

This work was supported by the Dean of Scientific Research/The Islamic University of Gaza. Thanks are extended to Dr. Abdel Rahman Eisa and the laboratory staff of Al-Nasser Pediatrics Hospital in Gaza, and to the families and their children.

References

1. Nime FA, Burek JD, Page DL, Holscher NA, Yardley JH. Acute enterocolitis in a human being infected with the protozoon *Cryptosporidium*. *Gastroenterology* 1976; 70: 592-8.
2. Chai JY, Shin SM, Yun CK, Yu JR, Lee SH. Experimental activation of cryptosporidiosis in mice by immunosuppression. *Korean J Parasitol* 1990; 28: 31-7.
3. Rhee JK, Seu YS, Park BK. Isolation and identification of *Cryptosporidium* from various animals in Korea. I. Prevalence of *Cryptosporidium* in various animals. *Korean J Parasitol* 1991; 29: 139-48.
4. Wee SH, Joo HD, Kang YB. Evaluation for detection of *Cryptosporidium* oocysts in diarrheal feces of calves. *Korean J Parasitol* 1996; 34: 121-6.
5. Miller K, Duran-Pinales C, Cruz-Lopez A, Morales-Lechuga L, Taren D, Enriquez FJ. *Cryptosporidium parvum* in children with diarrhoea in Mexico. *Am J Trop Med Hyg* 1994; 51: 322-5.
6. Ramratnam B, Flanigan TP. Cryptosporidiosis in persons with HIV infection. *Postgrad Med J* 1997; 73: 713-6.
7. Palit A, Sur D, MitraDhar K, Saha MR. Asymptomatic cryptosporidiosis in periurban slum setting in Kolkata, India – a pilot study. *Jpn J Infect Dis* 2005; 58: 110-1.

8. Pettoello-Mantovani M, Di Martino L, Dettori G, Vajro P, Scotti S, Ditullio MT et al. Asymptomatic carriage of intestinal *Cryptosporidium* in immunocompetent and immunodeficient children: a prospective study. *Pediatr Infect Dis J* 1995; 14: 1042-7.
9. Vuorio AF, Jokipii AM, Jokipii L. *Cryptosporidium* in asymptomatic children. *Rev Infect Dis* 1991; 13: 261-4.
10. EL-Mohamady H, Abdel-Messih IA, Youssef FG, Said M, Farag H, Shaheen HI et al. Enteric pathogens associated with diarrhea in children in Fayoum, Egypt. *Diagn Microbiol Infect Dis* 2006; 56(1): 1-5.
11. Abdel-Messih IA, Wierzba TF, Abu-Elyazeed R, Ibrahim AF, Ahmed SF, Kamal K et al. Diarrhoea associated with *Cryptosporidium parvum* among young children of the Nile River Delta in Egypt. *J Trop Pediatr* 2005; 51(3): 154-9.
12. Miron D, Colodner R, Kenes Y. Age-related seroprevalence of *Cryptosporidium* in northern Israel. *Isr Med Assoc J* 2000; 2(5): 343-5.
13. Mahgoub ES, Almahbashi A, Abdulatif B. Cryptosporidiosis in children in a north Jordanian paediatric hospital. *East Mediterr Health J* 2004; 10(4-5): 494-501.
14. Soave R, Ruiz J, Garcia-Saucedo V, Garrocho C, Kean BH. Cryptosporidiosis in a rural community in central Mexico [letter]. *J Infect Dis* 1989; 159: 1160-2.
15. Shubair ME, Yassin MM, Al-Hindi AI, al-Wahaidi AA, Jadallah SY, Abu Shaaban ND. Intestinal parasites in relation to haemoglobin level and nutritional status of school children in Gaza. *J Egypt Soc Parasitol* 2000; 30 (2): 365-75.
16. Current WL. *Cryptosporidium parvum*: household transmission. *Ann Intern Med* 1994; 120: 518-9.
17. Sallon S, El Showwa R, El Masri M, Khalil M, Blundell N, Hart CA. Cryptosporidiosis in children in Gaza. *Ann Trop Paediatr* 1991; 11: 277-81.
18. Osman GA, Makled KM, El-Shakankiry HM, Metwali DM, Abdel-Aziz SS, Saafan HH. Coccidian parasites as a cause of watery diarrhoea among protein energy malnourished and other immunocompromised Egyptian children. *J Egypt Soc Parasitol* 1999; 29(3): 653-68.
19. Melvin CM, Brooke MM. Laboratory procedures for the diagnosis of intestinal parasites, 3rd ed. US Department of Health, Education and Welfare publication no. (CDC) 82-8282. Centers for Disease Control, Atlanta, 1982.
20. Procedures for the recovery and identification of parasites from the intestinal tract. Proposed Guideline. NCCLS Document M28-p, 1993; 13: 6.
21. Kehl KS, Cicirello H, Havens PL. Comparison of four different methods for detection of *Cryptosporidium* species. *J Clin Microbiol*. 1995; 33(2): 416-8.
22. Rodriguez Hernandez J, Canut Blasco A, Martin Sanchez AM. [Epidemiology and diagnosis of *Cryptosporidium* spp. parasitosis in children: usefulness of the serologic study]. *Rev Clin Esp*. 1994; 194(5): 330-3.
23. Iqbal J, Hira PR, Al-Ali F, Philip R. Cryptosporidiosis in Kuwaiti children: seasonality and endemicity. *Clin Microbiol Infect*. 2001; 7(5): 261-6.
24. Reinthaler FF, Mascher F, Sixl W, Enayat U, Marth E. Cryptosporidiosis in children in Idukki District in southern India. *J Diarrhoeal Dis Res* 1989; 7(3-4): 89-91.
25. Macpherson DW, Stephenson BJ. Natural history of cryptosporidiosis infection in household contact. Abstract in 43rd Annual Meeting American Society of Tropical Medicine and Hygiene, 1994.
26. Adegbola RA, Demba E, De Veer G, Todd J. *Cryptosporidium* infection in Gambian children less than 5 years of age. *J Trop Med Hyg* 1994; 97: 103-7.
27. Chacin-Bonilla L, Bonilla MC, Soto-Torres L, Rios-Cadida Y, Sardina M, Enmanuel C et al. *Cryptosporidium parvum* in children with diarrhea in Zulia state, Venezuela. *Am J Trop Med Hyg* 1997; 56(4): 365-9.
28. Abdel-Maboud AI, Rossignol JF, El-Kady MS, Mos Tafa MS, Kabil SM. *Cryptosporidium* in Banha, study of some recent modalities in diagnosis and treatment. *J Egypt Soc Parasitol* 2000; 30(3): 717-25.
29. Current WL, Bick PH. Immunobiology of *Cryptosporidium* spp. *Pathol Immunopathol Res* 1989; 8: 141-60.
30. Lindo JF, Levy VA, Baum MK, Palmer CJ. Epidemiology of giardiasis and cryptosporidiosis in Jamaica. *Am J Trop Med Hyg* 1998; 59: 717-21.
31. Al-Braiken FA, Amin A, Beeching NJ, Hommel M, Hart CA. Detection of *Cryptosporidium* amongst diarrhoeic and asymptomatic children in Jeddah, Saudi Arabia. *Ann Trop Med Parasitol* 2003; 97(5): 505-10.