

1-1-2007

Effects of Cooking Techniques on Antioxidant Enzyme Activities of Some Fruits and Vegetables

İMGE B. ERGÜDER

ASLIHAN AVCI

ERDİNÇ DEVRİM

İLKER DURAK

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

ERGÜDER, İMGE B.; AVCI, ASLIHAN; DEVRİM, ERDİNÇ; and DURAK, İLKER (2007) "Effects of Cooking Techniques on Antioxidant Enzyme Activities of Some Fruits and Vegetables," *Turkish Journal of Medical Sciences*: Vol. 37: No. 3, Article 7. Available at: <https://journals.tubitak.gov.tr/medical/vol37/iss3/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Effects of Cooking Techniques on Antioxidant Enzyme Activities of Some Fruits and Vegetables

B. İmge ERGÜDER

Aslıhan AVCI

Erdoğan DEVRİM

İlker DURAK

Aim: The aim of this study was to investigate possible effects of cooking techniques on antioxidant enzyme activities in broccoli, tomato, red cabbage, parsley, carrot, green pepper, lemon, onion, and garlic, which are consumed frequently in our daily diet.

Materials and Methods: Superoxide dismutase (SOD), glutathione peroxidase (GSH-Px) and catalase (CAT) activities were measured in fresh and thermally treated (boiling, microwaving and baking) fruits and vegetables.

Results: When compared to raw vegetables, SOD, CAT and GSH-Px activities were found to diminish in most of the thermally treated vegetables except tomato, in which GSH-Px and CAT activities were found to increase. Our results show that antioxidant enzyme activities decrease in most of the fruits and vegetables after they are thermally treated, except in the tomato, in which increases are observed. It is possible that this property of the tomato is one of the reasons for its increased bioavailability after being thermally treated. We think that lycopene transfers from trans form to cis form, which is a more bioactive form, after being thermally treated.

Conclusions: We suggest that fruits and vegetables should be consumed raw if possible and tomatoes after thermal treatment.

Key Words: Antioxidant enzymes, influence of cooking

Department of Biochemistry,
Faculty of Medicine,
Ankara University,
06100 Ankara -TURKEY

Bazı Meyve ve Sebzelerin Antioksidan Enzim Aktiviteleri Üzerinde Pişirme Tekniklerinin Etkileri

Amaç: Bu çalışmanın amacı günlük diyetimizde sık olarak tükettiğimiz brokoli, domates, kırmızı lahana, maydanoz, havuç, yeşil biber, limon, soğan ve sarımsak gibi bazı meyve ve sebzelerde pişirme tekniklerinin antioksidan enzim aktiviteleri üzerine olası etkilerini incelemektir.

Yöntem ve Gereç: Çiğ ve pişirilmiş (kaynatma, mikrodalga ve fırın) meyve ve sebzelerde süperoksit dismutaz (SOD), glutatyon peroksidaz (GSH-Px) ve katalaz (KAT) enzim aktivitelerinin ölçümleri yapıldı.

Bulgular: Çiğ değerleriyle karşılaştırıldığında, SOD, KAT ve GSH-Px aktiviteleri, domates hariç, pişirilmiş diğer meyve ve sebzelerin tamamına yakınında azalmış olarak bulundu. Pişirilmiş domateste ise GSH-Px ve KAT aktiviteleri artmış olarak bulundu. Sonuçlarımız göstermiştir ki pişirme işlemleri bu yiyeceklerdeki antioksidan enzimlerde aktivite azalmasına yol açmaktadır, fakat domatesin pişirilmesi antioksidan enzimlerde aktivite artışına neden olmaktadır. Bunun nedeninin likopenin ısıyla muamele sonrasında trans formundan biyoyararlanımı daha yüksek olan cis formuna dönüşmesi olabileceğini düşünmekteyiz.

Sonuç: Sonuç olarak meyve ve sebzelerin mümkünse çiğ olarak tüketilmesinin, domatesin ise pişirme sonrası tüketilmesinin daha yararlı olacağını önerebiliriz.

Anahtar Sözcükler: Antioksidan enzimler, pişirmenin etkileri

Received: November 06, 2006

Accepted: March 22, 2007

Correspondence

B. İmge ERGÜDER

Department of Biochemistry,

Faculty of Medicine,

Ankara University,

06100 Sıhhiye, Ankara - TURKEY

imgeerguder@yahoo.com

Introduction

Epidemiological studies show that consumption of fruits and vegetables is associated with reduced risk of chronic diseases (1). Nutritional recommendations to reduce the risk of coronary heart disease have focused largely on the intake of nutrients that affect the risk factors, including plasma lipid and lipoprotein levels, blood pressure, and body weight, etc. Over the past two decades, considerable evidence has been gathered in support of the hypothesis that free-radical-mediated oxidative processes and specific end products play a key role in atherogenesis. This may involve effects on intimal

proliferation, fibrosis, calcification, endothelial function, vasoreactivity, plaque rupture, and thrombosis. Antioxidants constitute a diverse group of compounds with different properties. They operate by inhibiting oxidant formation, intercepting oxidants once they have formed, and repairing oxidant-induced injury. In terms of the coronary heart disease process, several points of antioxidant intervention have been proposed. Inhibition of low density lipoprotein (LDL) oxidation is the most well-known. Antioxidants create resistance to the oxidants. Although there is an effective antioxidant defense mechanism to protect the body against oxidant attacks, sometimes it cannot cope with oxidant load in the body, and additional dietary antioxidants are needed. Among them, vitamin C, vitamin E, flavonoids, and carotenes have received the greatest attention with regard to coronary heart disease prevention. A number of other dietary constituents are also proposed to act as antioxidants and to protect against coronary heart disease (2-11). Antioxidant properties of carotenoids are thought to be at least partly responsible for the protective effects of fruits and vegetables rich in carotenoids against colon cancer (12). Beta-carotene has been shown to enhance immune functions in humans (13).

Green leafy vegetables, fruits and vegetable oils are excellent sources of antioxidant components. Most vegetables undergo a cooking process rather than being eaten raw. Cooking practices may affect the antioxidant content and properties in vegetables (14); however, little is known in this regard.

Therefore, this study was carried out to determine the effects of domestic cooking practices such as boiling, microwaving and baking on antioxidant enzyme activities. We investigated activities of antioxidant enzymes, namely superoxide dismutase (SOD), glutathione peroxidase (GSH-Px) and catalase (CAT), in fresh and thermally treated fruits and vegetables, including lemon, tomato, red cabbage, parsley, carrot, green pepper, broccoli, onion and garlic.

Materials and Methods

All fruits and vegetables (lemon, broccoli, tomato, red cabbage, parsley, carrot, green pepper, onion, garlic) used in this study were purchased from a market and immediately cleaned, washed, sliced and homogenized in distilled water (2 g vegetable in 10 ml distilled water).

Fruits and vegetables were then allocated to four groups as raw and thermally treated (boiling, microwaving and baking). Of the three thermally treated groups, one was boiled for 20 minutes, one microwave-baked for one minute (only lemon for 30 sec.) at 360 watt and the third classical-oven baked for 10-20 minutes at 200°C. Cooking times were determined according to common cooking techniques. All samples were filtered using filter paper. GSH-Px activity was measured by following changes in NADPH absorbance at 340 nm (15). CAT activity was determined by measuring decrease in H₂O₂ absorbance at 240 nm (16). In the activity calculations, extinction coefficients of H₂O₂ and NADPH were used for CAT and GSH-Px, respectively. SOD activity was measured by the method based on nitro blue tetrazolium (NBT) reduction rate (17). One unit for SOD activity was expressed as the enzyme amount causing 50% inhibition in NBT reduction rate. Activity results were expressed as unit / g weight of related vegetable. In this study, all measurements were carried out three times and their mean values were used in the calculations.

Statistics

One sample t test was used in the statistical evaluation of enzyme activities between raw and thermally treated vegetables. P values lower than 0.05 were considered as significant.

Results

Mean enzyme activities are given in Figures 1-3 and percent changes in thermally treated fruits and vegetables relative to their raw forms are given in Tables 1-3. When compared to raw forms, SOD, CAT and GSH-Px activities were found to diminish significantly in most of the thermally treated fruits and vegetables. GSH-Px and CAT activities, however, were found to increase in the thermally treated tomato.

Discussion

Reactive oxygen species (ROS) have been implicated in tissue injury. The main ROS to be considered are superoxide anion (O₂⁻), which is predominantly generated by the mitochondria; hydrogen peroxide (H₂O₂) produced from O₂⁻ by the action of SOD, and peroxynitrite (ONOO⁻), generated by the reaction of O₂⁻ with nitric oxide (NO). These ROS produced are scavenged by SOD, GSH-Px and




Figure 1. Mean SOD activities in raw and thermally treated fruits and vegetables.

B: Broccoli. T: Tomato. RC: Red cabbage.
 P: Parsley. C: Carrot. GP: Green pepper.
 L: Lemon. O: Onion. G: Garlic.




Figure 2. Mean GSH-Px activities in raw and thermally treated fruits and vegetables.

B: Broccoli. T: Tomato. RC: Red cabbage.
 P: Parsley. C: Carrot. GP: Green pepper.
 L: Lemon. O: Onion. G: Garlic.




Figure 3. Mean CAT activities in raw and thermally treated fruits and vegetables.

B: Broccoli. T: Tomato. RC: Red cabbage.
 P: Parsley. C: Carrot. GP: Green pepper.
 L: Lemon. O: Onion. G: Garlic.

CAT. Under some circumstances, these endogenous antioxidant defenses are likely to be perturbed as a result of overproduction of oxygen radicals, inactivation of detoxification systems, consumption of antioxidants, and failure to adequately replenish antioxidants in tissue. It

has been demonstrated in numerous studies that ROS are directly involved in oxidative damage of cellular macromolecules such as lipids, proteins, and nucleic acids in tissues (18).

Table 1. Percent changes in SOD activities of thermally treated fruits and vegetables relative to their raw forms.

	Broccoli	Tomato	Red cabbage	Parsley	Carrot	G pepper	Lemon	Onion	Garlic
Boiling	-36.51 ^a	0.46	-39.01 ^a	-70.40 ^a	-51.47 ^a	-25.66 ^a	-4.78	-60.11 ^a	-85.57 ^a
Microwaving	-45.17 ^b	-13.92 ^b	-64.84 ^b	-78.16 ^b	-53.13 ^b	-26.99 ^b	3.35	-77.32 ^b	-96.22 ^b
Baking	-28.81 ^c	-8.32 ^c	-46.81 ^c	-75.43 ^c	-68.03 ^c	-26.74 ^c	-0.01	-26.38 ^c	-89.59 ^c

^a: P < 0.005; One sample t-test against raw vegetables.

^b: P < 0.005; One sample t-test against raw vegetables.

^c: P < 0.005; One sample t-test against raw vegetables.

Table 2. Percent changes in GSH-Px activities of thermally treated fruits and vegetables relative to their raw forms.

	Broccoli	Tomato	Red cabbage	Parsley	Carrot	G pepper	Lemon	Onion	Garlic
Boiling	0	400 ^a	-83.33 ^a	-33.33 ^a	-75 ^a	-100 ^a	0	-66.67 ^a	-85.71 ^a
Microwaving	-100 ^b	1300 ^b	-33.33 ^b	100 ^b	-100 ^b	100 ^b	0	-66.67 ^b	-100 ^b
Baking	133 ^c	200 ^c	-83.33 ^c	-100 ^c	0	100 ^c	0	-66.67 ^c	-85.71 ^c

^a: P < 0.005; One sample t-test against raw vegetables.

^b: P < 0.005; One sample t-test against raw vegetables.

^c: P < 0.005; One sample t-test against raw vegetables.

Table 3. Percent changes in CAT activities of thermally treated fruits and vegetables relative to their raw forms.

	Broccoli	Tomato	Red cabbage	Parsley	Carrot	G pepper	Lemon	Onion	Garlic
Boiling	-15.38 ^a	180 ^a	-97.30 ^a	-81.25 ^a	-44.44 ^a	-40 ^a	-50.00 ^a	-50.00 ^a	-95 ^a
Microwaving	-38.46 ^b	380 ^b	-86.49 ^b	-62.50 ^b	-66.67 ^b	-50 ^b	-66.67 ^b	-83.33 ^b	-85 ^b
Baking	-76.92 ^c	20 ^c	-91.89 ^c	-87.50 ^c	-88.89 ^c	-90 ^c	-50.00 ^c	-33.33 ^c	-70 ^c

^a: P < 0.005; One sample t-test against raw vegetables.

^b: P < 0.005; One sample t-test against raw vegetables.

^c: P < 0.005; One sample t-test against raw vegetables.

Domestic cooking practices, such as boiling, microwaving and frying, provoke a partial loss of the flavonols (19). There is considerable cooking loss of vitamins, and information on vitamin contents of cooked foods is essential for assessing the adequacy of vitamin intakes. The cooking losses were very high in case of ascorbic acid, thiamine, riboflavin, beta-carotene and folic acid (20). In one study, cooking of zucchini, beans and carrots with smaller amounts of water resulted in significantly higher content of phenolic phytochemicals in

the vegetables compared to cooking with larger water volumes (21). Lycopene is a bioactive carotenoid present in many fruits and vegetables. Tomatoes constitute the major dietary source of lycopene. Subjecting tomato juice to cooking temperatures in the presence of corn oil resulted in the formation of the cis form, which was considered to be more bioavailable (22). In addition to lycopene, tomato is a rich source of beta-carotene, folate, potassium, vitamin C, flavonoids, and vitamin E. The processing of tomatoes may significantly affect the

bioavailability of these nutrients. Homogenization, heat treatment, and the incorporation of oil in processed tomato products lead to increased lycopene bioavailability, while some of the same processes cause significant loss of other nutrients (23). Onion is a major source of flavonoids and is cooked in various ways throughout the world. The major flavonoids in onion are two quercetin glycosides, quercetin 4'-O-beta-glucoside (Q4'G) and quercetin 3,4'-O-beta-diglucosides (Q3,4'G), which are recognized as bioactive substances that are good for our health. Boiling, frying with oil and butter, and microwave cooking were examined in one study. The investigators reported that various cooking methods do not consider the degradation of quercetin conjugates when cooking onion. Microwave cooking without water better retains flavonoids and ascorbic acid. Frying does not affect flavonoid intake. The boiling of onion leads to about 30% loss of quercetin glycosides, which transfers to the boiling water (24).

As far as we know, this is the first study investigating the possible effects of cooking techniques on antioxidant enzyme activities in the fruits and vegetables consumed frequently in our daily diet. The results of the present study showed that all antioxidant enzyme activities decrease in the vegetables/fruits, except tomato, when they are thermally treated (boiling, microwaving and baking). However, we observed that GSH-Px and CAT

activities increase in the thermally treated tomato (boiling, microwaving and baking), and SOD activity increases in the boiled tomato whereas it decreases in the microwaved and baked tomato. Additionally, we determined that lemon has no GSH-Px activity in raw and thermally treated forms; this finding may be due to its acidic pH. On the other hand, GSH-Px activity was found to increase in microwaved parsley and in microwaved and baked green pepper.

Our results suggest that thermal treatments lead to loss in the activities of the antioxidant enzymes in most fruits and vegetables, which may cause decreases in their antioxidant potentials. However, the tomato appears to exhibit special properties in this regard, which indicate that thermal treatments cause significant increases in its antioxidant enzyme content activities. It is possible that this property of the tomato is one of the reasons for its increased bioavailability after cooking, which was also observed in a previous study (23). Hence, thermal treatments may lead to an increase in the antioxidant power of tomato, in contrast to observations in most other fruits and vegetables.

In conclusion, we suggest that it is better to consume most fruits and vegetables in their raw forms. Tomato is an exception in this regard, because the antioxidant property of tomato seems to increase after cooking.

References

1. Chu YF, Sun J, Wu X, Liu RH. Antioxidant and antiproliferative activities of common vegetables. *J Agric Food Chem* 2002; 50: 6910-6916.
2. Gokce N, Frei B. Basic research in antioxidant inhibition of steps in atherogenesis. *J Cardiovasc Risk* 1996; 3: 352-357.
3. Diaz MN, Frei B, Vita JA, Keaney JF Jr. Antioxidants and atherosclerotic heart disease. *N Engl J Med* 1997; 337: 408-416.
4. Hennekens CH, Gaziano JM, Manson JE, Buring JE. Antioxidant vitamin-cardiovascular disease hypothesis is still promising, but still unproven: the need for randomized trials. *Am J Clin Nutr* 1995; 62: 1377-1380.
5. Tavani A, Negri E, D'Avanzo B, La Vecchia C. Beta-carotene intake and risk of nonfatal acute myocardial infarction in women. *Eur J Epidemiol* 1997; 13: 631-637.
6. Kritchevsky SB, Tell GS, Shimakawa T, Dennis B, Li R, Kohlmeier L et al. Provitamin A carotenoid intake and carotid artery plaques: the Atherosclerosis Risk in Communities Study. *Am J Clin Nutr* 1998; 68: 726-733.
7. Jha P, Flather M, Lonn E, Farkouh M, Yusuf S. The antioxidant vitamins and cardiovascular disease: a critical review of epidemiologic and clinical trial data. *Ann Intern Med* 1995; 123: 860-872.
8. Stephens NG, Parsons A, Schofield PM, Kelly F, Cheeseman K, Mitchinson MJ. Randomised controlled trial of vitamin E in patients with coronary disease: Cambridge Heart Antioxidant Study. *Lancet* 1996; 347: 781-786.
9. Rapola JM, Virtamo J, Ripatti S, Huttunen JK, Albanes D, Taylor PR et al. Randomised trial of -tocopherol and -carotene supplements on incidence of major coronary events in men with previous myocardial infarction. *Lancet* 1997; 349: 1715-1720.
10. Tardif J-C, Cote G, Lesperance J, Bourassa M, Lambert J, Doucet S et al. Probuco and multivitamins in the prevention of restenosis after coronary angioplasty: multivitamins and Probuco Study Group. *N Engl J Med* 1997; 337: 365-372.

11. Yokoi H, Daida H, Kuwabara Y, Nishikawa H, Takatsu F, Tomihara H et al. Effectiveness of an antioxidant in preventing restenosis after percutaneous transluminal coronary angioplasty: the Probucol Angioplasty Restenosis Trial. *J Am Coll Cardiol* 1997; 30: 855-862.
12. Briviba K, Schnabele K, Rechkemmer G, Bub A. Supplementation of a diet low in carotenoids with tomato or carrot juice does not affect lipid peroxidation in plasma and feces of healthy men. *J Nutr* 2004; 134: 1801-1803.
13. Watzl B, Bub A, Briviba K, Rechkemmer G. Supplementation of a low-carotenoid diet with tomato or carrot juice modulates immune functions in healthy men. *Ann Nutr Metab* 2003; 47: 255-261.
14. Agostini LR, Moron Jimenez MJ, Ramon AN, Ayala Gomez A. [Determination of the antioxidant capacity of flavonoids in fruits and fresh and thermally treated vegetables.] *Arch Latinoam Nutr* 2004; 54: 89-92.
15. Paglia DE, Valentine WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med* 1967; 70: 158-169.
16. Aebi H. Catalase. In: Bergmayer HU, editor. *Methods of Enzymatic Analysis*. New York and London: Academic Press Inc; 1974. p.673-7.
17. Durak I, Canbolat O, Kavutcu M, Ozturk HS, Yurtarslani Z. Activities of total cytoplasmic and mitochondrial superoxide dismutase enzymes in sera and pleural fluids from patients with lung cancer. *J Clin Lab Anal* 1996; 10: 17-20.
18. Irmak MK, Fadilloğlu E, Güleç M, Erdoğan H, Yağmurca M, Akyol Ö. Effects of electromagnetic radiation from a cellular telephone on the oxidant and antioxidant levels in rabbits. *Cell Biochem Funct* 2002; 20: 279-283.
19. Tudela JA, Cantos E, Espin JC, Tomas-Barberan FA, Gil MI. Induction of antioxidant flavonol biosynthesis in fresh-cut potatoes. Effect of domestic cooking. *J Agric Food Chem* 2002; 50: 5925-5931.
20. Agte V, Tarwadi K, Mengale S, Hinge A, Chiplonkar S. Vitamin profile of cooked foods: how healthy is the practice of ready-to-eat foods? *Int J Food Sci Nutr* 2002; 53: 197-208.
21. Andlauer W, Stumpf C, Hubert M, Rings A, Furst P. Influence of cooking process on phenolic marker compounds of vegetables. *Int J Vitam Nutr Res* 2003; 73: 152-159.
22. Agarwal A, Shen H, Agarwal S, Rao AV. Lycopene content of tomato products: its stability, bioavailability and in vivo antioxidant properties. *J Med Food* 2001; 4: 9-15.
23. Willcox JK, Catignani GL, Lazarus S. Tomatoes and cardiovascular health. *Crit Rev Food Sci Nutr* 2003; 43: 1-18.
24. Ioku K, Aoyama Y, Tokuno A, Terao J, Nakatani N, Takei Y. Various cooking methods and the flavonoid content in onion. *J Nutr Sci Vitaminol* 2001; 47: 78-83.