

1-1-2006

Measurement of Cerebral and Cerebellar Volumes in Children with Fragile X Syndrome

NİMET KABAKUŞ

MUSTAFA AYDIN

AHMET KAVAKLI

HANEFİ YILDIRIM

HALUK AKIN

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

KABAKUŞ, NİMET; AYDIN, MUSTAFA; KAVAKLI, AHMET; YILDIRIM, HANEFİ; and AKIN, HALUK (2006) "Measurement of Cerebral and Cerebellar Volumes in Children with Fragile X Syndrome," *Turkish Journal of Medical Sciences*: Vol. 36: No. 3, Article 3. Available at: <https://journals.tubitak.gov.tr/medical/vol36/iss3/3>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Measurement of Cerebral and Cerebellar Volumes in Children with Fragile X Syndrome

Nimet KABAKUŞ¹
Mustafa AYDIN¹
Ahmet KAVAKLI²
Hanefi YILDIRIM³
Haluk AKIN⁴

Background and Aim: Fragile X syndrome (FXS) is a genetic disease associated with mental retardation, speech and memory deficits, developmental/learning disabilities, and, frequently, also with neuropsychological findings. It is known that all these abnormal brain functions could be associated with morphological anomalies of various brain regions. We investigated possible brain volume variations in children with FXS who had significant neurobehavioral symptoms and cognitive deficits.

Patients and Methods: This study was performed with 8 male FXS patients who were diagnosed cytogenetically. The Leiter International Performance Scales-Revised, the Stanford-Binet Intelligence Scale, the Vineland Social Maturity Scale, and DSM IV criteria were used for cognitive and behavioral assessment. The Cavalieri method, which is used to analyze the volumetric properties of smooth objects, was used for cranial volume measurements.

Results: The mean age of the patients was 8.8 ± 4.4 years (range: 5-18 years). The mean IQ level was 59.8 ± 10.3 (range: 40-76). The vermis/total cerebellum ratios of the patients decreased with age ($r = -0.851$, $P < 0.01$). There was a correlation between this condition and neurobehavioral symptoms that increased with age.

Conclusions: The study patients had decreased vermis/total cerebellum ratios, indicating the development of vermal atrophy with age. This vermal atrophy may have led to neurophysiological dysfunctions, cognitive deficits, and neurobehavioral phenotypes in the FXS patients.

Key Words: Brain volume, Cavalieri method, fragile X syndrome, magnetic resonance imaging, mental retardation

- ¹ Department of Pediatric Neurology, Faculty of Medicine, Firat University, Elazığ - TURKEY
² Department of Anatomy, Faculty of Medicine, Firat University, Elazığ - TURKEY
³ Department of Neuroradiology, Faculty of Medicine, Firat University, Elazığ - TURKEY
⁴ Department of Genetics, Faculty of Medicine, Firat University, Elazığ - TURKEY

Frajil X Sendromlu Çocuklarda Serebral ve Serebellar Volümlerin Ölçülmesi

Giriş ve Amaç: Frajil X sendromu (FXS) mental gerilik, dil ve hafıza kusurları, gelişim / öğrenim yetersizlikleri ile ilişkili ve ayrıca nöropsikolojik bulguların da sıkça görüldüğü kalıtsal bir hastalıktır. Tüm bu anormal beyin işlevlerinin beynin değişik bölgelerinin morfolojik anomalileri ile ilişkili olabileceği bilinmektedir. Bu çalışmada, belirgin nörodavranışsal semptomlar ve bilişsel kusurları olan FXS'li çocukların muhtemel beyin volüm değişikliklerinin incelenmesi amaçlandı.

Hastalar ve Yöntem: Çalışmaya sitogenetik olarak FXS tanısı konulan sekiz erkek hasta alındı. Bilişsel ve davranışsal değerlendirme için Yeniden Düzenlenmiş-Uluslararası Leiter Performans Ölçeği, Stanford Binet Zeka Ölçeği, Vineland Sosyal Gelişim Ölçeği ve DSM IV ölçütleri kullanıldı. Beyin volüm ölçümleri için, düz nesnelerin volümetrik özelliklerini değerlendirmede kullanılan, Cavalieri metodu kullanıldı.

Bulgular: Olguların yaş ortalaması $8,8 \pm 4,4$ (yaş aralığı 5-18) yıl idi. Ortalama IQ düzeyi $59,8 \pm 10,3$ (sınır aralığı 40-76) olarak belirlendi. Hastaların vermis / total serebellum oranları yaş ile azalma gösterdi ($r = -0,851$, $P < 0,01$). Yaş ile nörodavranışsal semptomların artması ve bu durum arasında korelasyon vardı.

Sonuç: Çalışma hastalarında, yaşla beraber vermal atrofi geliştiğine işaret eden, azalmış vermiş / total serebellum oranları vardı. Bu vermal atrofi, FXS hastalarda gözlenen nörofizyolojik fonksiyon bozuklukları, bilişsel kusurlar ve nörodavranışsal fenotiplere yol açabilir.

Anahtar Sözcükler: Beyin volümü, Cavalieri metodu, frajil X sendromu, manyetik rezonans görüntüleme, mental gerilik

Received: November 18, 2005
Accepted: June 15, 2006

Correspondence

Nimet KABAKUŞ
Department of Pediatric
Neurology, Faculty of Medicine,
Firat University,
23119 Elazığ - TURKEY
nimetkabakus@yahoo.com

Introduction

Recent advances in molecular genetics, human brain imaging, and behavioral studies have started to unravel the complex pathways leading to the cognitive, psychiatric, and physical features that are unique to fragile X syndrome (FXS). It has long been suggested that the decrease or absence of FMR1 protein production associated with FXS may lead to brain abnormalities in affected individuals (1-6). Postmortem microscopic examinations of brain tissue from patients with FXS have revealed that, although the number of neurons falls within the normal range, abnormalities in neuronal dendritic spines exist. These abnormalities are characterized by unusually long and thin, tortuous spines, with very few of the stubby, mushroom-shaped spines found in unaffected controls, and a higher density of spines along the dendrites, suggesting a possible failure of synapse elimination (3,7).

Neuroimaging studies have shown selective changes in brain size in FXS, which include reductions in the posterior cerebellar vermis, age-dependent increases in hippocampal volume, and an enlarged caudate nucleus and thalamus. Structural magnetic resonance imaging (MRI) studies of FXS have further explored the effects of the FMR1 full mutation on neuroanatomy. The first brain abnormality reported in a MRI study was hypoplasia of the cerebellar vermis, the connecting tissue between the right and left hemispheres of the cerebellum (8).

In the present study, we studied the relationship between volumetric changes in the gross structures of the brains of subjects with FXS and neurobehavioral signs, symptoms, and cognitive disorders, which might be observed in FXS patients.

Patients and Methods

This study was performed with 8 FXS patients who were diagnosed cytogenetically by lymphocyte culture in the presence of a folic acid antagonist. For this study, institutional ethical approval and permission from the parents were obtained. MRIs of the patients were evaluated, along with their histories, physical and neurological examinations, and their general laboratory findings.

The Leiter International Performance Scales-Revised, the Stanford-Binet Intelligence Scale, the Vineland Social

Maturity Scale, and DSM IV criteria were used by a psychologist for cognitive and behavioral assessment (9-11).

MRI was performed on a 1.5-T open scanner (High Speed Model, General Electric Signa) with 3-mm slice thickness. During cranial analysis, the following sequences were imaged: Spin Echo (SE) T1 (Fov: 240; TR: 500; TE: 20; FA: 90; thickness: 8 mm; interval: 10 mm; matrix: 256 x 256; NSA: 3; multislice: 12; projection: 160); Proton Dansite (PD) (Fov: 240; TR: 4000; TE: 26; FA: 90; Thickness: 8 mm; interval: 10 mm; NSA: 2; matrix: 256 x 256; projection: 192); Fast Spin Echo (FSE) T2 (Fov: 240; TR: 4000; TE: 117; FA: 90; Thickness: 8; interval: 10; multislice: 12; NSA: 4; projection: 160). To evaluate the hippocampus, a 3D coronal spoiled gradient echo (SGE) sequence was added (Fov: 200; TR: 70; TE: 18; FA: 50; Thickness: 3 mm; Interval: 3 mm; projection: 192; NSA: 3; matrix: 256 x 256; multislice: 32). After obtaining the sequences, the brain structures of the subjects were evaluated.

The Cavalieri method, which is used to analyze the volumetric properties of smooth objects, was used for cranial volume measurements (12). According to this method, first the size of the area (a) for each MRI plane section was calculated (cm^2). By multiplying the area (a) by the thickness of each MRI section (t), we obtained section volume ($V = ta$). This equation was extended according to general mathematical rules and the resulting formula [$V=t(a_1+a_2+.....+a_n) cm^3$] was obtained. The resulting equation was utilized in calculating the volumes. Volumetric measurements of the subjects were carried out by an experienced neuroradiologist and an anatomist. Volumetric measurements were taken of the right and left cerebral hemispheres, total cerebrum, cerebellum, and cerebellar vermis. Total brain volume was calculated by adding the volumes of the cerebrum and cerebellum. Total brain volume did not include the volume of the ventricles or the corpus callosum. We did not have volumetric measurements from normal children to compare with the values obtained from FXS subjects. Consequently, we used values from the literature of age- and sex-matched normal children (13-15).

The data were evaluated in the SPSS 11.0 program. The obtained data are presented with standard deviation. In the evaluation of the relationship between some of the parameters, Spearman's correlation analysis was used.

Results

The average age of the male patients was 8.8 ± 4.4 years (range: 5-18). The most common findings in their physical examinations were large ears (8/8, 100%), elongated face (6/8, 75%), protruding mandible (5/8, 62.5%), and macroorchidism (5/8, 62.5%). In addition to these, in each patient, hypospadias, strabismus, myopia, and macrocephaly were noted (12.5%) as separate cases. All patients had mental retardation (100%). Their average IQ level was 59.8 ± 10.3 (range: 40-76). Moreover, 3 patients had attention deficit hyperactivity disorder (ADHD) (37.5%), one had autism (12.5%), one had stereotyped behavior and speech apraxia, and one had social anxiety (Table).

While MRIs of 6 patients were found to be qualitatively normal, in 2 of the subjects (2/8, 25%), abnormalities were noted in cranial MRIs. These abnormalities were severe cerebellar/vermal hypoplasia (no: 2) and cerebral hemispherical asymmetry (no: 5, right: 699.2 cm^3 ; left: 533.6 cm^3). The vermis/total cerebellum ratios of the patients decreased with age ($r = -0.851$, $P < 0.01$) (Figure), while neurobehavioral symptoms appeared to increase (Table).

Discussion

The neuroanatomical basis for the cognitive deficits and neurobehavioral symptoms observed in FXS patients has not been satisfactorily established. The possibility that the brain dysfunctions noted in these patients are primarily the result of abnormalities in structural development has not been ruled out as yet (16-20).

Abnormal dendritic spine morphology in the cerebral cortex has been observed in neuropathological studies of FXS (3,21-23). However, others (24) found no differences in total cerebral volume in either males or females with FXS. Reiss and colleagues (25,26) found that the posterior cerebellar vermis region was reduced in both males and females with FRX compared to sex-matched controls. Another study by the same researchers, but with a larger sample (27), confirmed the absolute and relative decrease in posterior vermis (lobules VI-X) size in males with FXS ($n = 32$) when compared with both normal ($n = 38$) and developmentally delayed ($n = 28$) controls.

Associations between hypoplasia of the cerebellar vermis and behavioral and cognitive dysfunction in FXS have been reported in 2 studies (28,29). In an investigation of 30 school-age girls with FXS, Mazzocco and colleagues (28) reported a significant negative correlation between the area of lobules VI and VII, and parental ratings of frequency and severity of autistic behaviors, particularly in social communication and stereotypic behaviors. In contrast, measures of anxiety did not correlate with IQ or with the size of the posterior cerebellar vermis, suggesting that anxiety, which is included in the behavioral phenotype of girls with FXS, may have a different neuropathological mechanism that is unrelated to morphology of the posterior cerebellar vermis (1,29). In our study, no correlation between IQ and brain volume was established. This finding suggests that cellular disorders and the abnormalities in transmitter levels previously demonstrated in FXS subjects might have additional negative influence on IQ level (30-33).




Figure. Decreases in the ratio of vermis/total cerebellum with age ($r = -0.851$; $P < 0.01$).

Table. Cerebral and cerebellar volumes (cm³) of the patients with Fragile X syndrome.

Patient No	Age (year)	Right cerebral hemisphere	Left cerebral hemisphere	Cerebrum	Cerebellar vermis	Total cerebellum	Vermis / total cerebellum	Total brain volume	IQ	Neuro-behavioral sings and symptoms
1	5	623.7	628.8	1252.4	67.2	129.7	0.52	1382.1	40	MR
2*	5	614.1	624.2	1238.3	45.4	92.1	0.49	1330.3	60	MR / ADHD
3	6	627.9	627.9	1255.8	69.2	135.6	0.51	1391.4	65	MR
4	7	679.1	672.4	1351.6	73.6	144.9	0.50	1496.4	66	MR / ADHD
5*	8	699.2	533.6	1232.7	68.5	137.7	0.49	1370.5	56	MR / autism
6	10	649.8	643.6	1293.4	74.9	148.0	0.50	1442.4	76	MR / ADHD
7	12	666.9	662.9	1329.8	70.3	146.4	0.48	1476.2	60	MR / Stereotyped behavior and speech apraxia
8	18	671.1	671.1	1342.3	68.9	149.2	0.46	1491.4	56	MR / social anxiety
Mean ± SD	¹ 8.8 ± 4.4	654.0 ± 30.1	633.1 ± 44.8	1287.0 ± 48.7	67.2 ± 9.2	135.6 ± 18.9	¹ 0.49 ± 0.2	1422.6 ± 62.4	59.8 ± 10.3	
Reference 11	9.8	535.5	507.9		129.6					
Reference 12	10.8							1316.3		
Reference 13	4.4				71.4					

*: patients with cerebral malformations ¹; r=-0.851, P < 0.01 (Spearman's correlation analysis was used)

Reference 13: Hong et al., 2002; Reference 14: Moore et al., 2000; Reference 15: Pierce K et al., 2001.

ADHD: Attention-deficit / hyperactivity disorder

IQ: Intelligence quotient

MR: Mental retardation

SD: Standart deviation

It has been shown that the cerebellar vermis is connected to the frontal lobe and regulates higher cognitive functions (29). The most important finding obtained from our FXS series is that vermis/total cerebellum ratio decreased in these patients as the subjects aged, whereas neurobehavioral symptoms increased. Similar findings have been reported by others who have studied FXS subjects (24,29,32-34). Two brain regions, the cerebellar vermis, which is important for a variety of cognitive tasks and regulation of motor behavior, and the superior temporal gyrus, which is involved in processing complex auditory stimuli, are reported to be reduced in size in FXS patients relative to controls (25-27). The vermis, which is anatomically connected to limbic structures, including the hippocampus and amygdala, has been implicated in the execution and regulation of motor behavior (35), visual saccades (36), auditory processing (37), and some aspects of language (38,39). Abnormalities of the cerebellar vermis may, therefore, be linked to some of the behavioral anomalies associated with FXS, including hyperactivity and repetitive movements, tactile defensiveness, attention deficits, and language dysfunction (1).

Cranial MRI studies performed in our patients indicated that only 2 subjects had qualitative abnormalities, while qualitatively normal MRI findings were obtained in 6. Overall, these findings might be interpreted to mean that in FXS, even though structural brain abnormalities might exist, these changes are not unique to the syndrome.

The mean volumes of the right and left cerebral hemispheres in our FXS series were 654 ± 30.1 and $633.1 \pm 44.8 \text{ cm}^3$, respectively. The mean volumes from age- and sex-matched control subjects obtained from the literature were 535.5 and 507.9 cm^3 , respectively (13).

These findings demonstrated that our subjects had 120 to 125 cm^3 more mean mass on both the right and left hemispheres. Total brain volumes of our subjects were also higher with a mean volume of 1422.6 ± 62.4 vs. 1316.3 cm^3 as published in the literature for the normal 10.8-year-old age group (14). Cerebellar vermis volumes in our series (mean: $67.2 \pm 9.2 \text{ cm}^3$) were lower than those reported for the normal 4.4-year-old age group (mean: 71.4 cm^3). However, total cerebellum volumes (mean: $135.6 \pm 18.9 \text{ cm}^3$) were similar to those reported (mean: 129.6 cm^3) for the age- and sex-matched control subjects (13,15). These findings show that the FXS series studied in the current report had higher cerebral volume, yet lower cerebellar vermis volume. We are not able to provide a satisfactory explanation for this observation. Contrary to the reported findings (24-26), this discrepancy with the normal values in the literature (13-15) might have been caused by the increased total cerebral volume and decreased cerebellar vermis volume associated with aging and/or by racial or geographical differences existing between our study subjects and those reported in the literature. The lack of local control subjects in our study prevents us from further delineating the origin of this discrepancy.

In conclusion, the observed vermal volumetric changes in FXS patients might be responsible for some of the neurobehavioral signs, symptoms, and cognitive disorders associated with FXS. However, in order for all of these possible relationships to be brought to light, studies with larger patient populations, more sensitive combined research methods (positron emission tomography, diffusion-weighted imaging, proton magnetic resonance spectroscopy, and exciter/inhibitor neurotransmitters and performance tests), and multicenter design are needed.

References

1. Hessler D, Rivera SM, Reiss AL. The neuroanatomy and neuroendocrinology of fragile X syndrome. *Ment Retard Dev Disabil Res Rev* 2004; 10: 17-24.
2. Devys D, Lutz Y, Rouyer N, Belloccq JP, Mandel JL. The FMR-1 protein is cytoplasmic, most abundant in neurons and appears normal in carriers of a fragile X premutation. *Nat Genet* 1993; 4: 335-340.
3. Hinton VJ, Brown WT, Wisniewski K, Rudelli RD. Analysis of neocortex in three males with the fragile X syndrome. *Am J Med Genet* 1991; 41: 289-294.
4. Rudelli RD, Brown WT, Wisniewski K, Jenkins EC, Laure-Kamionowska M, Connell F et al. Adult fragile X syndrome. Clinico-neuropathologic findings. *Acta Neuropathol (Berl)* 1985; 67: 289-295.
5. Tamanini F, Willemsen R, van Unen L, Bontekoe C, Galjaard H, Oostra BA et al. Differential expression of FMR1, FXR1 and FXR2 proteins in human brain and testis. *Hum Mol Genet* 1997; 6: 1315-1322.

6. Wisniewski KE, Segan SM, Miezieski CM, Sersen EA, Rudelli RD. The Fra (X) syndrome: neurological, electrophysiological, and neuropathological abnormalities. *Am J Med Genet* 1991; 38: 476-480.
7. Irwin SA, Galvez R, Greenough WT. Dendritic spine structural anomalies in fragile-X mental retardation syndrome. *Cereb Cortex* 2000; 10: 1038-1044.
8. Reiss AL, Patel S, Kumar AJ, Freund L. Preliminary communication: neuroanatomical variations of the posterior fossa in men with the fragile X (Martin-Bell) syndrome. *Am J Med Genet* 1988; 31: 407-414.
9. Roid GH, Miller LJ. *Leiter International Performance Scale-Revised*. Wood Dale: Stoelting Company; 1997.
10. *Diagnostic and Statistical Manual of Mental disorders (DSM IV)*. Task Force on DSM-IV. American Psychiatric Association; 2000.
11. Thorndike RL, Hagen EP, Sattler JM. *The Stanford-Binet Intelligence Scale*. 4th ed. Chicago: The Riverside Publishing Company; 1986.
12. Milhorat TH, Chou MW, Trinidad EM, Kula RW, Mandell M, Wolpert C et al. Chiari I malformation redefined: clinical and radiographic findings for 364 symptomatic patients. *Neurosurgery* 1999; 44: 1005-1017.
13. Hong KE, Ock SM, Kang MH, Kim CE, Bae JN, Lim MK et al. The segmented regional volumes of the cerebrum and cerebellum in boys with Tourette syndrome. *J Korean Med Sci* 2002; 17: 530-536.
14. Moore BD, Slopis JM, Jackson EF, De Winter AE, Leeds NE. Brain volume in children with neurofibromatosis type 1. *Neurology* 2000; 54: 914-920.
15. Pierce K, Courchesne E. Evidence for a cerebellar role in reduced exploration and stereotyped behavior in autism. *Biol Psychiatry* 2001; 49: 655-664.
16. Munir F, Cornish KM, Wilding J. A neuropsychological profile of attention deficits in young males with fragile X syndrome. *Neuropsychologia* 2000; 38: 1261-1270.
17. Ferrando LMT, Banus GP, Lopez PG. Aspects of cognition and language in children with fragile X syndrome. *Rev Neurol* 2003; 36: 137-142.
18. Guerreiro MM, Camargo EE, Kato M, Marques-de-Faria AP, Ciasca SM, Guerreiro CA et al. Fragile X syndrome. Clinical, electroencephalographic and neuroimaging characteristics. *Arq Neuropsiquiatr* 1998; 56: 18-23.
19. Schapiro MB, Murphy DG, Hagerman RJ, Azari NP, Alexander GE, Miezieski CM et al. Adult fragile X syndrome: neuropsychology, brain anatomy, and metabolism. *Am J Med Genet* 1995; 18: 480-493.
20. Eliez S, Reiss AL. Genetics of childhood disorders: XI. Fragile X syndrome. *J Am Acad Child Adolesc Psychiatry* 2000; 39: 264-266.
21. Irwin SA, Patel B, Idupulapati M, Harris JB, Crisostomo RA, Larsen BP et al. Abnormal dendritic spine characteristics in the temporal and visual cortices of patients with fragile-X syndrome: a quantitative examination. *Am J Med Genet* 2001; 98: 161-167.
22. Kaufmann WE, Moser HW. Dendritic anomalies in disorders associated with mental retardation. *Cereb Cortex* 2000; 10: 981-991.
23. Comery TA, Harris JB, Willems PJ, Oostra BA, Irwin SA, Weiler IJ et al. Abnormal dendritic spines in fragile X knockout mice: maturation and pruning deficits. *Proc Natl Acad Sci USA* 1997; 94: 5401-5404.
24. Eliez S, Blasey C, Freund LS, Hastie T, Reiss AL. Brain anatomy, gender, and IQ in children and adolescents with fragile X syndrome. *Brain* 2001; 124: 1610-1618.
25. Reiss AL, Aylward E, Freund LS, Joshi PK, Bryan RN. Neuroanatomy of fragile X syndrome: the posterior fossa. *Ann Neurol* 1991; 29: 26-32.
26. Reiss AL, Freund L, Tseng JE, Joshi PK. Neuroanatomy in fragile X females: the posterior fossa. *Am J Hum Genet* 1991; 49: 279-288.
27. Mostofsky SH, Mazzocco MM, Aakalu G, Warsofsky IS, Denckla MB, Reiss AL. Decreased cerebellar posterior vermis size in fragile X syndrome: correlation with neurocognitive performance. *Neurology* 1998; 50: 121-130.
28. Mazzocco MM, Kates WR, Baumgardner TL, Freund LS, Reiss AL. Autistic behaviors among girls with fragile X syndrome. *J Autism Dev Disord* 1997; 27: 415-435.
29. Kates WR, Folley BS, Lanham DC, Capone GT, Kaufmann WE. Cerebral growth in fragile X syndrome: Review and comparison with Down syndrome. *Microscop Res Tech* 2002; 57: 159-167.
30. Todd PK, Mack KJ, Malter JS. The fragile X mental retardation protein is required for type-I metabotropic glutamate receptor-dependent translation of PSD-95. *Proc Natl Acad Sci USA* 2003; 100: 14374-14378.
31. Denman RB. Methylation of the arginine-glycine-rich region in the fragile X mental retardation protein FMRP differentially affects RNA binding. *Cell Mol Biol Lett* 2002; 7: 877-883.
32. Gruss M, Braun K. Alterations of amino acids and monoamine metabolism in male Fmr1 knockout mice: a putative animal model of the human fragile X mental retardation syndrome. *Neural Plast* 2001; 8: 285-298.
33. Gruss M, Braun K. Age- and region-specific imbalances of basal amino acids and monoamine metabolism in limbic regions of female Fmr 1 knock-out mice. *Neurochem Int* 2004; 45: 81-88.
34. Johnston C, Eliez S, Dyer-Friedman J, Hessel D, Glaser B, Blasey C et al. Neurobehavioral phenotype in carriers of the fragile X premutation. *Am J Med Genet* 2001; 103: 314-319.
35. Rosenthal G, Gilman S, Koeppe RA, Klun KJ, Markel DS, Junck L et al. Motor dysfunction in olivopontocerebellar atrophy is related to cerebral metabolic rate studied with positron emission tomography. *Ann Neurol* 1988; 24: 414-419.

36. Hayakawa Y, Nakajima T, Takagi M, Fukuhara N, Abe H. Human cerebellar activation in relation to saccadic eye movements: A functional magnetic resonance imaging study. *Ophthalmologica* 2002; 216: 399-405.
37. Huang CM, Burkard R. Frequency sensitivities of auditory neurons in the cerebellum of the cat. *Brain Res* 1986; 371: 101-108.
38. Moretti R, Bava A, Torre P, Antonello RM, Cazzato G. Reading errors in patients with cerebellar vermis lesions. *J Neurol* 2002; 249: 461-468.
39. Schmitt JE, Eliez S, Warsofsky IS, Bellugi U, Reiss AL. Enlarged cerebellar vermis in Williams syndrome. *J Psychiatr Res* 2001; 35: 225-229.