

1-1-2022

The conservation method effects of the Barley-Vetch mixture planted together on the performance of pregnant ewes

HASAN HÜSEYİN ŞENYÜZ

ENGİN ÜNAY


ABDULKADİR ERİŞEK

ARZU EROL TUNÇ

ÇAĞATAY YILDIRIM

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/veterinary>

 Part of the [Animal Sciences Commons](#), and the [Veterinary Medicine Commons](#)

Recommended Citation

ŞENYÜZ, HASAN HÜSEYİN; ÜNAY, ENGİN; ERİŞEK, ABDULKADİR; TUNÇ, ARZU EROL; YILDIRIM, ÇAĞATAY; ERDAL, NURGÜL; COŞKUN, MUHAMMED İKBAL; and MARAŞ, HALİL (2022) "The conservation method effects of the Barley-Vetch mixture planted together on the performance of pregnant ewes," *Turkish Journal of Veterinary & Animal Sciences*: Vol. 46: No. 2, Article 6. <https://doi.org/10.55730/1300-0128.4169>

Available at: <https://journals.tubitak.gov.tr/veterinary/vol46/iss2/6>









This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Veterinary & Animal Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

The conservation method effects of the Barley-Vetch mixture planted together on the performance of pregnant ewes

Authors

HASAN HÜSEYİN ŞENYÜZ, ENGİN ÜNAY, ABDULKADİR ERİŞEK, ARZU EROL TUNÇ, ÇAĞATAY YILDIRIM, NURGÜL ERDAL, MUHAMMED İKBAL COŞKUN, and HALİL MARAŞ

The conservation method effects of the Barley-Vetch mixture planted together on the performance of pregnant ewes

Hasan Hüseyin ŞENYÜZ* , Engin ÜNAY , Abdulkadir ERİŞEK , Arzu EROL TUNÇ ,
Çağatay YILDIRIM , Nurgül ERDAL , Muhammed İkbâl COŞKUN , Halil MARAŞ 
International Center for Livestock Research and Training, Mamak, Ankara, Turkey

Received: 17.08.2021 • Accepted/Published Online: 04.02.2022 • Final Version: 25.04.2022

Abstract: The aim of this study was to examine the effects of barley-vetch hay and barley-vetch silage on liveweight change in pregnant sheep and birthweight in lambs and to determine the optimal form of barley-vetch roughage usage in sheep feeding. Forty-eight (Kıvırcık x Akkaraman (G₁)) sheep were blocked according to age and randomly distributed to three groups: barley-vetch hay (H, n = 16), barley-vetch silage (S, n = 16), and the control group (C, n = 16). The H group was fed barley-vetch hay *ad libitum* and limited concentrate, the S group was fed barley-vetch silage *ad libitum* and limited concentrate, and the C group was fed barleygrass straw + alfalfa hay (50/50%) and limited concentrate feed. The trial was started in the final trimester period of the pregnancy and lasted 90 days. Liveweight and feed intake of the pregnant sheep were recorded in the last period of pregnancy. The liveweight of the lambs were measured at lambing and then on d 15 and 30. There is no diet effect on liveweights measure 1 and 2 months after lambing ($p > 0.05$). The liveweight value of sheep at lambing was 67.01 ± 1.32 , 64.07 ± 1.54 , and 64.89 ± 1.48 kg for the groups H, S, and C, respectively. The birth type was significant on birth weight ($p < 0.05$). Lamb liveweights on d 15 and 30 for the group H was greater than those for the group S ($p < 0.05$). Lambs' birth weight and weights on d 15 and 30 were 4.94 ± 0.13 , 4.44 ± 0.15 , and 4.77 ± 0.17 kg; 9.30 ± 0.30 , 8.03 ± 0.30 , and 8.94 ± 0.34 , 13.36 ± 0.43 , 10.85 ± 0.56 , and 12.37 ± 0.41 for groups respectively. In conclusion, barley-vetch mixture can be used as hay and silage in pregnant sheep feeding, being hay superior to silage.

Key words: Barley-vetch, silage, lamb performance

1. Introduction

Animal food has an important role in human nutrition. Among these foods, meat, milk, and eggs are the main ones [1,2]. It is estimated that the world population will be 9 billion in 2050 and the need for animal products will increase by 60-70% [3,4]. Also, the global pandemic conditions affected animal production negatively. Therefore, in order to increase the yield and quality of livestock enterprises, alternative production models should be applied, and the use of existing crops should be integrated [5,6]. In order to maintain the current production, it is necessary to increase the number of animals or increase the productivity [7]. Considering these facts, adequate nutrition intake of animals should be ensured in order to increase the livestock products [8].

Small ruminant has an important role in meat and milk production in Turkey as well as in the world [9,10]. Mutton production occupies a larger place among the small ruminant. Meat obtained from small ruminant meets 11.3% of the world's total meat production and 25% in Turkey [4]. Serious yield can be achieved due to

the increase of the yield per unit animal. Feeding of sheep is mostly based on pasture. Important periods in sheep nutrition include flushing, gestation, and lactation periods [11]. The growth of fetus to about 70% of adult growth takes place in the last 1/3 period of the pregnancy. The quality and adequate feed should be offered to the animals in this period [12].

The wheatgrass straw and dry grass are preferred as roughage for feeding pregnant sheep in Turkey conditions. [13]. However, wheatgrass straw is not enough to meet the nutritional needs of the animals. Therefore, it is necessary to increase the production of quality roughage resources to meet the needs of livestock enterprises. Considering the reasons such as decreasing water resources in the world, climate change, and the deterioration of soil structure, the production and use of roughage resources, which require minimum water, should be increased [14]. For this reason, two croppings and evaluation of fallow fields are widely used around the world [15,16–17]. One of the roughage sources is the vetch that can be evaluated for this purpose. Vetch can be fed to animals in the form of grass, silage, or

* Correspondence: hasansenyuzvet@yahoo.com

grain [18]. It is recommended that vetch is to be planted as a mixture with grains, as the stem problem occurs during the end of the flowering period [19]. It is preferred to plant it together with barley or oats and obtain successful results from ensiled as grass-legume hay [20]. It could be more appropriate to consider it as silage, due to the drying problems in the heavy rainy regions [21]. There are many studies on the ratio of vetch and wheatgrass mixed plantings, and although regional differences are important, it is recommended to use 60%–75% of vetch [22,23,24].

Studies on sheep and lambs are generally dealt with grain vetch [25,26]. The study on Akkaraman lambs and yards performance [27], there was no difference between the performance of the animals that consumed *ad libitum* vetch+barley and 500 g/d concentrate. In another study examining the effects of oat vetch on the fattening performance of weaned lambs [28], lambs fed oat vetch had better daily liveweight gain, and body weight at the end of fattening, as well as better carcass weight, and carcass efficiency as compared to the lambs that were grazing on natural pasture. In a comparison of lentil straw and vetch straw to alfalfa hay and wheatgrass straw [29], it was observed that sheep fed vetch straw gained more body weight, and that the nutritional value of vetch straw was much better than wheatgrass straw. Furthermore, it was stated that the use of oat+vetch hay was economical [30], and 100 g concentrate and 200 g oat vetch mixtures added to pasture were suitable and profitable in lamb fattening [31]. In a study examining the effect of the vetch, which was left in the field to analyze the lamb fattening performance [32], the lambs were allowed to graze on vetch stubble, 50/50% vetch+wheat stubble, and wheat stubble for 6 h, and it was observed that the lambs grazing only on vetch and vetch+wheat stubble gained 3 times more liveweight than those grazing only on wheat stubble. The purpose of this study was to investigate the form of barley-vetch as hay and silage in pregnant sheep feeding and its carry-over effects on lamb performance.

2. Material and methods

This study was carried out by International Center for Livestock Research and Training Ethical Committee Report No:167. The animal material of the study was obtained from the Department of Small Ruminants, International Center for Livestock Research and Training (ICLRT), which was comprised of 3–5 years old, 48 Lalahan (Kivircik x Akkaraman, B₁) sheep and lambs. Sheep were blocked by the age (3,4 and 5) and assigned into three diets: the control group was fed barleygrass straw + alfalfa hay (50/50%) (C, n = 16), barley-vetch hay (H, n = 16), and barley-vetch silage (S, n = 16). All groups had limited concentrate. Ages, initial liveweights, and days of gestation are given in Table 1.

Table 1. The age, initial liveweight, and day of gestation of sheep.

Groups	Age	LWMS (kg)	Gestation (d)
Hay	4.63 ± 0,29	57.63 ± 1,45	98.44 ± 3,07
Silage	4.53 ± 0,29	57.04 ± 1,37	95.67 ± 3,57
Control	4.79 ± 0,30	57.80 ± 1,16	103.36 ± 3,81
<i>P value</i>	0.833	0.918	0.308

LWMS: Live weight at the mating season. Data are Mean ± SE. N per group is 16.

The nutrient requirements of animals [33] were determined after a week of an adaptation period. The composition of the concentrate and ingredients of rations are given in Table 2.

The roughages were planted as 70% Hungarian vetch and 30% barley (Tarım 92) in the fields of ICLRT in November and harvested in June. Barleygrass straw was obtained from ICLRT and alfalfa hay local market. Fresh grass and hay yield was calculated during the harvest time. It was calculated by throwing a circle on an area of 1 m². Approximately 3 tons of the grasses were cut in 2–4 cm sizes in the silage machine and then ensiled using the classical method. The other grass was dried in the field for 1 day before baling.

After grinding to pass a 1 mm sieve, the feed samples were analyzed for DM, OM, CA, CP, and EE [34]. Neutral detergent fiber, ADF, ADL, and CS was analyzed according to Van Soest et al. [35] using Ankom 200. Organic acid levels (lactic, acetic, propionic, butyric, isobutyric, isovaleric acid) were determined in high-performance liquid chromatography (HPLC, Hewlett Packard, Series 1100) [36] after preparing silage samples as outlined by Tjardes et al. [37]. The metabolic energy of the concentrates was calculated according to TSE 9610 [38] modified by Sauvante and Morand [39].

To determine in situ DM, OM, and CP of barley-vetch, 3 ruminally cannulated Holstein cows were used. Approximately 5 g sample with 2 mm particle size introduced into nylon bag with 45 µ pore size. Samples were placed to rumen for 0, 2, 4, 8, 16, 24, 48, and 72 h for incubation [40]. The DM, OM, and CP degradability of feeds were calculated with following equation: $a+b(1-e^{-ct})$. Also in vitro OM digestibility of silage samples was also determined based on Tilley and Terry [41] modified by Marten and Barnes [42] using the Daisy II incubator (ANKOM®, Fairport, New York, USA).

After testing for normality, the difference between the groups and their interactions by the time were analyzed using one-way ANOVA. The group differences were attained using the Tukey test in commercial software (Minitab16) [43].

Table 2. Composition of concentrate and ration of consumed feed by animals.

Concentrate %						
Feed	Barley	Wheat bran	Sun flower seed meal	Calcium	Salt	Vit-Min
Rate (%)	62	17.4	18	2	0.5	0.1
Feed Consumption, g						
	1. Month			2. and 3. Months		
Feeds (g)	Hay	Silage	Control	Hay	Silage	Control
Concentrate	875	875	875	1000	1000	1000
Barley-Vetch hay	1500	-	-	1500	-	-
Barley-Vetch silage	-	2125	-	-	2125	-
Alfalfa hay	-	-	750	-	-	750
Barleygrass straw	-	-	625	-	-	625

3. Results

The amount of fresh hay of barley-vetch was 2120.50 kg/da, and the hay yield was 782.70 kg/da. Dry matter, nature DM, CA, CP, EE, NDF, ADF, ADL, ME, and in vitro digestibility of feeds are given in Table 3. Initial liveweights, age, and day of pregnancy were similar. Liveweights of sheep at lambing and on d 15 and 30 are given in Table 4. The consumption of nutrients and requirements of sheep [33] at the beginning of the experiment, late gestation, and early lactation period are given in Table 5.

4. Discussion and conclusion

Drying and silage are important storage method for green roughage. In this study, the effects of barley-vetch hay and silage were investigated on the live weight of pregnant sheep. Although it is considered that the fresh yield varies according to the climate and region, the fresh yield was obtained in the study was higher than Acar and Mülayim [44] and Ay and Mut [45], and similar to Balabanlı et al [6]. The reason for this difference may depend on the climatic structure of the regions and the planting period) must remove. A high DM value is related to the harvest time. The DM of the barley-vetch hay was similar with Bingöl et al. [5], and Civaner [8].

No difference is expected between fresh and silage DM. However, there was a difference in DM because the time between the cutting and ensiling was prolonged. It is important to silo in a short time after the cutting. The silage DM was higher in the study than in the former studies. [1, 46, 47].

The CA, EE, NDF and ADF values of barley-vetch hay and silage were similar in the study. Only the CP was higher in barley-vetch hay. In general, the CP was lower than similar studies [5, 45, 47]. It is thought that the low

CP is associated with the DM. The CA, NDF, and ADF was similar to the former studies [1, 5, 6]. The literature [1, 5] supports that the CP, NDF and ADF levels are related to the cutting time.

In vitro digestibility indicates that the digestibility of feeds. Digestibility decreases when NDF and ADF increased in the feeds [12]. In vitro digestibility of barley-vetch hay was higher than silage. It is thought to be due to the high NDF and ADF content of the silage. The literature [5, 48] supports that the low IVD values of feeds with high NDF and ADF content.

In situ digestibility is an important indicator for determining the digestibility levels of feeds in the rumen. In situ OM digestibility of barley-vetch hay and silage was similar. The 0th hour in situ CP digestibility of silage was higher than hay. This suggests that the NPN content of silage was high. Total CP digestibility of hay was higher than silage. This can be explained by the difference of NDF and ADF. In situ digestibility was lower than Karşlı et al. 2006 [49] and Turgut et al. [50].

Lamb development of 70% takes place in the last 1/3 of pregnancy [12]. Therefore, feeding in the last 1/3 of pregnancy affects of lamb development and birth weight. There was no difference between the groups in terms of birth weight of lambs ($p>0.05$). It was observed that the roughage type did not affect the birth weight. However, hay group lambs were better development in terms of 15th and 30th day liveweight ($p < 0.05$). This situation is thought to be due to the net nutrient consumed by the animal is high due to the low NDF and ADF content of barley-vetch hay. In the literature, sheep fed vetch straw were gained more liveweight and better digested than the fed wheat straw [50], and lambs fed vetch in different varieties were better than those fed alfalfa in terms of nutrient digestibility, live

Table 3. Chemical composition, in vitro digestibilities (%), and in situ degradabilities of feeds.

	DM	NDM	CA	CP	EE	NDF	ADF	ADL	ME (Mcal/kg)
Concentrate	89.65		3.93	13.28	2.13	29.9	12.39	2.61	2.9
Fresh barley-vetch		36.92	7.12	9.94	2.33	51.86	35.69	3.86	1.75
Barley-vetch hay	92.21		6.56	8.75	1.37	54.85	33.75	4.42	1.72
Barley-vetch silage		59.63	5.28	7.21	1.17	59.87	37.3	4.95	1.64
Alfalfa hay	91.96		12.71	13.72	1.76	50.09	38.74	8.55	2.08
Barleygrass straw	90.22		7.27	6.33	1.70	68.56	48.01	8.92	1.55
In-vitro Digestibility									
	IVDMD		IVCPD		IVNDFD		IVOMD		
Barley-vetch hay	65.22 ± 0,79		83.70 ± 0.29		44.83 ± 1.03		65.78 ± 1.08		
Barley-vetch silage	58.80 ± 1,18		70.45 ± 0.69		53.62 ± 1.50		60.06 ± 0.57		
In situ OM Degradability									
Groups	0	2	4	8	16	24	48	72	
BVH	32.71	43.94	47.49	51.83	58.75	60.34	69.73	71.85	
BVS	32.80	40.27	43.36	49.15	53.83	56.00	68.71	71.85	
In situ CP Degradability									
BVH	52.10	70.47	74.22	78.73	79.51	80.22	80.88	80.88	
BVS	63.79	64.79	68.70	74.11	74.88	75.02	75.02	75.17	

DM; Dry matter, NDM; nature dry matter, CA; crude ash, CP; crude protein, NDF; neutral detergent fiber, ADF; acid detergent fiber, ADL; acid detergent lignin, ME; metabolic energy.

IVDMD; in vitro dry matter digestibility, IVCPD; in vitro crude protein digestibility, IVNDFD; in vitro neutral detergent fiber digestibility, IVOMD; in vitro organic matter digestibility.

BVH: Barle-vetch hay, BVS: barley-vetch silage.

Table 4. Liveweights of sheep at lambing and on d 15 and 30.

Groups	n	Birth LW (kg)	n	15 th Day LW (kg)	n	30 th Day LW (kg)
Hay	16	4.94 ± 0.13	15	9.30 ± 0.30 ^a	12	13.36 ± 0.43 ^a
Silage	21	4.44 ± 0.15	12	8.03 ± 0.29 ^b	6	10.85 ± 0.56 ^b
Control	18	4.77 ± 0.17	12	8.94 ± 0.34 ^{ab}	11	12.37 ± 0.41 ^{ab}
P values		0.064		0.018		0.006

weight gain, and performance [48]. Again, Haddad and Husein [29] reported that vetch hay instead of wheat straw gives better results on live weight gain of lambs. It has been reported that 33% and more vetch hay has a positive effect on lamb performance in the fattening period [51].

There was no difference between the groups in terms of live weight of sheep during early pregnancy, late pregnancy, and early lactation period. All groups had live

weight gain in accordance with the gestational period. The DM, CP and ME consumptions of the animals were higher in the group H than in the group S. Increasing the DM consumption has positive effects on the development of pregnant animals. Since feed intake is inversely proportional to NDF, DM consumption was higher in hay group. In addition, dry matter consumption is higher in rations which ingredient 50% moisture roughage [33]. In

Table 5. Nutrient consumption and requirements of sheep in the beginning of fattening, late gestation, and early lactation periods (38).

Beginning of fattening (pregnancy of 100. day)													
Groups	n	Gestation Day	LW (Kg)	LWG (kg)	RDM (Kg)	CDM (Kg)	DMCR %	RCP (g)	CCP (g)	CPCR %	RME (g)	CME (g)	MECR %
Hay	16	98.44 ± 3.07	61.08 ± 1.34	0.034 ± 0.01	1.69 ± 0.03	1.83 ± 0.04 ^a	1.08 ± 0.01	128.94 ± 4.88	184.39 ± 4.03 ^a	1.45 ± 0.05 ^a	3.05 ± 0.11	3.64 ± 0.08 ^a	1.21 ± 0.04 ^a
Silage	16	95.67 ± 3.57	60.60 ± 1.25	0.037 ± 0.01	1.64 ± 0.04	1.67 ± 0.03 ^b	1.03 ± 0.02	136.67 ± 5.12	153.12 ± 3.16 ^b	1.14 ± 0.05 ^b	3.18 ± 0.12	3.24 ± 0.07 ^b	1.04 ± 0.04 ^b
Control	16	103.36 ± 3.81	61.77 ± 1.58	0.037 ± 0.01	1.65 ± 0.04	1.73 ± 0.04 ^{ab}	1.05 ± 0.03	136.93 ± 6.66	190.02 ± 4.86 ^a	1.42 ± 0.06 ^a	3.15 ± 0.15	3.20 ± 0.08 ^b	1.03 ± 0.04 ^b
P values		0.308	0.841	0.941	0.571	0.022	0.127	0.506	0.001	0.001	0.734	0.001	0.001
Late Gestation													
Hay	16	122.44 ± 3.07	68.18 ± 1.47	0.30 ± 0.02	1.81 ± 0.03	2.17 ± 0.05 ^a	1.20 ± 0.01 ^a	144.81 ± 4.11	225.20 ± 4.86 ^a	1.56 ± 0.03 ^a	3.37 ± 0.11	4.44 ± 0.10 ^a	1.32 ± 0.03 ^a
Silage	15	119.67 ± 3.57	67.34 ± 1.53	0.28 ± 0.02	1.82 ± 0.03	1.91 ± 0.04 ^b	1.05 ± 0.01 ^c	148.73 ± 5.01	191.28 ± 4.35 ^b	1.30 ± 0.03 ^b	3.49 ± 0.13	3.62 ± 0.08 ^b	1.05 ± 0.03 ^c
Control	14	127.36 ± 3.81	68.06 ± 1.90	0.26 ± 0.03	1.81 ± 0.04	2.05 ± 0.06 ^{ab}	1.13 ± 0.01 ^b	146.57 ± 6.39	230.21 ± 6.44 ^a	1.59 ± 0.04 ^a	3.43 ± 0.16	3.94 ± 0.11 ^b	1.16 ± 0.03 ^b
P values			0.925	0.626	0.975	0.002	0.001	0.862	0.001	0.001	0.815	0.001	0.001
Early Lactation													
Hay	16	149.75 ± 0.54	67.01 ± 1.32		1.92 ± 0.05	2.28 ± 0.05 ^a	1.19 ± 0.02 ^a	209.69 ± 7.80	240.08 ± 4.71 ^a	1.16 ± 0.03 ^a	3.55 ± 0.11	4.72 ± 0.09 ^a	1.34 ± 0.03 ^a
Silage	15	152.67 ± 1.74	64.07 ± 1.54		1.88 ± 0.05	2.02 ± 0.05 ^b	1.08 ± 0.02 ^b	225.80 ± 9.14	206.17 ± 4.96 ^b	0.93 ± 0.04 ^b	3.75 ± 0.13	3.91 ± 0.09 ^b	1.05 ± 0.03 ^c
Control	14	149.86 ± 0.82	64.89 ± 1.48		1.85 ± 0.06	2.16 ± 0.05 ^{ab}	1.18 ± 0.02 ^a	217.60 ± 11.70	245.09 ± 5.60 ^a	1.15 ± 0.04 ^a	3.66 ± 0.16	4.23 ± 0.10 ^b	1.17 ± 0.03 ^b
P values		0.137	0.330		0.581	0.002	0.001	0.485	0.001	0.001	0.570	0.001	0.001

LW; live weight, LWG; live weight gain, RDM; requirements of dry matter, CDM; consumption of DM, DMCR; DM coverage rate, RCP; requirements of crude protein, CCP; consumption of CP, CPCR; CP coverage rate, RME; requirements of metabolic energy, CME; consumption of ME, MECR; ME coverage rate. Data are Mean ± SE.

this study, DM consumption was lower in the group S due to high amount of moisture. This agrees with the literature [33]. The pregnant sheep were fed with alfalfa and vetch hay [52]; it was stated that no adverse events were found, and vetch can be given safely.

In summary, barley-vetch mixture hay and silage are quality roughage sources for sheep. Both forms can be used safely alone in the feeding of pregnant sheep, and it does not cause any health problems or low productivity. However, considering lamb development, the H form was superior to the S form.

References

- Kennedy PC, Dawson LER, Lively FO, Steen RWJ, Fearon AM, et al. Effects of offering lupins/triticale and vetch/barley silages alone or in combination with grass silage on animal performance, meat quality and the fatty acid composition of lean meat from beef cattle. *The Journal of Agricultural Science* 2018; 156 (8): 1005-1016.
- Şenyüz HH, Erat S, Karlı MA, Soydemir İ. Comparison of fattening performance of Angus, Charolais, Limousine and Simmental cattle imported to Turkey. *Livestock Studies* 2020; 60 (1): 1-4. doi: 10.46897/lahaed.779712
- Makkar HPS, Tran G, Heuz'e V, Giger-Reverdin S, Lessire M, et al. Seaweeds for livestock diets: a review. *Animal Feed Science and Technology* 2015; 212: 1-17. doi: 10.1016/j.anifeedsci.2015.09.018
- FAO. (2019). FAO Statistics.
- Bingöl NT, Karlı MA, Yılmaz IH, Bolat D. The effects of planting time and combination on the nutrient composition and digestible dry matter yield of four mixtures of vetch varieties intercropped with barley. *Turkish Journal of Veterinary and Animal Sciences* 2007; 31 (5): 297-302.
- Balabanlı C, Albayrak S, Türk M, Yuksel O. A research on determination of hay yields and silage qualities of some vetch+cereal mixtures. *Turkish Journal of Field Crops* 2010; 15(2): 204-209.
- Karlı MA, Bingöl NT. The determination of planting density on herbage yield and silage quality of Jerusalem artichoke (*Helianthus tuberosus* L.) green mass. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 2009; 15 (4): 581-586.
- Civaner AG. Batı Akdeniz bölgesinde yetiştirilen bazı yem hammaddelerinin besin madde içeriklerinin belirlenmesi üzerine bir araştırma. PhD, Akdeniz University, Antalya, Turkey, 2015.
- Aygün T, Çelikyürek H. Recommended sheep husbandry model and body weights of Morkaraman lambs in Muş province. *Livestock Studies* 2020; 60 (1): 005-008. doi: 10.46897/lahaed.661067
- Güngör İ, Atasoy F. Use of multiple regression modeling for the evaluation of lactation characteristics of Awassi sheep. *Livestock Studies* 2020; 60(2): 46-53. doi: 10.46897/livestockstudies.846408
- Umberger SH. Feeding sheep. Virginia Cooperative Extension, Publication 2009; 410-853.
- Ergün A, Çolpan İ, Yıldız G, Küçükersan S, Tuncer ŞD et al. Hayvan Besleme ve Beslenme Hastalıkları., Geliştirilmiş 5. Baskı., 975-97808-2-8, Ankara, Turkey. Pozitif Yayıncılık; 2011.
- Bingöl NT, Bolat D, Karlı MA, Akça İ. Arpa hasılı ve korunga karışımı silaja farklı düzeylerde melas ilavesinin silaj kalitesi ve sindirilebilirliği üzerine etkileri. *Atatürk Üniversitesi Veteriner Bilimleri Dergisi* 2009; 4(1): 23-30. (in Turkish with an abstract in English)
- Huang YF, Gao XL, Nan ZB, Zhang ZX. Potential value of the common vetch (*Vicia sativa* L.) as an animal feedstuff: a review. *Journal of animal physiology and animal nutrition* 2017; 101(5): 807-823. doi: 10.1111/jpn.12617
- Garduño-Castro Y, Espinoza-Ortega A, González-Esquível CE, Mateo-Salazar B, Arriaga-Jordán CM. Intercropped oats (*Avena sativa*)-common vetch (*Vicia sativa*) silage in the dry season for small-scale dairy systems in the Highlands of Central Mexico. *Tropical animal health and production* 2009; 41 (5): 827-834. doi: 10.1007/s11250-008-9258-7
- Chen L, Guo G, Yu C, Zhang J, Shimojo M et al. The effects of replacement of whole-plant corn with oat and common vetch on the fermentation quality, chemical composition and aerobic stability of total mixed ration silage in Tibet. *Animal Science Journal* 2015; 86 (1): 69-76. doi: 10.1111/asj.12245
- İke F. Laktik asit bakterisi ve enzim ilavesinin adi fiğ, buğday, yulaf karışımı silajlarda fermentasyon ile aerobik stabilite özellikleri üzerine etkileri. MSc, Namık Kemal University, Tekirdağ, Turkey, 2019.

Acknowledgment

This study was funded by the General Directorate of Agricultural Research and Policies (GDARP) (Project number: TAGEM/HSGYAD/B/20/A3/P1/2085). Authors thank GDARP and International Center for Livestock Research and Training for supporting the project and animal material support.

Credit

This manuscript was presented orally at 6th International Congress on Advances in Veterinary Sciences & Technics (23-27 August 2021) (Abstract).

18. Keles G, Demirci U. The effect of homofermentative and heterofermentative lactic acid bacteria on conservation characteristics of baled triticale–Hungarian vetch silage and lamb performance. *Animal Feed Science and Technology* 2011; 164 (1-2): 21-28. doi: 10.1016/j.anifeeds.2010.11.017
19. Aşçı Ö, Eğritaş Ö. Yaygın fiğ-tahıl karışımlarında ot verimi, bazı kalite özellikleri ve rekabetin belirlenmesi. *Journal of Agricultural Sciences* 2017; 23 (2): 242-252. (in Turkish with an abstract in English)
20. Bingol NT, Karsli MA, Bolat D, Akca I. Effect of molasses and sulphuric acid addition to barley/hungarian vetch bi-crop silages. *Journal of Applied Animal Research* 2006; 30 (1): 29-32. doi: 10.1080/09712119.2006.9706819
21. Çerçi İH, Şahin K, Güler T. Farklı oranlarda silajlık mısır yonca kullanılarak yapılan silajların kalitesinin belirlenmesi. *Fırat Üniversitesi Sağlık Bilimleri Dergisi* 1996; 10 (2): 193-200. doi: 10.29278/azd.737554
22. Tan M. Fiğ+ tahıl karışımları için en uygun karışım oranları ve biçim zamanının belirlenmesi. PhD, Atatürk University, Erzurum, Turkey, 1995.
23. Yılmaz Ş, Özel A, Atak M, Erayman M. Effects of seeding rates on competition indices of barley and vetch intercropping systems in the Eastern Mediterranean. *Turkish Journal of Agriculture and Forestry* 2015; 39 (1): 135-143.
24. Leilei L, Hua D, Zheng X, Li C. Effects of moisture content and sowing-mix ratio on the quality of baled oat and common vetch/hairy vetch silage mixtures in the pastoral area of southern Qinghai. *Acta Prataculturae Sinica* 2018; 27 (7): 166-174.
25. Kaya İ, Yalçın S. The effects of rations containing different amounts of common vetch seed on growth performance, digestibility and some blood and rumen metabolites in male lambs. *Turkish Journal of Veterinary and Animal Sciences* 2000; 24 (4): 307-316. (in Turkish with an abstract in English)
26. Gül M, Yörük MA, Özudoğru Z, Timurkan S. İvesi kuzu rasyonlarına değişik oranlarda fiğ (*vicia sativa*) ilavesinin besi performansına ve karaciğerin histo-patolojisi üzerine etkisi. *Yüzüncü Yıl Üniversitesi Veteriner Fakültesi Dergisi* 2003; 14 (2): 6-9. (in Turkish with an abstract in English)
27. Karabulut A, Munzur M. Nadas alanlarına ekilen farklı karışımlar üzerinde otlatılan toklu ve süttan kesilmiş kuzuların besi güçleri üzerinde araştırmalar. *Uludağ Üniversitesi Ziraat Fakültesi Dergisi* 1987; 6: 47-55. (in Turkish)
28. Lema S. Comparison of oats/vetch fodder crop and natural pasture for fattening Ethiopian highland sheep. In 3. National Conference of the Ethiopian Society of Animal Production, Addis Abeba (Ethiopia), 27-29 Apr 1995. ESAP.
29. Haddad SG, Husein MQ. Nutritive value of lentil and vetch straws as compared with alfalfa hay and wheat straw for replacement ewe lambs. *Small Ruminant Research* 2001; 40(3): 255-260. doi: 10.1016/S0921-4488(01)00176-6
30. Mekonen T, Hailu A, Tigabe A, Abebe A, Alemayehu L. Demonstration of fattening of yearling sheep with vetch hay in gerakeya worda. *Results of Livestock Research* 2016; 129-134.
31. Mengistu S, Hawassa E. Assessment of forage production, feed resource utilization and substitution effect of oat-vetch forage for concentrate mix on performance of sheep fed desho grass as a basal diet in Damote Gale District of Wolaita Zone, SNNPR PhD, Hawassa University, Ethiopia, 2018.
32. Abidi S, Benyoussef S, Ben Salem H. Foraging behaviour, digestion and growth performance of sheep grazing on dried vetch pasture cropped under conservation agriculture. *Journal of Animal Physiology and Animal Nutrition* 2021; 105(1): 51-58. doi: 10.1111/jpn.13456
33. National Research Council, Committee on the Nutrient Requirements of Small Ruminants, Board on Agriculture, Division on Earth, & Life Studies. *Nutrient requirements of small ruminants: sheep, goats, cervids, and new world camelids*, 2007.
34. AOAC (1995). *Association of Official Analytical Chemists. Official Methods of Analysis (16th Edition)*. Arlington. Virginia. USA.
35. Van Soest PJ, Robertson JB, Lewis BA. Methods for dietary fiber, neutral detergent fiber and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 1991; 74: 3583-3597. doi: 10.3168/jds.S0022-0302(91)78551-2
36. Canale A, Valente ME, Ciotti A. Determination of volatile carboxylic acids (C1-C5i) and lactic acid in aqueous acid extracts of silage by high-performance liquid-chromatography. *Journal of the Science of Food and Agriculture* 1984; 35: 1178-1182. doi: 10.1002/jsfa.2740351106
37. Tjardes KE, Buskirk DD, Allen MS, Ames NK, Bourquin LD et al. Brown midrib-3 corn silage improves digestion but not performance of growing beef steers. *J. Anim. Sci.* 2000; 78: 2957-2965.
38. TSE 1994. TS 9610 Animal Feeds- Determination of Metabolizable Energy (Chemical Method)
39. Sauvant D, Morand-Fehr P. Analyse des variations individuelles en nutrition animale: application de l'analyse en composantes principales à l'étude de la sécrétion lipidique du lait de chèvre. In *Annales de zootechnie* 1978; 27 (1): 75-81.
40. Ørskov ER, McDonald I. The estimation of protein degradability in the rumen from incubation measurements weighted according to rate of passage. *The Journal of Agricultural Science* 1979; 92 (2): 499-503.
41. Tilley JMA, Terry DR. A two-stage technique for the in vitro digestion of forage crops. *Grass and forage science* 1963; 18(2): 104-111.
42. Marten GC, Barnes RF. Prediction of energy digestibility of forages with in vitro rumen fermentation and fungal enzyme systems. In *Standardization of analytical methodology for feeds: proceedings*. IDRC, Ottawa, ON, CA. 1979.
43. Minitab LLC. (2006). SAS, v. 19; Minitab. Inc.: State College, PA, USA.
44. Acar R, Mülayim M. Konyada bazı yem bitkilerinin doğrudan anıza ekim yöntemiyle ikinci ürün olarak yetiştirilmesi. *Bahri Dağdaş Bitkisel Araştırma Dergisi* 2014; 1(1-2): 20-25. (in Turkish with an abstract in English)

45. Ay İ, Mut H. Determination of suitable mixture ratio of common vetch and pea with oats and barley. COMU Journal of Agriculture Faculty 2017; 5 (2): 55-62. (in Turkish with an abstract in English)
46. Altınok S. Tüylü fiğ (*Vicia villosa* L.) ve koca fiğ (*Vicia narbonensis* L.)'in arpa (*Hordeum vulgare* L.) ile farklı oranlardaki karışımlarının silaj kalitesine etkileri. Tarım Bilimleri Dergisi 2002; 8 (3): 232-237. (in Turkish with an abstract in English)
47. Bengisu G. A study on the silage properties of hungarian vetch (*vicia pannonica* crantz.) and barley (*hordeum vulgare* L.) grass mixtures in different rates. Legume Research 2019; 42 (5): 680-683.
48. Huang YF, Matthew C, Li F, Nan ZB. Common vetch varietal differences in hay nutritive value, ruminal fermentation, nutrient digestibility and performance of fattening lambs. Animal 2021; 15(7): 100244. doi: 10.1016/j.animal.2021.100244.
49. Karşlı MA, Akdeniz H, Levendoğlu T, Terzioğlu Ö. Evaluation of the nutrient content and protein fractions of four different common vetch varieties. Turkish Journal of Veterinary and Animal Sciences (2006); 29 (6): 1291-1297.
50. Turgut L, Yanar M, Kaya A, Tan M. Farklı olgunluk dönemlerinde hasat edilen bazı fiğ türlerinin ham besin maddeleri içeriği ve bunların in situ rumen parçalanabilirlikleri. Atatürk Üniversitesi Ziraat Fakültesi Dergisi 2006; 37 (2): 181-186. (in Turkish with an abstract in English)
51. Gheshlagh, FMA, Benemar AH. Determination of nutritive value and feeding different levels of vetch-barley mixed culture on fattening performance of moghani male lamb. Research On Animal Production (Scientific and Research) 2016; 6 (12): 131-138.
52. Poland C, Faller T, Tisor L. Effect of chickling vetch (*Lathyrus sativus* L.) or alfalfa (*Medicago sativa*) hay in gestating ewe diets. Originally published in the Sheep Day Report, Hettinger RE Center 2003; 38-40.