

1-1-2022

## The effect of nongenetic factors on calf birth weight and growth performance in Anatolian buffaloes

KÜRŞAT ALKOYAK

SEZER ÖZ

Follow this and additional works at: <https://journals.tubitak.gov.tr/veterinary>



Part of the [Animal Sciences Commons](#), and the [Veterinary Medicine Commons](#)

---

### Recommended Citation

ALKOYAK, KÜRŞAT and ÖZ, SEZER (2022) "The effect of nongenetic factors on calf birth weight and growth performance in Anatolian buffaloes," *Turkish Journal of Veterinary & Animal Sciences*: Vol. 46: No. 4, Article 8. <https://doi.org/10.55730/1300-0128.4232>

Available at: <https://journals.tubitak.gov.tr/veterinary/vol46/iss4/8>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Veterinary & Animal Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## The effect of nongenetic factors on calf birth weight and growth performance in Anatolian buffaloes

Kürşat ALKOYAK<sup>\*</sup>, Sezer ÖZ

General Directorate of Agricultural Research and Policies, Ankara, Turkey

Received: 29.01.2022 • Accepted/Published Online: 13.07.2022 • Final Version: 03.08.2022

**Abstract:** The aim of the current study is to determine the growth performances of Anatolian buffaloes such as birth weight ( $BW_0$ ), body weight at 6 months ( $BW_6$ ), body weight at 12 months ( $BW_{12}$ ), average daily weight gain from birth to 6 months ( $ADG_{0-6}$ ), average daily weight gain from birth to 12 months ( $ADG_{0-12}$ ) and average daily weight gain from 6 to 12 months ( $ADG_{6-12}$ ), and to identify various nongenetic factors affecting these traits. In the study, the data of 2821 head buffalo calves born from buffaloes bred from the Anatolian Buffalo Breeding Project in Bartın province between 2015–2021 were used. In the present study, the effects of nongenetic factors such as district, calving year and season, calving age, and sex on  $BW_0$  and growth performance characteristics were investigated. The mean and standard errors of the  $BW_0$ ,  $BW_6$ ,  $BW_{12}$ ,  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  values of buffalo calves were calculated as  $28.33 \pm 0.090$  kg,  $119.13 \pm 0.459$  kg,  $173.53 \pm 0.743$  kg,  $504.64 \pm 2.31$  g,  $398.24 \pm 1.88$  g, and  $304.27 \pm 2.45$  g respectively. In the study, except for the effect of calving age on  $ADG_{0-6}$  and  $ADG_{6-12}$  ( $p > 0.05$ ), the effect of all other nongenetic factors on growth performance was found to be statistically significant ( $p \leq 0.05$ ). There is not enough research on environmental factors affecting growth performance at different ages in Anatolian buffaloes. The data on growth performances and significant nongenetic factors obtained in this study will be useful in the selection program and improvement in buffalo husbandry and enlighten future studies.

**Key words:** Anatolian buffalo, birth weight, daily weight gain, growth performance, nongenetic factors

### 1. Introduction

The existence of buffaloes in Turkey dates back to 3000 BC. Anatolian Buffalo breed is classified as Mediterranean buffalo among river-type buffaloes. This breed is spread all over Turkey, the majority of the buffalo population in Turkey is located in the Central Black Sea Region [1,2]. The current number of these animals is 192,489 and they are generally raised for meat and milk [3]<sup>1</sup>. Anatolian buffaloes are very resistant to sudden changes in feeding as well as diseases [4]. Buffalo milk is used in the production of cream, cheese, and ice cream. Buffalo meat is a promising market as it is preferred by consumers due to its excellent nutritional properties and taste and thus gaining popularity in many parts of the world [5,6,7]. However, only 0.38% and 0.007% of the milk and red meat produced from bovine animals in Turkey in 2019 are produced from buffaloes respectively, and these rates are quite low<sup>1</sup>. To increase milk and meat production, it is very important to know the factors affecting the growth performance as well as increasing the number of buffaloes.

Calf birth weight, which is an indicator of growth, is affected by genetic and nongenetic factors such as dam's age and calf sex in the prenatal period [8,9]. Birth weight ( $BW_0$ ) is associated with adaptation and survival [10], and is also used as an indicator feature in selection programs to reduce the risk of dystocia [11]. The growth characteristics of buffaloes are affected by many environmental factors as well as their genetic structure. Environmental factors can suppress the real growth potential of the animal and therefore make normal selection procedures ineffective [12]. In this respect, the success of the breeding program largely depends on understanding and knowing the relationship between genetic and environmental effects [13].

There is very little information in the literature about the growth performance of Anatolian buffaloes, especially about average daily weight gains (ADG) and the effect of nongenetic factors on growth performance. This study aimed to formulate stud selection and breeding programs and to contribute to future genetic studies by determining the effect of nongenetic factors affecting some growth

<sup>1</sup> TURKSTAT (2022). Turkish Statistical Institute Statistics. [online]. Website <http://www.turkstat.gov.tr/Start.do> [Accessed 01 Jan, 2022].

\* Correspondence: kursatalkoyak@gmail.com

performances of Anatolian buffaloes in farms where the breeding project is applied.

## 2. Material and method

### 2.1. Study area, animals, and data collection

The material of the study consisted of birth records of 2821 calves obtained from Anatolian buffaloes (41°38'4"N and 32°20'15"E) that gave birth in Bartın province between 2015 and 2021. The data on the growth performances of Anatolian buffaloes were obtained from the "Manda Yıldızı" program [14] in which the data were recorded within the scope of the National Project of Buffalo Breeding, supported by the General Directorate of Agricultural Research and Policies. Buffalo farming is carried out under extensive conditions in the area where the study was conducted. On days when the season is suitable, buffaloes are taken to the pasture in the morning and return in the evening. When the buffaloes are out on the pasture, additional feeding is not generally performed, but farmers do additional feeding according to the current feed (hay, silage dry alfalfa hay, etc.) in winter. In the farms within the scope of the breeding project where the study was carried out, the care and feeding methods for the buffaloes are generally similar to each other. The average number of female buffaloes per farm is 10 heads and the birth rate of buffaloes is approximately 44%. Within the farms, the buffalo cows are naturally inseminated by the bulls. Bulls are replaced every 2–3 years as for the project. Ear tags are applied to all buffalo calves born in the farms and their  $BW_0$  are weighed within 24 h with a digital scale (up to 10 g sensitive with a capacity of 50 kg). Meanwhile, the birth information (date of birth, sex, and dam's ear tag number) of the buffalo calves is taken and recorded. Buffalo calves suckle from their dams twice a day, morning and evening. After about 5–6 months of age, they go out to the pasture with their dams. Body weights of buffalo calves at the ages of 6 ( $BW_6$ ) and 12 months ( $BW_{12}$ ) are weighed and recorded with a scale (600 kg capacity, sensitive up to 100 g). Average daily weight gains from birth to 6 months ( $ADG_{0-6}$ ), from birth to 12 months ( $ADG_{0-12}$ ), and 6 to 12 months ( $ADG_{6-12}$ ) were calculated using these weights ( $BW_0$ ,  $BW_6$  ve  $BW_{12}$ ) [15,16]. The study was carried out in 4 districts of Bartın province; (1) Amasra, (2) Kurucaşile, (3) Center, and (4) Ulus. According to the climatic conditions, calving seasons were divided into four groups; (1) winter (December, January, and February), (2) spring (March, April, and May), (3) summer (June, July, and August), and (4) fall (September, October, and November). The calving age was divided into six groups: (1) age  $\leq$  4 years, (2) 4 < age  $\leq$  6, (3) 6 < age  $\leq$  8, (4) 8 < age  $\leq$  10, (5) 10 < age  $\leq$  12, (6) 12 < age.

### 2.2. Statistical analysis

In this study, the effects of the district, calving year and season, age, and sex on the growth performance of the

nongenetic factors were determined by using the "Least Squares Method". Checks for statistical significance of mean values were made by analysis of variance, and differences between significant means were made by Tukey's multiple comparison test. Due to insufficient data in subgroups, two or three-way interactions between the factors were not included in the analysis. The GLM (General Linear Model) method in the "Minitab-Version 18" program package was used for the statistical analysis of all data [17]. The effects of nongenetic factors examined in the present study on some growth characteristics ( $BW_0$ ,  $BW_6$ ,  $BW_{12}$ ,  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$ ) were investigated using the model below.

$$Y_{ijklmn} = \mu + D_i + Y_j + S_k + A_l + G_m + e_{ijklmn}$$

Where;

$Y_{ijklmn}$ : Level of productivity feature of any buffalo (i. district, j. year, k. season l. age, m. the observation value of the feature emphasized on sex)

$\mu$ : General (expected) mean,

$D_i$ : Effect of  $i^{\text{th}}$  district ( $i = 1,2,3,4$ )

$Y_j$ : Effect of  $j^{\text{th}}$  calving year ( $j = 2015, 2016, 2017, 2018, 2019, 2020$ ),

$S_k$ : Effect of  $k^{\text{th}}$  calving season ( $k = 1,2,3,4$ ),

$A_l$ : Effect of  $l^{\text{th}}$  calving age ( $l = 1,2,3,4,5,6$ )

$G_m$ : Effect of  $m^{\text{th}}$  sex ( $m = \text{male, female}$ )

$e_{ijklmn}$ : Random error which is assumed to be normally independently distributed with zero mean and constant variance ( $NID, 0, \sigma_2$ ).

## 3. Results

The effect of the nongenetic factors on these growth performances and the least squares averages are shown in Table 1 and Table 2. In this study,  $BW_0$ ,  $BW_6$ ,  $BW_{12}$  overall mean and standard error were determined as  $28.33 \pm 0.090$ ,  $119.13 \pm 0.459$ , and  $173.53 \pm 0.743$  kg, respectively (Table 1). The overall mean and standard error of  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  were  $504.64 \pm 2.31$ ,  $398.24 \pm 1.88$ , and  $304.27 \pm 2.45$  g, respectively (Table 2). The effects of nongenetic factors such as district, calving year, season, calving age and sex on these features were determined. Except for the effect of calving age on  $ADG_{0-6}$  and  $ADG_{6-12}$  ( $p > 0.05$ ), the effect of all other nongenetic factors on growth performance was found to be significant ( $p < 0.05$ ,  $p < 0.01$ ,  $p < 0.001$ ).

## 4. Discussion

The  $BW_0$  value found in the present study ( $28.33 \pm 0.090$  kg) (Table 1) is higher than the values in the research by Uğurlu et al. [18] ( $26.95 \pm 0.25$  kg) on Anatolian buffaloes. On the other hand, it is compatible with the study performed by Kul et al. [19] ( $29.3 \pm 0.43$  kg) on Anatolian buffaloes in Turkey. However, this value found in our research is lower than the  $BW_0$  found by many other researchers [20–27].

**Table 1.** Least squares means ( $\pm$  SE) of various growth periods in Anatolian buffaloes according to the district, calving year, season, age, and sex.

Nongenetic factors	BW <sub>0</sub> (kg)		BW <sub>6</sub> (kg)		BW <sub>12</sub> (kg)	
	n	Mean $\pm$ SEM	n	Mean $\pm$ SEM	n	Mean $\pm$ SEM
<b>Overall means</b>	2821	28.33 $\pm$ 0.090	2536	119.13 $\pm$ 0.459	2091	173.53 $\pm$ 0.743
District		***		***		***
Amasra	144	27.50 $\pm$ 0.210 <sup>c</sup>	126	114.72 $\pm$ 1.080 <sup>c</sup>	97	165.78 $\pm$ 1.650 <sup>c</sup>
Kurucaşile	132	29.16 $\pm$ 0.222 <sup>a</sup>	124	125.75 $\pm$ 1.100 <sup>a</sup>	113	184.42 $\pm$ 1.540 <sup>a</sup>
Centrum	2193	28.52 $\pm$ 0.061 <sup>b</sup>	2000	117.94 $\pm$ 0.308 <sup>b</sup>	1646	172.73 $\pm$ 0.532 <sup>b</sup>
Ulus	352	28.13 $\pm$ 0.140 <sup>c</sup>	286	118.13 $\pm$ 0.740 <sup>b</sup>	235	171.20 $\pm$ 1.120 <sup>b</sup>
<b>Calving year</b>		***		***		***
2015	287	26.00 $\pm$ 0.172 <sup>c</sup>	270	111.43 $\pm$ 0.854 <sup>c</sup>	243	160.36 $\pm$ 1.210 <sup>d</sup>
2016	405	27.00 $\pm$ 0.147 <sup>d</sup>	390	117.17 $\pm$ 0.725 <sup>d</sup>	373	167.72 $\pm$ 0.994 <sup>c</sup>
2017	518	28.27 $\pm$ 0.135 <sup>c</sup>	502	119.88 $\pm$ 0.664 <sup>bc</sup>	452	176.97 $\pm$ 0.937 <sup>ab</sup>
2018	555	28.62 $\pm$ 0.130 <sup>c</sup>	518	118.80 $\pm$ 0.644 <sup>cd</sup>	469	175.55 $\pm$ 0.894 <sup>b</sup>
2019	534	29.42 $\pm$ 0.129 <sup>b</sup>	526	121.88 $\pm$ 0.631 <sup>b</sup>	502	179.30 $\pm$ 0.873 <sup>a</sup>
2020	522	30.66 $\pm$ 0.131 <sup>a</sup>	330	125.65 $\pm$ 0.766 <sup>a</sup>	52	181.30 $\pm$ 2.290 <sup>ab</sup>
<b>Calving season</b>		*		***		**
Winter	457	28.25 $\pm$ 0.137 <sup>ab</sup>	377	117.92 $\pm$ 0.706 <sup>b</sup>	339	173.43 $\pm$ 1.010 <sup>ab</sup>
Spring	637	28.13 $\pm$ 0.129 <sup>b</sup>	592	118.18 $\pm$ 0.650 <sup>b</sup>	507	171.40 $\pm$ 0.980 <sup>b</sup>
Summer	969	28.43 $\pm$ 0.109 <sup>ab</sup>	904	121.27 $\pm$ 0.550 <sup>a</sup>	733	175.05 $\pm$ 0.895 <sup>a</sup>
Fall	758	28.50 $\pm$ 0.118 <sup>a</sup>	663	119.17 $\pm$ 0.610 <sup>b</sup>	512	174.25 $\pm$ 0.980 <sup>a</sup>
<b>Calving age (year)</b>		***		***		***
3–4	499	26.92 $\pm$ 0.134 <sup>d</sup>	456	116.74 $\pm$ 0.675 <sup>b</sup>	388	170.08 $\pm$ 1.010 <sup>b</sup>
5–6	661	27.90 $\pm$ 0.120 <sup>c</sup>	592	118.45 $\pm$ 0.613 <sup>ab</sup>	507	172.86 $\pm$ 0.920 <sup>ab</sup>
7–8	647	28.40 $\pm$ 0.121 <sup>b</sup>	589	119.50 $\pm$ 0.609 <sup>a</sup>	506	174.16 $\pm$ 0.922 <sup>a</sup>
9–10	521	28.81 $\pm$ 0.129 <sup>ab</sup>	469	120.14 $\pm$ 0.657 <sup>a</sup>	380	174.36 $\pm$ 1.000 <sup>a</sup>
11–12	320	28.88 $\pm$ 0.158 <sup>ab</sup>	275	119.06 $\pm$ 0.809 <sup>ab</sup>	209	172.86 $\pm$ 1.240 <sup>ab</sup>
13 $\leq$	173	29.06 $\pm$ 0.211 <sup>a</sup>	155	120.92 $\pm$ 1.070 <sup>a</sup>	101	176.86 $\pm$ 1.720 <sup>a</sup>
<b>Sex of calves</b>		***		***		***
Male	1429	28.66 $\pm$ 0.102 <sup>a</sup>	1283	120.77 $\pm$ 0.514 <sup>a</sup>	1039	176.96 $\pm$ 0.817 <sup>a</sup>
Female	1392	27.99 $\pm$ 0.102 <sup>b</sup>	1253	117.50 $\pm$ 0.519 <sup>b</sup>	1052	170.10 $\pm$ 0.817 <sup>b</sup>

 BW<sub>0</sub>: Birth weight; BW<sub>6</sub>: Body weight at 6 months; BW<sub>12</sub>: Body weight at 12 months

 \*:  $p < 0.05$  \*\*:  $p < 0.01$  \*\*\*:  $p < 0.001$ 

 a, b, c, d, e: Differences between averages with different letters in the same column are significant ( $p < 0.05$ ).

In Egyptian buffaloes, the values were reported as  $42.0 \pm 0.5$  kg by Marai et al. [21] and  $33.263 \pm 0.828$  kg by Kamal El-den et al. [22]; in Nili Ravi buffaloes, the values were reported as  $31.2 \pm 0.84$  kg by Charlini and Sinniah [20] and as  $36.1 \pm 3.23$  kg by Kuthu and Hussain [23]; in Anatolian buffaloes, they were reported as 30.4 kg by Çelikeloğlu et al. [24]; in Swamp buffaloes, they were reported as  $30.11 \pm 4.49$  kg by Thevamanoharan et al. [25]; in Murrah buffaloes, they were  $32.4 \pm 0.30$  kg by Thiruvenkadan et

al. [26], and in Iraqi buffaloes, the values were reported as  $37.711 \pm 0.231$  kg by Al-Khauzai [27]. These variations in BW<sub>0</sub> may result from differences in management and a lack of genetic improvement. Although BW<sub>0</sub> is the first appropriate criterion for growth, maternal influences are strong and should not be ignored [22]. The low BW<sub>0</sub> value in the present study may be due to the breeds of buffaloes in other studies, their higher productivity, and the genotype differences in the regions where the studies are carried out.

**Table 2.** Least squares means ( $\pm$  SE) of various growth periods in Anatolian buffaloes according to the district, calving year, season, age, and sex.

Non-Genetic factors	ADG <sub>0-6</sub> (g)		ADG <sub>0-12</sub> (g)		ADG <sub>6-12</sub> (g)	
	n	Mean $\pm$ SEM	n	Mean $\pm$ SEM	n	Mean $\pm$ SEM
<b>Overall means</b>	2536	504.64 $\pm$ 2.31	2091	398.24 $\pm$ 1.88	2091	304.27 $\pm$ 2.45
District		***		***		***
Amasra	126	485.15 $\pm$ 5.40 <sup>b</sup>	97	380.11 $\pm$ 4.15 <sup>c</sup>	97	291.78 $\pm$ 5.42 <sup>c</sup>
Kurucaşile	124	536.54 $\pm$ 5.53 <sup>a</sup>	113	425.47 $\pm$ 3.89 <sup>a</sup>	113	326.29 $\pm$ 5.07 <sup>a</sup>
Centrum	2000	496.81 $\pm$ 1.55 <sup>b</sup>	1646	395.23 $\pm$ 1.34 <sup>b</sup>	1646	306.99 $\pm$ 1.75 <sup>b</sup>
Ulus	286	500.05 $\pm$ 3.72 <sup>b</sup>	235	392.13 $\pm$ 2.84 <sup>bc</sup>	235	292.04 $\pm$ 3.70 <sup>c</sup>
<b>Calving year</b>		***		***		***
2015	270	474.39 $\pm$ 4.29 <sup>d</sup>	243	368.00 $\pm$ 3.05 <sup>d</sup>	243	270.07 $\pm$ 3.98 <sup>b</sup>
2016	390	500.87 $\pm$ 3.64 <sup>c</sup>	373	385.77 $\pm$ 2.51 <sup>c</sup>	373	281.32 $\pm$ 3.27 <sup>b</sup>
2017	502	508.78 $\pm$ 3.34 <sup>bc</sup>	452	407.55 $\pm$ 2.36 <sup>ab</sup>	452	316.92 $\pm$ 3.09 <sup>a</sup>
2018	518	501.09 $\pm$ 3.23 <sup>c</sup>	469	402.68 $\pm$ 2.26 <sup>b</sup>	469	314.24 $\pm$ 2.95 <sup>a</sup>
2019	526	513.76 $\pm$ 3.17 <sup>b</sup>	502	411.02 $\pm$ 2.20 <sup>a</sup>	502	319.75 $\pm$ 2.87 <sup>a</sup>
2020	330	528.93 $\pm$ 3.84 <sup>a</sup>	52	414.40 $\pm$ 5.77 <sup>ab</sup>	52	323.34 $\pm$ 7.54 <sup>a</sup>
<b>Calving season</b>		***		**		***
Winter	377	498.28 $\pm$ 3.55 <sup>b</sup>	339	398.18 $\pm$ 2.56 <sup>ab</sup>	339	310.11 $\pm$ 3.34 <sup>a</sup>
Spring	592	500.78 $\pm$ 3.26 <sup>b</sup>	507	393.11 $\pm$ 2.47 <sup>b</sup>	507	295.76 $\pm$ 3.23 <sup>b</sup>
Summer	904	515.72 $\pm$ 2.70 <sup>a</sup>	733	402.00 $\pm$ 2.26 <sup>a</sup>	733	299.44 $\pm$ 2.95 <sup>b</sup>
Fall	663	503.77 $\pm$ 3.06 <sup>b</sup>	512	399.66 $\pm$ 2.47 <sup>a</sup>	512	311.78 $\pm$ 3.23 <sup>a</sup>
<b>Calving age (year)</b>		NS		*		NS
3-4	456	499.07 $\pm$ 3.39	388	392.50 $\pm$ 2.55 <sup>b</sup>	388	300.22 $\pm$ 3.32
5-6	592	502.90 $\pm$ 3.08	507	397.35 $\pm$ 2.32 <sup>ab</sup>	507	304.70 $\pm$ 3.03
7-8	589	506.53 $\pm$ 3.06	506	399.87 $\pm$ 2.33 <sup>ab</sup>	506	304.11 $\pm$ 3.04
9-10	469	507.76 $\pm$ 3.30	380	399.33 $\pm$ 2.53 <sup>ab</sup>	380	302.84 $\pm$ 3.30
11-12	275	501.29 $\pm$ 4.06	209	395.19 $\pm$ 3.13 <sup>ab</sup>	209	303.69 $\pm$ 4.09
13 $\leq$	155	510.27 $\pm$ 5.36	101	405.17 $\pm$ 4.33 <sup>a</sup>	101	310.08 $\pm$ 5.66
<b>Sex of calves</b>		***		***		***
Male	1283	511.85 $\pm$ 2.58 <sup>a</sup>	1039	406.62 $\pm$ 2.06 <sup>a</sup>	1039	312.97 $\pm$ 2.69 <sup>a</sup>
Female	1253	497.43 $\pm$ 2.61 <sup>b</sup>	1052	389.85 $\pm$ 2.06 <sup>b</sup>	1052	295.58 $\pm$ 2.69 <sup>b</sup>

ADG<sub>0-6</sub>: Daily weight gain from birth to 6 months; ADG<sub>0-12</sub>: Daily weight gain from birth to 12 months; ADG<sub>6-12</sub>: Daily weight gain from 6 to 12 months.

\*:  $p < 0.05$  \*\*:  $p < 0.01$  \*\*\*:  $p < 0.001$  NS: not significant ( $p > 0.05$ )

a, b, c, d: Differences between averages with different letters in the same column are significant ( $p < 0.05$ ).

The  $BW_6$  value found in this study ( $119.13 \pm 0.459$  kg) is compatible with the study by Çelikeloğlu et al. [24] ( $118.46$  kg) on Anatolian buffaloes. The current study finding is lower than the values reported by Marai et al. [21] ( $134.8 \pm 0.4$  kg) on Egyptian buffaloes, Al-Khauzai [27] ( $126.095 \pm 0.393$  kg) on Iraqi buffaloes, and Shahjahan et al. [15] ( $144.14 \pm 4.10$  kg) in F1 crossbred (Indigenous  $\times$  Mediterranean) buffaloes. On the other

hand, it is higher than the values of Thiruvankadan et al. [26] ( $87.9 \pm 0.95$  kg) and Shahjahan et al. [15] ( $113.42 \pm 4.47$  kg) in indigenous buffaloes in Bangladesh. The  $BW_{12}$  value ( $173.53 \pm 0.743$  kg) found is higher than the values found in Murrah buffalo researches by Thiruvankadan et al. [26] ( $134.2 \pm 1.41$ ). However, it is lower than the values reported by Çelikeloğlu et al. [24] in Anatolian buffaloes ( $179.37$  kg), Kamal El-den et al. [22] ( $208,470 \pm 5,042$

kg) and Shahjahan et al. [15] in indigenous buffaloes in Bangladesh ( $194.88 \pm 9.55$  kg), and in F1 crossbred (Indigenous  $\times$  Mediterranean) buffaloes ( $219.70 \pm 6.36$  kg). These differences in  $BW_6$  and  $BW_{12}$  values may have resulted from the differences in genotypes of buffaloes in the study areas and the differences in care, feeding, and management practices in the farms.

In this study,  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  were significantly ( $p < 0.001$ ) affected by the district. Similar to this study, Ergüneş et al. [16] reported that the province has a significant effect on  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  in Anatolian buffaloes. On the other hand, contrary to this study, Çelikeloglu et al. [24] notified that the province did not have a significant effect on  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  in Anatolian buffaloes. In this study, the highest  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  values were obtained in Kurucaşile district, and the lowest  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  values were obtained in the Amasra district. These different body weight values in the region may have resulted from the breeding bulls and the different management practices of the breeders. In the present study, the effect of calving year on  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  was found to be significant ( $p < 0.001$ ) (Table 1). Similar to the current study, many researchers [16, 22–25] reported that the calving year had a significant effect on  $BW_0$ . Moreover, the effect of calving year on  $BW_6$  was significant in Murrah buffaloes [26]. In addition, significant effects of calving year on  $BW_6$  and  $BW_{12}$  were reported in F1 cross (Indigenous  $\times$  Mediterranean) buffaloes from Bangladesh [15]. On the other hand, unlike this study, it was found that the calving year did not have a significant effect on  $BW_0$  in Egyptian buffaloes [21], on  $BW_6$  and  $BW_{12}$  in Anatolian buffaloes [24], and in indigenous buffaloes in Bangladesh [15], and on  $BW_{12}$  in Murrah buffaloes [26]. In this study, there was a generally smooth increase in BW values as the years progressed, and this can be attributed to the experience gained by the breeders in care and feeding over the years.

In this study, calving season was found to be significant on  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  ( $p \leq 0.05$ ). Similar to this study, it was reported by many researchers [19, 21] that the calving season had a significant effect on  $BW_0$ , on  $BW_6$  in Anatolian buffaloes [24], and on  $BW_6$  and  $BW_{12}$  again in Anatolian buffaloes [28]. On the other hand, unlike this study, some researchers [22–24] stated that calving season did not have a significant effect on  $BW_0$ , on  $BW_6$  in Murrah buffaloes [21,26], and on  $BW_6$  and  $BW_{12}$  in Bangladesh buffaloes [15]. In this study, the lowest  $BW_0$  value was obtained in those born in spring ( $28.13 \pm 0.129$  kg), and the highest  $BW_0$  value was in those born in autumn ( $28.50 \pm 0.118$  kg). This result shows that buffaloes make good use of pasture in the spring and summer seasons during the last stages of their pregnancy and contribute to the increase in BW of their offspring. In addition, in this study, while  $BW_6$  and

$BW_{12}$  values were close to each other in buffalo calves born in autumn, winter, and spring, the highest  $BW_6$  and  $BW_{12}$  values were obtained in buffalo calves born in summer.

In this study, the effect of calving age on  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  was found to be significant ( $p < 0.001$ ) (Table 1). Consistent with this study, in studies conducted by some researchers [18,19] on Anatolian buffaloes, the calving age was significant on  $BW_0$ ; in some studies on Iraqi buffaloes [27], it was significant on  $BW_0$  and  $BW_6$ , and in some other studies on Anatolian buffaloes [29], it had a significant effect on  $BW_{12}$ . However, contrary to the current research, it was reported that the effect of calving age on  $BW_0$  and  $BW_6$  in Anatolian buffaloes [29] and the effect of calving age on  $BW_6$  and  $BW_{12}$  in Anatolian buffaloes [28] were not significant. In this study, the lowest  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  values were reached in those born from buffaloes  $\leq 4$  years of age at  $26.92 \pm 0.134$ ,  $116.74 \pm 0.675$ , and  $170.08 \pm 1.010$  kg, respectively; and the highest  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  values were  $29.06 \pm 0.211$ ,  $120.92 \pm 1.070$ , and  $176.86 \pm 1.720$  kg, respectively, in those born from buffaloes aged  $13 \leq$  in the present study, it was observed that the  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  values of buffalo calves increased regularly as the calving age increased. This result shows that reproductive performance increases with increasing age in buffaloes and contributes to the increase in  $BW_0$ ,  $BW_6$ , and  $BW_{12}$ . In this study, the effect of sex on  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  was found to be significant ( $p < 0.001$ ) (Table 1). Similar to this study, Kul et al. [29] reported that sex had a significant effect on  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  in Anatolian buffaloes, on  $BW_6$  in Iraqi buffaloes [27], and on  $BW_{12}$  in Egyptian buffaloes [22]. On the other hand, unlike this study, it was reported that sex did not have a significant effect on  $BW_6$  and  $BW_{12}$  in Anatolian buffaloes [24] and Bangladesh buffaloes [15], and on  $BW_0$  in Egyptian buffaloes [22]. It was also calculated in the present study that males have significantly higher  $BW_0$ ,  $BW_6$ , and  $BW_{12}$  than females.

The  $ADG_{0-6}$  value ( $504.64 \pm 2.31$  g) in this study was lower than in the studies by Marai et al. [21] ( $0.667.24 \pm 0.01$  kg) on Egyptian buffaloes, and in the studies by Shahjahan et al. [15] ( $599.66 \pm 22.21$  g) on F1 crossbred (Indigenous  $\times$  Mediterranean) buffaloes in Bangladesh. However, it was found to be higher than the value in the study on indigenous buffaloes in Bangladesh by Shahjahan et al. [15] ( $468.94 \pm 21.70$ ). It is thought that these different values may result from the different genotypes in the researched regions and the different management, care, and feeding regimes in the farms. The  $ADG_{0-12}$  ( $398.24 \pm 1.88$  g) value in the present study was found to be lower than in the studies on Egyptian buffaloes by Kamal Elden et al. [22] ( $0.479 \pm 0.013$  g), and than in the studies by Shahjahan et al. [15] on indigenous ( $456.85 \pm 25.69$  g) and F1 crossbred (Indigenous  $\times$  Mediterranean) buffaloes ( $506.09 \pm 16.82$  g) in Bangladesh ( $456.85 \pm 25.69$  g), and

this result can be attributed to the breeds and higher productivity of buffaloes in other studies. In the current study,  $ADG_{6-12}$  values were found to be  $304.27 \pm 2.45$  g, and there is not a study in the literature on this feature in Anatolian buffaloes and other buffalo breeds.

In the study, the effect of the district on  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  was found to be significant ( $p < 0.001$ ) (Table 2). In this study, the highest  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  values were found in the Kurucaşile district as  $536.54 \pm 5.53$ ,  $425.47 \pm 3.89$ , and  $326.29 \pm 5.07$  g, respectively; and the lowest  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  values were found in Amasra district as  $485.15 \pm 5.40$ ,  $380.11 \pm 4.15$ , and  $291.78 \pm 5.42$  g, respectively. The difference in ADG values between districts may be the result of the farms and the different management practices of the breeders. In the present study, the effect of calving year on  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  was found to be significant ( $p < 0.001$ ) (Table 2). In this study, the lowest  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  values were obtained as  $474.39 \pm 4.29$ ,  $368.00 \pm 3.05$ , and  $270.07 \pm 3.98$  g, respectively, in 2015; and the highest  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  values were obtained in 2020 as  $528.93 \pm 3.84$ ,  $414.40 \pm 5.77$ , and  $323.34 \pm 7.54$  g, respectively. In general, it is seen that the ADG increases as the years' progress. This result may be due to the training given to the breeders each year and the improvement in management on the farm. Similar to the results of the present study, calving year had a significant effect on  $ADG_{0-6}$  and  $ADG_{0-12}$  in F1 crossbred (Indigenous  $\times$  Mediterranean) buffaloes [15], and on  $ADG_{0-12}$  in Egyptian buffaloes [22]. However, unlike in the present study, it was reported in the studies that the effect of calving year on  $ADG_{0-6}$  and  $ADG_{0-12}$  in Bangladesh indigenous buffaloes [15] and on  $ADG_{0-6}$  in Egyptian buffaloes [21] were not significant.

In this study, the effect of calving season on  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  was found to be significant ( $p \leq 0.01$ ). The highest  $ADG_{0-6}$  and  $ADG_{0-12}$  values were obtained in summer as  $515.72 \pm 2.70$  and  $402.00 \pm 2.26$  g, respectively, and the highest  $ADG_{6-12}$  value was obtained in autumn as  $311.78 \pm 3.23$  g. Similar to the current study, the effect of calving season on  $ADG_{0-6}$  was found to be significant in studies on Bangladesh indigenous buffalo [15] and Egyptian buffalo [21]. However, contrary to this study, it was reported that the effect of the calving season on  $ADG_{0-6}$  and  $ADG_{0-12}$  was not significant in studies conducted in F1 crossbred (Indigenous  $\times$  Mediterranean) buffaloes [15]. In addition, it was reported that the effect of the calving season on  $ADG_{0-12}$  was not significant in the study on Egyptian buffaloes [22]. These differences may have resulted from changes in management and climatic conditions at different times of the year or over a period of

time. The effect of calving age on  $ADG_{0-6}$  and  $ADG_{6-12}$  was not significant ( $p > 0.05$ ), while the effect on  $ADG_{0-12}$  was significant ( $p < 0.001$ ). The lowest  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  values were obtained as  $499.07 \pm 3.39$ ,  $392.50 \pm 2.55$ , and  $300.22 \pm 3.32$  g respectively in buffaloes calving  $\leq 4$  years of age; and the highest  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  values were obtained as  $510.27 \pm 5.36$ ,  $405.17 \pm 4.33$ , and  $310.08 \pm 5.66$  g respectively in buffaloes calving  $13 \leq$  years of age. In general, it is seen that ADG increases with increasing calving age. This result can be attributed to the fact that the reproductive performance increases and contributes to the increase in ADG with the advancing age of the buffalo, and to the increased experience of the breeders in farm management. In the present study, the effect of sex on  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  was found to be significant ( $p < 0.001$ ). In this study, the  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  values in males were  $511.85 \pm 2.58$ ,  $406.62 \pm 2.06$ , and  $312.97 \pm 2.69$  g, respectively; and  $ADG_{0-6}$ ,  $ADG_{0-12}$ , and  $ADG_{6-12}$  values in females were  $497.43 \pm 2.61$ ,  $389.85 \pm 2.06$ , and  $295.58 \pm 2.69$  g, respectively. Similar to the current research, the effect of sex on  $ADG_{0-12}$  was found to be significant in a study on Egyptian buffaloes [22]. On the other hand, contrary to this study, it was reported that the effect of sex on  $ADG_{0-6}$  and  $ADG_{0-12}$  was not significant in the study on Bangladesh indigenous and F1 crossbred (Indigenous  $\times$  Mediterranean) buffaloes [15]. In this study, it was seen that male buffalo calves have higher ADG than female buffalo calves. This increase is probably due to the increasing differences in the endocrine system between males and females [30].

In this study, nongenetic factors significantly affected the weight and growth performance of buffalo calves at all ages. Body weights and growth performance characteristics at different ages obtained in the present study, which are economically important, should be taken into account to better predicted parameters such as heritability in future genetic studies. In addition, it was concluded that the growth performance of buffaloes can increase if the environmental factors that are important are taken into account in the selection.

#### Conflict of interest

The authors declare that they have no conflict of interest.

#### Acknowledgments

We would like to thank the Ministry of Agriculture and Forestry, General Directorate of Agricultural Research and Policies (project number: TAGEM/74MANDA2013-01), Bartın Buffalo Breeders' Association and Project Technical Staff for their contribution to this study.

## References

1. Soysal Mİ, Tuna YT, Gürçan EK. An investigation on the water buffalo breeding in Danamandira village of Silivri district of Istanbul province of Turkey. *Journal of Tekirdag Agricultural Faculty* 2005; 2: 73-78.
2. Soysal Mİ. Anatolian water buffaloes husbandry in Turkey. *Buffalo Bulletin* 2013; 32 (Special Issue 1): 293-309.
3. Gargani M, Pariset L, Soysal Mİ, Özkan E, Valentini A. Genetic variation and relationships among Turkish water buffalo populations. *Animal Genetics* 2010; 41: 93-96.
4. Soysal Mİ. Native Animal Genetic Researches of Turkey. 1st ed. Tekirdağ, Turkey: Kırklareli Feed Industry; 2010.
5. Giordano P, Guarini G, Ferrari P, Biondi-Zoccai G, Schiavone B et al. Beneficial impact on cardiovascular risk profile of water buffalo meat consumption. *European Journal of Clinical Nutrition* 2010; 64: 1000-1006.
6. Giuffrida-Mendoza M, Arenas de Moreno L, HuertaLeidenz N, Uzcátegui-Bracho S, Valero-Leal K et al. Cholesterol and fatty acid composition of longissimus thoracis from water buffalo (*Bubalus bubalis*) and Brahman-influenced cattle raised under savannah conditions. *Meat Science* 2015; 106: 44-49.
7. Huerta-Leidenz N, Rodas-González A, Vidal A, Lopez-Nuñez J, Colina O. Carcass cutout 651 value and eating quality of longissimus muscle from serially harvested savannah-raised 652 Brahman-influenced cattle and water buffaloes in Venezuela. *Animal Production Science* 2016; 56 (12): 2093-2104.
8. Akçapınar H, Özbeyaz C. Fundamental animal breeding. 1st ed. Ankara, Turkey: Kariyer Press; 1999.
9. Akdağ F, Arslan S, Caynak A, Teke B. The relationships phenotype, genotype and some environmental factors with weight in Jersey calves. *African Journal Biotechnology* 2011; 10: 7308-7313.
10. Zaman G, Goswami RN, Aziz A. Factors affecting gestation period and birth weight in Swamp Buffaloes of Assam. *Indian Journal of Animal Health* 2007; 46 (1): 33-36.
11. Hossein-Zadeh N, Madad M, Shadparvar AA, Kianzad D. An observational analysis of secondary sex ratio, stillbirth and birth weight in Iranian Buffaloes (*Bubalus Bubalis*). *Journal of Agricultural Science and Technology* 2012; 14: 1477-1484.
12. Thevarnanoharan K, Vandepitte W, Mohiuddin G, Chantalakhana C. Environmental factors affecting various growth traits of Swam: P Buffalo calves. *Pakistan Journal of Agricultural Sciences* 2001; 38 (3-4): 5-10.
13. Massey ME, Benyshek LL. Genetic, phenotypic and environmental correlations among performance traits estimated from Limousine Field Data. *Journal of Animal Science* 1982; 54 (1): 46-50.
14. Tekerli M. Manda Yıldızı data records, account and project tracking program, v5.04; 2019.
15. Shahjahan M, Khatun A, Khatun S, Hoque MM, Hossain S et al. Study on growth traits at weaning and yearling stages of indigenous and F1 crossbred buffalo in Bangladesh. *Asian Journal of Medical and Biological Research* 2017; 3 (4): 499-503. doi: 10.3329/ajmbr.v3i4.35341
16. Ergüneş Berkin E, Küçükkepapçı M, Kaptan C. Effects of some environmental factors on the some growth and lactation characteristics of Anatolian water buffalo in Balıkesir, Düzce and Kütahya. *Black Sea Journal of Engineering and Science* 2020; 3 (4): 146-150. doi: 10.34248/bsengineering.782547
17. Minitab. Minitab statistical software version 18.1; 2017.
18. Uğurlu M, Kaya İ, Saray M. Effects of some environmental factors on calf birth weight and milk yield of Anatolian Water Buffalo (*Bubalus bubalis*). *Bulgarian Journal of Agricultural Science* 2016; 22 (6): 995-998.
19. Kul E, Filik G, Şahin A, Çayıroğlu H, Uğurlutepe E et al. Effects of some environmental factors on birth weight of Anatolian buffalo calves. *Turkish Journal of Agriculture - Food Science and Technology* 2018; 6 (4): 444-446.
20. Charlini BC, Sinniah J. Performance of Murrah, Surti, Nili-Ravi buffaloes and their crosses in the intermediate zone of Sri Lanka. *Livestock Research for Rural Development* 2015; 27 (3): 47.
21. Marai IFM, Daader AH, Soliman AM, El-Menshawey SMS. Non-genetic factors affecting growth and reproduction traits of buffaloes under dry management housing (in sub-tropical environment) in Egypt. *Livestock Research for Rural Development* 2009; 21 (3).
22. Kamal El-den MA, Mohammed KM, Saudi EM. Estimation of genetic and non-genetic factors influencing growth traits performance of Egyptian buffalo. *Journal of Animal and Poultry Production Mansoura Univ* 2020; 11 (10): 383-388. doi: 10.21608/jappmu.2020.123613
23. Kuthu ZH, Hussain A. Effects of some environmental sources of variation on birth weight in Nili-Ravi buffalo calves. *Buffalo Bulletin* 2020; 39 (1): 47-52.
24. Çelikeloğlu K, Kocak S, Erdoğan M, Bozkurt Z, Tekerli M. The investigation of viability and body measurements for water buffalo calves. *Turkish Journal of Veterinary and Animal Sciences* 2019; 43 (1): 60-67.
25. Thevamanoharan K, Vandepitte W, Mohiuddin G, Chantalakhana C. Restricted maximum likelihood animal model estimates of heritability for various growth traits and body measurements of swamp buffaloes. *Pakistan Journal of Agricultural Science* 2001; 38 (1-2): 19-22.
26. Thiruvankadan AK, Panneer SS, Rajendran R. Non-Genetic and genetic factors influencing growth performance in Murrah buffalos. *South African journal of Animal Science* 2009; 39 (5): 102-106.

27. Al-Khauzai ALD. Estimation of genetic parameters for weights at different ages in local Iraqi buffaloes. *Plant Archives* 2020; 20 (2): 1801-1804.
28. Kul E, Tepe Ö. The effects of dam's milk yield and milk components on calf birth weight and some growth traits in Anatolian buffaloes. In: *Proceedings of the 12th World Buffalo Congress*; İstanbul, Turkey; 2019. pp.10.
29. Kul E, Şahin A, Çayıroğlu H, Filik G, Uğurlutepe E et al. Non genetic factors affecting some growth traits in Anatolian buffaloes. In: *Proceedings of the 8th Balkan Conference on Animal Science Conference (Balnimalcon)*; Prizren, Kosova; 2017. pp.171.
30. Swenson MJ, Reece WO. 1990. *Duke's physiology of domestic animals* (11th edn.) Cornell University Press, Ithaka, New York; 1990.