

1-1-2001

Allele Frequencies of the HumF13B Str Locus in the Çukurova Region

LALE DÖNBAK

SERPİL SALAÇIN

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

DÖNBAK, LALE and SALAÇIN, SERPİL (2001) "Allele Frequencies of the HumF13B Str Locus in the Çukurova Region," *Turkish Journal of Medical Sciences*: Vol. 31: No. 5, Article 7. Available at: <https://journals.tubitak.gov.tr/medical/vol31/iss5/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Allele Frequencies of the HumF13B Str Locus in the Çukurova Region

Received: October 10, 2000

¹Department of Biology, Faculty of Science and Art, Kahramanmaraş Sütçü İmam University, Karacasu/Kahramanmaraş.

²Department of Forensic Medicine, Medical School, Dokuz Eylül University, İzmir - TURKEY

Abstract: In this study, the STR system HumF13B polymorphism and forensic efficiency values were investigated in a population sample from the Çukurova region. The F13B phenotypes were analysed using polyacrylamide gel electrophoresis (PAGE) following a polymerase chain reaction (PCR) and were visualised by silver staining. Allele frequencies were calculated by the gene counting method. Forensic efficiency values were estimated using the obtained gene frequencies of F13B.

In this population sample, a total of five alleles, 6, 7, 8, 9 and 10 of F13B, were

observed. The calculated frequencies of these alleles were as follows: 6=0.0910; 7=0.0045; 8=0.3273; 9=0.2045; 10=0.3727. In the Çukurova region, the forensic efficiency values H-obs, MEC, pM and pD of F13B locus were estimated to be 0.69092, 0.44507, 0.14032 and 0.85968 respectively.

According to these values, the HumF13B STR locus may be useful in criminal and paternity cases in the Çukurova region.

Key Words: DNA, STRs, HumF13B, Allele frequencies, Turkey

Introduction

The forensic usefulness of STRs has been demonstrated in various studies (1-3). Before introducing STRs into forensic casework, allele frequency distributions in the relevant population must be known for statistical evaluation of the evidence. However, our knowledge about the allelic distributions of STRs routinely used in forensic applications is limited in Turkey to a few studies. This paper presents the allele frequency data of the HumF13B STR locus (4) in a Turkish population sample.

Materials and Methods

DNA was extracted from bloodstains (200 µl blood, air-dried on filter paper) from 110 healthy unrelated individuals residing in the Çukurova region of Turkey, using the Chelex-100 and Proteinase K method (5). The reaction assay amplifications were carried out according to the commercially available Gene Print STR system-F13B (Promega USA). The PCR products were separated in 6% non-denaturing polyacrylamide gels with the locus specific allelic ladder and visualised by silver staining (6). F13B alleles were typed by comparison with the allelic ladder components.

Allele frequencies were calculated by the gene counting method. Possible divergence from the Hardy-Weinberg equilibrium was determined using the conventional χ^2 -test. From the forensic efficiency values, the expected heterozygosity rate (H-exp)(7), the mean exclusion change (MEC)(8), the probability of match (pM) and the power of discrimination (pD) (9) were calculated. The allele frequency comparisons of different populations were performed by RxN contingency tables calculating χ^2 .

Results

In this population sample, 5 alleles and 10 F13B genotypes (3 homozygotes and 7 heterozygotes) were determined. Microvariant alleles were not observed. The calculated allele frequencies of F13B are shown in Table 1, together with frequencies in the other Asian and European populations. As shown in Table 1, the most common alleles were 10 ($f=0.3727$), 8 ($f=0.3273$) and 9 ($f=0.2045$). The results were in good agreement with the Hardy-Weinberg equilibrium ($\chi^2=0.9593$, $df=7$, $p=0.9955$). The forensic efficiency values determined for this locus are listed in Table 2.

Table 1. Allele frequencies of F13B in the Turkish population and in other European and Asian populations.

Allele	Allele frequency							
	Turks* n:110	Turks ¹⁰ n:200	Italians ¹¹ n:200	Hungarians ¹² n:223	Austrians ¹³ n:216	Egyptians ¹⁴ n:100	Yemenians ¹⁴ n:100	Japanese ¹⁵ n:367
6	0.0910	0.0750	0.1000	0.0920	0.0900	0.1700	0.1450	-
7	0.0045	0.0300	0.0175	0.0040	0.0160	0.0400	0.0500	0.0030
8	0.3273	0.3150	0.2625	0.2780	0.2450	0.3050	0.2200	0.0650
9	0.2045	0.2450	0.2450	0.2170	0.2250	0.2250	0.3100	0.2030
10	0.3727	0.3350	0.3725	0.4010	0.4190	0.2600	0.2750	0.7250
11	-	0.0030	0.0025	0.0070	0.0050	-	-	0.0040

* The results of this study

Table 2. The forensic efficiency values of HumF13B in the Çukurova region.

Statistical parametres	F13B
H-obs	0.69092
H-exp	0.70386
MEC	0.44507
pM	0.14032
pD	0.85968

Discussion

STR typing of medicolegal evidence in forensic applications requires knowledge of the suspect's ethnic group and the allele frequencies in that population.

Meyer et al. (16) investigated the allele frequencies distribution of 6 STR systems (THO1, VWA, ACTBP2, FES, F13B, D21S11) in 7 human populations (Germans, Turks, Moroccans, Japanese, Chinese, Papuans and Ovambos). Depending on the statistical data, they stated that significant allele frequency differences existed between major ethnic groups. They also showed that the most discriminative STRs between major ethnic groups are THO1, F13B and FES.

The allele frequency distribution of F13B in this Turkish population residing in the southern part of

Turkey are compared with that of some European and Asian populations in Table 1. Depending on interpopulation comparisons, the allele frequency distribution in this Turkish population was found to be similar to that in an Italian (Tur.-Ita.: $\chi^2=5.61$, $df=5$, $p=0.3450$), a Hungarian (Tur.-Hun.: $\chi^2=5.61$, $df=5$, $p=0.6821$), and an Austrian population (Tur.-Aus.: $\chi^2=7.26$, $df=5$, $p=0.2022$). Furthermore, the frequency profile was similar to that of an Adana area population ($\chi^2=6.92$, $df=5$, $p=0.2265$). However, the statistical data revealed significant differences from those in an Egyptian (Tur.-Egy.: $\chi^2=15.78$, $df=4$, $p=0.0033$), a Yemenian (Tur.-Yem.: $\chi^2=22.90$, $df=4$, $p<0.0001$) and a Japanese population (Tur.-Jap.: $\chi^2=192.69$, $df=5$, $p<0.00001$) mainly due to alleles 8 and 10 (10-15).

The obtained forensic efficiency values demonstrated that the HumF13B STR locus may be used in personal identification and paternity cases in the Çukurova region.

Correspondence author:

Lale DÖNBAK

Kahramanmaraş Sütçü İmam University,

Science and Art Faculty, Department of Biology,

Karacasu/Kahramanmaraş, Turkey

References

1. Brinkmann B. The Use of STRs in Stain Analysis. Proceedings from the Third International Symposium on Human Identification. Promega Corporation, Madison, USA 1992, pp:357-373.
2. Hammond HA, Jin L, Zhong Y, Caskey CT, Chakraborty R. Evaluation of 13 short tandem repeat loci for use in personal identification applications. Am J Hum Genet 55:175-189, 1994.
3. Alford RL, Hammond HA, Coto I, Caskey CT. Rapid and efficient resolution of parentage by amplification of short tandem repeats. Am J Hum Genet 55:190-195, 1994.

4. Nishimura DY, Murray JC. A tetranucleotide repeat for the F13B locus. *Nucleic Acids Res* 20:1167, 1992.
5. Wiegand P, Bajanowsky T, Brinkmann B. DNA typing of debris from fingernails. *Int J Leg Med* 106:81-83, 1993.
6. Budowle B, Chakraborty R, Guisti AM, Eisenberg AJ, Allen RC. Analysis of the variable number of tandem repeat locus D1S80 by the polymerase chain reaction followed by high resolution polyacrylamide gel electrophoresis. *Am J Hum Genet* 48:137-144, 1991.
7. Weaver RF, Hedrick PW. *Genetics*. W.M.C. Brown Publisher, Iowa 1989, pp:499-503.
8. Krüger J, Fuhrman W, Lichte KH, Steffens C. Zur verwendung des polymorphisms der sauren erythrocytenphosphotase bei der vaterschaftsbegutachtung. *Dtsch Z Gerichtl Med* 64:127-146, 1968.
9. Sensabaugh GF. Biochemical markers of individuality. *Forensic Science Handbook* (Eds. R. Saferstein) Simon and Schuster Company, New Jersey 1982, pp:358-373.
10. Alper B, Meyer E, Schürenkamp M, Brinkmann B. HumFES/FPS and HumF13B: Turkish and German population data. *Int J Leg Med* 108: 93-95, 1995.
11. Garofano L, Lago G, Vecchio C, Pizzamiglio M, Zanon C, Virgili A, Albonici L, Manzari V, Budowle B. Italian population data on the polymarker system and on the five short tandem repeat loci CSF1PO, TPOX, THO1, F13B and vWA. *J Forensic Sci* 43: 837-840, 1998.
12. Füredi S, Woller J, Padar Z. A population study of the STR loci HumLPL, HumF13B and HumF13A01 in Hungary. *Int J Leg Med* 110: 107-108, 1997.
13. Neuhuber F, Radacher M, Krasa B. F13B and CD4 allele frequencies in an Austrian population sample. *Int J Leg Med* 108: 227-228, 1996.
14. Klintschar M, Al-Hammadi N, Lux T, Reichenpfader B. Genetic variation at the short tandem repeat loci-HumvWA, HumFXIII B and HumFES/FPS in the Egyptian and Yemenian populations. *J Forensic Sci* 43: 850-853, 1998.
15. Nagai A, Yamada S, Watanabe Y, Bunai Y, Ohya I. Analysis of the STR loci HumF13A01, HumFXIII B, HumLIPOL, HumTHO1, HumTPOX and HumVWA31 in a Japanese population. *Int J Leg Med* 109: 34-36, 1996.
16. Meyer E, Wiegand P, Brinkmann B. Phenotype differences of STRs in 7 human populations. *Int J Leg Med* 107: 314-322, 1995.