

1-1-2001

Expression of the 8 kDa Heat Shock Protein (Ubiquitin) in Psoriasis

İBRAHİM PİRİM

TEOMAN ERDEM

NECMETTİN AKDENİZ

YAŞAR NURİ ŞAHİN

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

PİRİM, İBRAHİM; ERDEM, TEOMAN; AKDENİZ, NECMETTİN; and ŞAHİN, YAŞAR NURİ (2001) "Expression of the 8 kDa Heat Shock Protein (Ubiquitin) in Psoriasis," *Turkish Journal of Medical Sciences*: Vol. 31: No. 1, Article 11. Available at: <https://journals.tubitak.gov.tr/medical/vol31/iss1/11>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

İbrahim PİRİM¹
Teoman ERDEM²
Necmettin AKDENİZ²
Yaşar Nuri ŞAHİN¹

Expression of the 8 kDa Heat Shock Protein (Ubiquitin) in Psoriasis*

Received: December 1, 1999

Abstract: The 8kDa protein (ubiquitin) is a member of the small heat shock protein (hsb) family. Ubiquitin may additionally be involved in the pathogenesis of various diseases. Intracellular accumulation of ubiquitin has been detected in patients with neurodegenerative disease, brain ischemia, cancer, rheumatoid arthritis and haemodialysis.

Sixteen biopsy samples from patients with psoriasis who were treated by PUVA and the same samples from these patients before PUVA treatment were investigated by immunohistochemistry on formalin-fixed paraffin- embedded tissue sections, using anti-ubiquitin antibody.

In psoriatic epidermis, ubiquitin was mostly expressed in basal layers with a cytoplasmic staining pattern. Various cells of the dermis layer with psoriasis were stained with anti-ubiquitin antibody. Staining was again confined to the cytoplasmic region and lesions with or without PUVA treatment did not exhibit a different staining pattern. It did, however, seem to have higher staining intensity in PUVA-treated psoriatic lesions than untreated ones. We conclude that ubiquitin might be regarded as a useful marker of cell injury during the stressful conditions of tissue and as an indication of cell vitality and viability.

Key Words: Ubiquitin, psoriasis, immunohistochemistry.

Departments of ¹Biochemistry, ²Dermatology, Faculty of Medicine, Atatürk University, 25240, Erzurum-TURKEY

Introduction

All cells show a heat shock or stress response when exposed to heat or other influences such as toxic substances, heavy metals, treatment with ethanol, amino acid analogues etc. The response induces the specific proteins called heat-shock proteins (1,2). Human skin is at the front line with respect to insults from the environment and it is therefore of interest to understand its stress response. In addition, heat and cold treatment is used in a number of medical treatments for skin diseases. It is therefore of interest to know what the heat-shock response is and whether it contributes to or is deleterious to the medical treatment. Moreover, it is important to know whether the disease condition stimulates the heat shock itself.

Ubiquitin (8 kDa) is a 76 amino acid protein and is perhaps the most conserved gene product in evolution (3). It modulates degradation of abnormal or damaged proteins and belongs to the class of heat-shock proteins induced in conditions of cell stress (4,5). Many studies show that ubiquitin is involved in several chronic degenerative diseases characterised by the formation of cellular inclusion bodies. The ubiquitin response to cell injury appears to be cytoprotective and is particularly important in diseases of the nervous system (6-8). The present study was performed to investigate the pattern of ubiquitin expression in human epidermis and dermis with psoriasis, and to correlate ubiquitin expression with proliferative activity and degree of differentiation of cells. In addition it is known that psoriasis, a hyperkeratotic skin condition the aetiology of which is unknown, may be

* This article was presented in III. International Congress of Pathophysiology (28 June-3 July, 1998, Lahti, Finland)

treated by PUVA (psoralen plus UV light, A band). This treatment may also easily induce hsb production. Formallin-fixed, paraffin-embedded tissue samples from skin with psoriasis which were treated with PUVA, and untreated samples were investigated by immunohistochemistry, using a polyclonal antibody specific for ubiquitin (8 kDa hsb).

Materials and Methods

The study was conducted on 16 patients with psoriasis who were exposed to PUVA therapy for almost 10 or 15 sessions (total mean dosage of PUVA 22-38 joule/cm²). Biopsy samples for each patient was taken afterwards for immunohistochemical analysis.

Tissues were fixed in formalin and embedded in paraffin. Paraffin-embedded blocks were cut at a thickness of 5 mm on a vibrotome. Sections were dried in an incubator overnight at 40°C and deparaffinised in xylol and dehydrated through graded alcohol. Sections were later left in phosphate buffered saline (PBS). Sections were incubated with anti-ubiquitin antibody (1:250 dilution in PBS) for 4 hours at room temperature. Subsequent incubations were performed with peroxidase conjugated rabbit anti-mouse immunoglobulin antibody (Dakopotts, Copenhagen, Denmark) (1:500 dilution in PBS). The final colour product was developed by incubating for 10 min in diaminobenzidine alone (DAB; 0.5 ng/ml in PBS buffer containing 0.01 H₂O₂) for a brown reaction product. Counterstaining was performed with Harris's haematoxylin (Merck). Sections were coverslipped and examined under a light microscope.

Results

Keratinocytes of the basal layer, spinosum layer and granulosum layer were positive with anti-ubiquitin antibody (Fig. 1). There was, however, no reaction on the corneum layer with the antibody (Fig. 2). Staining intensity appeared not to vary from the upper layers of epidermis to the dermis.

Staining or labelling was heterogeneously distributed throughout the cells, corresponding to a cytoplasmic expression of hsb. All ubiquitin staining was confined to the cytoplasmic site of cells (Fig. 3).




Figure 1. Immunological reactivity of ubiquitin in epidermis and dermis layers of skin from patients with psoriasis. Stained with anti-ubiquitin antibody. The structures were visualised by the indirect peroxidase method. Magnification (x100).

In spite of the fact that ubiquitin is found in all eukaryotic cells and is ubiquitous, the immunohistochemical staining of it within the cells can only take place as its concentration is elevated. Control normal human skin sections (5 volunteers) were almost free of staining with anti-ubiquitin antibody (unstained normal skin sections are not shown).

It was interesting to discover that biopsy sections stained with anti-ubiquitin antibody did not display a distinct staining pattern between psoriatic lesions and PUVA-treated psoriatic lesions, in spite of having stronger antibody staining with PUVA-treated sections than not untreated ones. Furthermore, biopsies taken from the patients covered non-lesional regions where there was no significant ubiquitin staining after PUVA treatment (data not shown).




Figure 2. Ubiquitin reactive and non-reactive layers of skin with psoriasis. Arrow indicates corneum layer. The structures were visualised by the indirect peroxidase method. Magnification (x200).




Figure 3. Immunological reactivity of ubiquitin in skin sections with psoriasis. Anti-ubiquitin staining is confined to cytoplasm of cells. The structures were visualised by the indirect peroxidase method. Magnification (x200).

Discussion

In this study, we demonstrated by immunohistochemistry that ubiquitin is expressed in human skin with psoriasis and might be expressed with PUVA treatment as well. Most of the basal layer of epidermal cells displayed higher levels of ubiquitin than cells of the dermis. This could be an indication of stress effects on the first lines. The psoriatic cases we used in our study had been treated by PUVA. PUVA has the potential to induce stress in epidermal cells since the skin is exposed to increasing temperature during the time in the PUVA cabinet. When the stress response is triggered, the synthesis of most of the cellular proteins is switched off temporarily; consequently cell proliferation would be inhibited at least temporarily and it is known that PUVA treatment is basically antiproliferative. These effects may be more important if local PUVA becomes more widely accepted. The biopsies in the present work were taken from the skin of patients with psoriasis, before and after treatment with PUVA.

It is worth noting that the non-psoriatic regions of the biopsies with PUVA did not show anti-ubiquitin antibody staining. However, the staining of the sections with PUVA treatment appeared to be much denser than untreated biopsy sections. This difference remains unsolved, whether PUVA increases the ubiquitin activities in psoriatic lesions where the cells are highly susceptible to the disease and the cells have already had elevated ubiquitin activity or non-psoriatic regions with PUVA response to synthesis of ubiquitin, but it is not as detectable as the disease stimulation.

Ubiquitin is a critical factor in the ATP-dependent pathway for protein degradation in eukaryotic cells (9). The discovery that ubiquitin is a hsp suggests an important role for protein breakdown in the protection of cells from high temperatures and other types of stress, for instance inflammatory effect and radiation effect.

Recent studies demonstrated that other hsp were induced following PUVA treatment (10,11) or other disease conditions (12-14). Changes in the expression of hsp often occur under conditions that are ultimately lethal to the cells, indicating protein damage and denaturation as the common mechanism of indication (15).

There is concern that long-term PUVA may generate activated oxygen species, such as the superoxide radical anion (O_2^-), the hydroxyl radical (OH), or hydrogen

peroxide, in the epidermal cells, including DNA strand breaks and peroxidation of protein. Most protein oxidation is due to the oxidation of sulphhydryl groups causing disulphide crosslink changes, which inactivate or denature proteins (16,17).

In conclusion, using a polyclonal anti-ubiquitin antibody, we demonstrated the presence of the disease

and PUVA-induced ubiquitination. The ubiquitin produced may be a response which prevents cell degeneration.

Correspondence author

İbrahim PİRİM

*Medical School, Department of Biochemistry
Atatürk University, 25240 Erzurum-TURKEY*

References

1. Lidquist S. Heat shock proteins. *Ann. Rev. Biochem.* 55: 1151-91, 1986.
2. Morimoto RI, Tissieres A, Gregopoulos C. Stress proteins in biology and medicine. Cold Harbor Laboratory Press, 1990.
3. Schlesinger Dh, Golstein G, Niall HD. The complete amino acid sequence of ubiquitin. An adenylate cyclase stimulating polypeptide probably universal in living cells. *Biochemistry* 14: 2214-18, 1975.
4. Bond U, Schlesinger M. Ubiquitin is a heat shock protein in chicken embryo fibroblasts. *Mol. Cell. Biol.* 5: 945-53, 1985.
5. Finley D, Ozkaynak E, Varshovsky A. The yeast polyubiquitin gene is essential for resistance to high temperatures starvation and other stresses. *Cell.* 48: 1035-45, 1987.
6. Magnusson K, Wieoch K. Impairment of protein ubiquitination may cause delayed neural death. *Neuroscience letters.* 96: 264-70, 1989.
7. Mayer RJ, Lowe J, London M. Ubiquitin and the molecular pathology of chronic degenerative disease. *J. Pathol.* 163: 279-84, 1991.
8. Mayer RJ, Arnold J, Laszlo L, London M. Ubiquitin in health and disease. *Biochimica et Biophysica Acta.* 1089: 127-39, 1991.
9. Jentsch S, Seufert W, Hauser HP. Genetic analysis of the ubiquitin system. *Biochimica et Biophysica Acta.* 1089:127-39, 1991.
10. Musaud AS, Cunliffe WJ, Hollond DB. Are stress proteins induced during puva therapy. *British J Dermatology* 134:892-899, 1996.
11. Muramatsu T, Yamoshino Y, Toda H. 8-methoxypsoralen plus PUVA Induces the 72 kDa heat shock protein in organ-cultured normal human skin. *Photochem Photobiol.* 58:809-12, 1993.
12. Trautinger F, Mugge IK, Dekrout B, Knobler RM, Metz D. Expression of the 27 kDa hsp in human epidermis and in epidermal neoplasms. *British J Dermatology* 133: 199-202, 1995.
13. Pirim İ, Gündoğdu C, Sari I. Amyloid precursor protein Fragment (A4) and heat shock protein (8 kDa Hsp) Immunoreaktivitiy during ischemia. *Turkish J of Medical Science.* 24: 149-251, 1995.
14. Izaki S, Goto Y, Kaburagi Y. Antibody Production to hsp with Mr 65 kDa (Hsp 65) in cutaneous Inflammation: A possible relation to focal infection. *Acta Otolaryngol (stockh), Supple* 523: 197-200, 1996.
15. Lee KJ, Hohn GM. Abnormal proteins as the trigger for the induction of stress responses; heat, diamide and sodium arsenite. *J. Cell Physiol* 136: 411-20, 1988.
16. Carbonare MD, Pathak MA. Skin photosensitising agents and the role of reactive oxygen species in photoaging. *J Photochem Photobiol B: Biol.* 14: 105-124, 1992.
17. Paillous N, Vicendo P. Mechanisms of photosensitised DNA cleavage. *J Photochem Photobiol B: Biol.* 203-09, 1993.