

1-1-2001

Pulmonary Infections Due to Legionella pneumophila in Immunocompromised Patients

DEMET KAYA

SİNAN ERGİNEL

MUZAFFER METİNTAŞ

ZEKİ ÜSTÜNER

İRFAN UÇGUN

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>

 Part of the [Medical Sciences Commons](#)

Recommended Citation

KAYA, DEMET; ERGİNEL, SİNAN; METİNTAŞ, MUZAFFER; ÜSTÜNER, ZEKİ; UÇGUN, İRFAN; and ÖZDEMİR, NECLA (2001) "Pulmonary Infections Due to Legionella pneumophila in Immunocompromised Patients," *Turkish Journal of Medical Sciences*: Vol. 31: No. 1, Article 12. Available at: <https://journals.tubitak.gov.tr/medical/vol31/iss1/12>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Pulmonary Infections Due to *Legionella pneumophila* in Immunocompromised Patients

Authors

DEMET KAYA, SINAN ERGİNEL, MUZAFFER METİNTAŞ, ZEKİ ÜSTÜNER, İRFAN UÇGUN, and NECLA ÖZDEMİR

Demet KAYA¹
Sinan ERGİNEL²
Muzaffer METİN TAŞ²
Zeki ÜSTÜNER³
İrfan UÇGUN²
Necla ÖZDEMİR²

Pulmonary Infections Due to *Legionella pneumophila* in Immunocompromised Patients

Received: March 05, 1999

Abstract: The numbers of immunocompromised individuals increase as programmes for organ transplantation and chemotherapy of malignant disease advance. Pulmonary infections, which progress rapidly, are one of the major clinical problems in these groups. This study was designed to evaluate the microbiology of pneumonia in immunocompromised patients and to determine the role of *Legionella pneumophila*.

Immunocompromised patients receiving antineoplastic chemotherapy, diagnosed as pneumonia with clinical and radiographic findings and routine laboratory tests were included in the study. For microbiological diagnosis, expectorated sputum was cultured for *L. pneumophila*. In addition to standard bacterial, fungal and mycobacterial media, specimens were examined microscopically. Direct fluorescent antibody (DFA) assay was used for antigen detection of *L. pneumophila*. By indirect fluorescent antibody (IFA) assay the sera of the patients were screened for *L. pneumophila*.

Of 74 sputum samples, 7 (9.5%) specimens yielded *L. pneumophila*. The other isolates were *Escherichia coli* (n:3), *Enterobacter* sp. (n:2), *Aspergillus* sp. (n:3), *Mycobacterium tuberculosis* (n:3), *Staphylococcus aureus* (n:1), yeasts (n:10), *Nocardia* sp. (n:2) and flora members (n:43). Antibiotic supplemented media and acid treatment was found superior for the isolation of *L. pneumophila*. Direct microscopic evaluation was necessary to support the culture results. DFA assay had 85% sensitivity and 100% specificity. Of 3 sera screened by IFA for *L. pneumophila* antibodies, 2 had the titres 1/256, indicating the disease.

Our results showed that *L. pneumophila* has an important role among the respiratory pathogens in immunocompromised patients, and it may be useful if techniques for isolation and identification of this pathogen are added to routine culturing programmes.

Key Words: *L. pneumophila*, immunocompromised patients, pulmonary infection, diagnosis.

¹Department of Microbiology and Clinical Microbiology, Faculty of Medicine, Abant İzzet Baysal University, Düzce - TURKEY
Departments of ²Chest Disease, ³Haematology, Faculty of Medicine, Osmangazi University, Eskişehir - TURKEY

Introduction

During the past decade the number of immunocompromised patients has increased in parallel to the increased use of high-dose chemotherapy and immunosuppressive drugs. A compromised host who has one or more defects in the body's natural defence mechanisms, is predisposed to severe, often life-threatening infections (1,2).

A sensible approach to the diagnosis and treatment of infections in these patients is to begin with an appreciation of the predisposing factors, because each is associated with an array of pathogens. The isolation and identification of a specific pathogen permits appropriate antibiotic management of patients with pulmonary infections (1,2).

This study was designed to evaluate the microbiology of pneumonia in immunocompromised patients at the Osmangazi University Medical Faculty Hospital and to compare the results with those of previous reports. The role of *L. pneumophila* and microbiological techniques for isolation and identification of *L. pneumophila* were also discussed.

Materials and Methods

Immunocompromised patients with pneumonia admitted to the Department of Chest Disease and Haematology Clinics of the Medical Faculty of Osmangazi University over a 1- yr period from 1 November 1995 to 31 March 1997 were eligible for the study. Criteria for hospitalization were at the discretion of the clinic from

which the patient was admitted. Outpatient diagnostic and therapeutic interventions were performed. Inclusion criteria were as follows:

- 1) patients > 20 yr of age
- 2) pulmonary infiltrate on chest radiograph, and
- 3) clinical findings of one "major criterium": cough, sputum production of fever > 37.8°C; or two "minor criteria": pleuritic chest pain, dyspnea, altered mental status, pulmonary consolidation by physical examination or white blood cell count > 12.000/mm³. Patients were excluded if chest roentgenograms did not reveal that infiltrate or radiographic abnormalities were attributable to a non-infectious aetiology. Neutropenic patients receiving antineoplastic chemotherapy were included in the study. The patients were considered neutropenic if the granulocyte count was < 1.000/mm³.

Routine tests obtained upon admission included chest roentgenograms, complete blood count and serum electrolytes (3).

For microbiological diagnosis, expectorated sputum was obtained. Standard bacterial cultures were performed by inoculation of the specimen onto 5% sheep blood, chocolate, and Eosin Methylene Blue (EMB) (Difco) agars. Mycobacterial cultures (Bactec system and Löwenstein Jensen medium-Difco) and fungal cultures (Sabouraud Dextrose Agar-SDA, Difco) were also made.

The culture for *Legionella* sp. was made using Buffered Charcoal Yeast Extract Agar (BCYE) and BMPA containing cefamandol, polymyxin B and anisomycin (BCYE-selective agars, BCYE-s) (Oxoid). The culture results before and after acid treatment of sputum (HCl-KCl buffer, ph: 2.2) with BCYE and BCYE-selective agars were compared. All plates were incubated at 37°C for 24-28 h in an aerobic atmosphere. After examination and recording of the results, blood agar, EMB and SDA plates were discarded, whereas BCYE and BCYE-s agar plates were incubated for 5 days. All isolates were identified by standard procedures and then by the Sceptor system. Colonies suspected for *Legionella* were removed and stained with Gram's stain. When Gram-negative coccobacillary or filamentous forms resembling the members of the family Legionellaceae were observed, biochemical tests were performed, including catalase, oxidase, urease, nitrate, gelatinase and hippurate hydrolysis tests.

DFA for the antigen of *L. pneumophila* serogroup 1-14 was done on sputum (Fresenius Diagnostics). For *L. pneumophila* serogroup 1-8, serologic tests were done by IFA (Fresenius Diagnostics). For this purpose, a single set of sera was collected from 3 patients and screened for *L. pneumophila* serogroup 1-8 by IFA assay.

For microscopic evaluation, all sputum samples were stained with Gram's and Ehrlich Ziehl Neelsen (EZN) stains. Sputum smears with a paucity of flora members and a large number of neutrophils and small, pleomorphic faintly staining Gram-negative rods supported the diagnosis if growth was obtained by culture (3,4).

A total of 74 immunocompromised patients were included in this study. The distributions according to age, sex and primary pathology are listed in Table 1.

Table 1. Characteristics of patients.

Characteristics	No of patients (No: 74)
Mean age, yr (range)	57.3 (22-75)
Male / Female	39 / 9
Primary pathology	
CLL	7
NHL	9
ANLL	3
HD	5
ALL	1
Renal transplant	2
Bronchial carcinoma	44
Malignant mesothelioma	3

ANLL: Acute nonlymphocytic leukemia
 NHL: Non-Hodgkin lymphoma
 ALL: Acute lymphoblastic leukemia
 HD: Hodgkin's disease
 CLL: Chronic lymphocytic leukemia

If the patients were diagnosed with pneumonia due to *L. pneumophila*, they were treated with erythromycin or roxitromycin. Improvement of the patient was defined as clinical improvement with negative microbiological diagnostic test.

Results

Of 74 sputum specimens, 7 specimens yielded *L. pneumophila* by culturing and direct microscopic examination. The distribution of organisms isolated from sputum samples is listed in Table 2.

L. pneumophila was isolated from the sputum samples of 7 patients. The characteristics and outcomes of these patients are given in Table 3.

Table 2. The distribution of organisms isolated from sputum samples.

ISOLATE	No	%
<i>L. pneumophila</i>	7	9.5
<i>E. coli</i>	3	4.1
Enterobacter sp.	2	2.7
Aspergillus sp.	3	4.1
<i>M. tuberculosis</i>	3	4.1
<i>S. aureus</i>	1	1.3
Yeast	10	13.5
Nocardia sp.	2	2.7
Flora members	43	58.0

Of 7 sputum specimens including *L. pneumophila*, DFA assay yielded positive results for 6 specimens (85% sensitivity) and no cross reaction was present. Of 3 sera screened by IFA for *L. pneumophila* antibodies, the titres were 1/256 for 2 patients, indicating acute disease, and negative for one patient. DFA and antibody detection results supported the results obtained by culture and microscopic evaluation. The comparison of culturing results with DFA and serology results is shown in Table 4.

Table 3. The characteristics of the patients.

Patient No	Age (yr)	Sex	Diagnosis	Cytotoxic therapy and/or Corticosteroid	Symptoms	Neutropenia	Chest X-ray	Therapy	Outcome
1(SC)	62	M	Bronchial Cancer	Yes	Cough, malaise	No	Heterogenic density	Roxitromycin	Improved
2(IY)	65	M	Bronchial Cancer	Yes	Dyspnea, productive cough, fever, Malaise	No	Bilateral patchy infiltrate	Roxitromycin	Improved
3(EÇ)	51	M	Bronchial Cancer	Yes	Bloody sputum Fever, malaise	No	Infiltration and cavitaray lesion in left lower lobe	Roxitromycin	Improved
4(NY)	70	M	Bronchial Cancer	Yes	Dyspnea, cough, Chest pain, fever Malaise	No	Patchy density in right lower lobe, pleural effusion	(He refused therapy)	Improved
5(NB)	52	M	SLE	Yes	Cough	Yes	Bilateral patchy infiltrate	Eryhromycin	Improved
6(AP)	69	M	KLL	Yes	Productive cough, Fever	Yes	Bilateral patchy infiltrate	Eryhromycin	Improved
7(AY)	63	F	Lymphoma	Yes	Non-productive cough, fever	Yes	Unilateral patchy density	Eryhromycin	Improved

Patient No	CULTURE				DFA	SEROLOGY
	BCYE		BCYE-S			
	Acid treatment	No acid treatment	Acid treatment	No acid treatment		
1	Positive	Positive	Positive	Positive	Positive	NT
2	Positive	Positive	Positive	Positive	Positive	1/256
3	Positive	Positive	Positive	Positive	Negative	1/256
4	Positive	Positive	Positive	Positive	Positive	NT
5	Positive	Positive	Positive	Positive	Positive	NT
6	Negative	Negative	Positive	Negative	Positive	NT
7	Positive	Negative	Positive	Positive	Positive	Negative

Table 4. The results obtained by culturing, DFA and IFA assays.

NT: Not tested

The isolation of *L. pneumophila* was found to be difficult with BCYE agar. However, BCYE-s agar allowed us to see the typical colonies easily, and pure culture was easily obtainable from this medium. In our study, all *L. pneumophila* isolates could be isolated as a pure culture by using BCYE-s agar whereas the isolates were inhibited by the flora members and not distinguished by BCYE agar.

After acid treatment of sputum samples, the flora members were inhibited, and thus isolation was easy, as BCYE-s agar. When the sputum was acid treated and inoculated onto BCYE agar, growth was obtained in 6 samples, similar to inoculating onto BCYE-s agar without acid treatment. But all samples were found positive for *L. pneumophila* when both acid treatment and inoculating onto BCYE-s agar was done. Thus, both acid treatment and using selective agar were found to be more useful for the isolation of *L. pneumophila*.

Discussion

In 1976, an outbreak of pneumonia occurred at a hotel at the site of the American Legion Convention in Philadelphia. The discovery of *L. pneumophila* and Legionnaires' disease was after this outbreak. *L. pneumophila* can survive in a wide range of environmental conditions. The distribution of *L. pneumophila* on man-made aquatic reservoirs, including water distribution systems and cooling towers, are related to epidemiology. The incidence of Legionnaires' disease is dependent on the degree of contamination of the organism in the aquatic reservoir and the susceptibility of the persons, availability of specialized laboratory tests and their application to the infected patients (1,5-7).

Numerous studies have shown *L. pneumophila* among the three most common microbial aetiologies of community-acquired pneumonia. It is suggested by CDC investigations that only 3% of sporadic cases of Legionnaires' disease are correctly diagnosed. The incidence of nosocomial pneumonia by Legionella sp. is dependent on the degree of colonization, the number of immunosuppressed host and the availability of culture methods. Cigarette smoking, chronic lung disease, advanced age and immunosuppression have been implicated as risk factors. Excess alcohol intake, renal failure and surgery are major predisposing factors, with transplant recipients at highest risk (1,7).

The aim of this study was to determine the distribution of respiratory pathogens in immunocompromised patients and the role of *L. pneumophila*. All of the patients included in this study were receiving immunosuppressive therapy for the treatment of their primary pathologies.

In granulocytopenia and cellular immune dysfunction, Gram-negative bacterial infections due to the members of Enterobacteriaceae and *L. pneumophila* is common. In our study, *E. coli* was isolated from 3, Enterobacter sp. was isolated from 2 and *L. pneumophila* was isolated from 7 sputum samples. Granulocytopenic patients are also susceptible to infections caused by Gram-positive bacteria, including *S. aureus*, *S. epidermidis* and streptococci. Also, fungi such as candida sp. and Aspergillus sp. cause infections. In our study, Aspergillus sp. was isolated from 3 specimens, yeast from 10 specimens and *S. aureus* from 1 specimen. *M. tuberculosis* and Nocardia sp, which are also common in cellular immune dysfunction, were also isolated. This distribution of causative agents is similar to that in the literature (1-3, 8-10).

Microbiological diagnosis of nosocomial pneumonia is particularly difficult, and sputum cultures have very low sensitivity and specificity. When microorganisms such as *E. coli*, *S. aureus* or Candida sp. are isolated, they may be aetiologic agents or represent oropharyngeal colonization. Thus, the approach used in differentiating them is very important.

The incidence of *L. pneumophila* varies widely from study to study, ranging from 1 to 16%. Mortality rates range from 5% in nonimmunosuppressed patients to 80% in compromised patients. Pneumonia with *L. pneumophila* and other Legionella sp. may occur in patients with a variety of defects, and its diagnosis is particularly difficult because of its weak Gram staining and the delay associated with a serological response of difficult cultures. For culturing this organism, BCYE agar is one of the media used. BCYE agar can be used for the isolation of the organism, but as respiratory specimens, especially sputum samples, include flora members, their growth does not allow easy differentiation of *L. pneumophila* colonies. Therefore, a more selective medium is recommended. BCYE-selective agar including antimicrobials for the inhibition of flora members is more useful (3,4). In our study, the identification of *L. pneumophila* was much easier when BCYE-s agar was used.

Another approach for the isolation of *L. pneumophila* is acid decontamination. If the sputum samples are acid treated and then cultured, flora members are inhibited, so isolation may be easier. In our study the results of acid treatment were similar to those obtained by culturing onto BCYE-s agar. These two techniques can be recommended and if both methods are used, accurate results may be obtained (11-13).

L. pneumophila is a faintly staining organism, so microscopic examination of the sputum specimens is usually insufficient. Microscopic examination is useful only when used in combination with another identification technique.

There is increasing use of fluorescent antibodies for identification of *L. pneumophila* in clinical specimens, suggesting that these methods have good sensitivity. In our study, the DFA test was applied to the sputum samples, and in 6 specimens that yielded *L. pneumophila*, positive results were obtained (85% sensitivity). Cross reaction with other organisms was not seen (100% specificity).

For serological diagnosis of Legionnaires' disease, a four-fold increase in antibody titres is required. A single

high titre ($\geq 1/128$) may indicate acute disease if the prevalence is low in that population. In our study, by IFA assay the sera of 3 patients were screened for *L. pneumophila* antibodies; 1/256 titre was obtained for 2 patients, indicating acute disease. We can conclude that IFA is not sufficient for diagnosis; it must be performed in addition to another technique, especially culturing.

In conclusion, immunocompromised patients are susceptible to infections caused by a variety of infectious agents. *L. pneumophila* is one of the pathogens that cannot be isolated by routine diagnostic methods. Our study showed that it plays an important role in the pneumonia of immunocompromised patients. For microbiological diagnosis of pneumonia caused by *L. pneumophila*, culturing is the 'gold standard' especially when the specimen is acid decontaminated or inoculated onto antibiotic-supplemented media. Immunofluorescence techniques and direct microscopic examination are of diagnostic value when combined with culturing methods. Therefore, a combination of techniques must be added to the routine culturing programmes for the isolation and identification of *L. pneumophila*.

References

- Schimpff SC. Infections in the cancer patient- Diagnosis, prevention and treatment. Mandell, Douglas and Bennetts' Principles and Practice of Infectious Diseases (Eds. Mandell GL, Bennett JE, Dolin R.) 4th ed Vol 2 Churchill Livingstone, New York, 1995, pp: 2666-2675.
- Hopkin JM. Respiratory disease in the immunocompromised host. Non-AIDS. Respiratory Medicine (Eds. Brews RAL, Gibson GL, Gedde DM) Bailliere Tindall WB Saunders, 1990, pp: 961-73.
- Mundy LM, Auwaerter PG, Oldach D, Warner ML, Burton A, Vance E, Gaydos CA, Joseph JM, Gopalan R, Moore R., Quino TC, Charache P, Bartlett JG: Community-acquired pneumonia: Impact of Immune status, Am J Respir Crit Care Med 152: 1309-1315, 1995.
- Baron EJ, Peterson LR, Finegold SM. Bailey and Scott's Diagnostic Microbiology. 9th ed Mosby Co., USA, 1994, pp: 570-575.
- Edelstein P. Legionnaires' disease. Clin Infect Dis 16: 741-747, 1993.
- Yu VL. Legionella pneumophila (Legionnaires' Disease) Mandell, Douglas and Bennetts' Principles and Practice of Infectious Diseases (Eds. Mandell GL, Bennett JE, Dolin R.) 4th ed. Vol 2 Churchill Livingstone, New York, 1995, pp: 2087, 2097.
- Meyer RD, Edelstein PH, Kirby BD, Louie MH, Mulligan ME, Morgenstein AA, Finegold JM. Legionnaires' disease. Unusual clinical and laboratory features, Ann Intern Med 93: 240-243, 1980.
- Meunier F. Infections in patient with acute leukemia and lymphoma. Mandell, Douglas and Bennetts' Principles and Practice of Infectious Diseases (Eds. Mandell GL, Bennett JE, Dolin R) 4th ed. Vol 2 Churchill Livingstone, New York, 1995, pp: 2675-2686.
- Mandell LA. Community-acquired pneumonia. Etiology, epidemiology and treatment. Chest 108: 355-425, 1995.
- Woodhead MA, MacFarlane JT, Rodgers FG. Aetiology and outcome of severe community acquired pneumonia. J Infect Dis 10: 204-210, 1985.
- Kerr E, Snell H, Black BL, Storey M, Colby WD. Isolation of Nocardia asteroides from respiratory specimens by using selective buffered charcoal-yeast extract agar J Clin Microbiol 30(5): 1320-1322, 1992.
- Vickers RM, Rish JD, Yu VL. Clinical demonstration of isolation of Nocardia asteroides on buffered charcoal-yeast extract media. J Clin Microbiol 30(1): 227-228, 1992.
- Garrett MA, Holmes HT, Nolte FS. Selective buffered charcoal-yeast extract medium for isolation of Nocardia from mixed cultures. J Clin Microbiol 30(7): 1891-1892, 1992.