

1-1-2000

Acute Dynamic Exercise Reduces Intraocular Pressure

İBRAHİM KOÇER

ŞENOL DANE

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

KOÇER, İBRAHİM and DANE, ŞENOL (2000) "Acute Dynamic Exercise Reduces Intraocular Pressure," *Turkish Journal of Medical Sciences*: Vol. 30: No. 2, Article 20. Available at: <https://journals.tubitak.gov.tr/medical/vol30/iss2/20>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Ibrahim KOÇER¹
Şenol DANE²

Acute Dynamic Exercise Reduces Intraocular Pressure

Departments of¹ Ophtalmology, ²Physiology,
Faculty of Medicine, Atatürk University,
Erzurum- Turkey

Received: July 27, 1999

In recent years, several studies have shown that exercise reduces intraocular pressure (1-3). In this study, the effect of exercise on intraocular pressure was evaluated in normal healthy young subjects. Subjects were 17 male and 11 female high school students, 18-22 years of age (M= 19.5, SD= 1.67). Subjects who had ocular problems were excluded from the study.

The resting intraocular pressures of both eyes were assessed by aplanation tonometry. Then, subjects ran at 30% of their maximum rate for 6 minutes. Intraocular pressures were assessed after exercise.

Student's t test was used for statistical evaluation, with the Statgraphics computer program.

The mean intraocular pressures of the right and left eyes before exercise were 16.05±2.39 mmHg and 16.11±2.12 mmHg respectively. The mean intraocular pressures of the right and left eyes after exercise were 12.21±2.22 mmHg and 12.59±1.38 mmHg respectively. The differences in mean intraocular pressures between, before and after exercise for the right and left eyes were statistically significant (t= 11.84, p= 0.0; t= 9.64, p= 0.00).

The findings of the present study are consistent with the literature (1-3). As a consequence, acute dynamic exercise has been found to reduce intraocular pressure.

Quereshi has reported that physical fitness causes significant attenuation in the intraocular pressure (4). Quereshi et al. concluded that the intensity of exercise seems to be responsible for the magnitude of the decrease in the resting intraocular pressure after short-term exercise (5). Martin et al. claimed that acute dynamic exercise seem to change intraocular pressure through changes in plasma colloid osmotic pressure (6). Kiuchi et al. have reported that the amount of intraocular pressure reduction after short-term exercise seems to depend on the intensity of exercise, not on the duration or the quantity of exercise (7).

Passo et al. have reported that regular aerobic exercise is associated with a reduction in elevated intraocular pressure (8). Thus we conclude that acute dynamic exercise may represent an effective non-pharmacologic intervention for patients suspected of having glaucoma.

References

1. Quereshi IA. Effects of mild, moderate and severe exercise on intraocular pressure of sedentary subjects. *Ann Hum Biol*, 22 (6): 545-553, 1995.
2. Harris A, Malinovsky V, Martin B. Correlates of acute exercise-induced ocular hypotension. *Invest Ophtalmol Vis Sci*, 35 (11): 3852-3857, 1994.
3. Quereshi IA, Xi XR, Wu XD, Zharf J, Shiarhov F. The effect of physical fitness on intraocular pressure in chinese medical students. *Chung Hua I Hsuah Tsa Chih*, 58 (5): 317-322, 1996.

4. Quereshi IA. Effects of exercise on intraocular pressure in physically fit subjects. *Clin Exp Pharmacol Physiol*, 23 (8): 648-652, 1996.
5. Quereshi IA, Xi XR, Huang YB, Wu XD. Magnitude of decrease in intraocular pressure depend upon intensity of exercise. *Korean J Ophthalmol*, 10 (2): 109-115, 1996.
6. Martin B, Harris A, Hammel T, Malinovsky V. Mechanism of exercise-induced ocular hypotension. *Invest Ophthalmol Vis Sci*, 40 (5): 1011-1015, 1999.
7. Kiuchi Y, Mishima HK, Hotehama Y, Furumoto A, Hiroto A, Onari K. Exercise intensity determines the magnitude of intraocular pressure decrease after running. *Jpn J Ophthalmol*, 38 (2): 191-195, 1994.
8. Passo MS, Goldberg L, Elliot DL, Van Buskirk EM. Exercise training reduces intraocular pressure among subjects suspected of having glaucoma. *Arch Ophthalmol*, 109 (8): 1096-1098, 1991.