

1-1-1999

Flexion Contractures in a Diabetic Young Man(Rosenbloom Syndrome)

SERPİL SAVAŞ

YAVUZ BARDAK

ŞEVKET ERCAN TUNÇ

MELTEM ÇETİN

SELAMİİ AKKUŞ

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

SAVAŞ, SERPİL; BARDAK, YAVUZ; TUNÇ, ŞEVKET ERCAN; ÇETİN, MELTEM; and AKKUŞ, SELAMİİ (1999) "Flexion Contractures in a Diabetic Young Man(Rosenbloom Syndrome)," *Turkish Journal of Medical Sciences*: Vol. 29: No. 6, Article 23. Available at: <https://journals.tubitak.gov.tr/medical/vol29/iss6/23>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Serpil SAVAŞ¹
Yavuz BARDAK²
Şevket Ercan TUNÇ³
Meltem ÇETİN⁴
Selami AKKUŞ¹

Flexion Contractures in a Diabetic Young Man (Rosenbloom Syndrome)

Departments of ¹Physical Medicine and
Rehabilitatoin, ²Ophthalmology, ³Internall
Medicine, ⁴Radiology, Faculty of Medicine,
Süleyman Demirel University, Isparta-Turkey

Received: 03.07.1998

Key Words: Diabetes Mellitus, Rosenbloom
syndrome, limited joint mobility

Introduction

Limited joint mobility syndrome was initially described by Rosenbloom as painless limitation of the fingers and large joints in association with short stature, thick, tight, waxy skin, delayed sexual maturation, and early microvascular complications (1). Limited joint mobility (LJM) is thought to be a manifestation of the diffuse collagen abnormalities in diabetic patients (2). Limited joint mobility is of particular interest because it is common in young patients and associated with an increased risk for the microvascular complications of diabetes (2,3). A 21 year-old diabetic man who had multiple joint contractures and nonfamilial short stature with microvascular complications is presented.

Case Report

A 21-year-old man with insulin-dependent diabetes mellitus (IDDM) complained of stiffness and progressive limitation of the mobility of his joints for one year. He also complained of severe paresthesias and pain in his hands and feet. He had had IDDM for 17 years. He had never experienced arthritis and there was no family history of rheumatic disease. His mother had non-insulin-dependent diabetes mellitus (NIDDM).

His weight was 40 kg and his height was 146 cm. He had flexion contractures at the distal interphalangeal, proximal interphalangeal and metacarpophalangeal joints of his hands and elbows, and the metatarsophalangeal, proximal interphalangeal and distal interphalangeal joints of feet (Figure 1). His hands and feet were more severely affected. He was unable to use his hands properly and wear shoes. The range of motion in his wrists, knees, ankles and subtalar joints was severely limited. He was unable to oppose the palms of his hands (Prayer sign) (Figure 2). Lumbosacral movement was limited in all directions. Modified Schober test was 13 cm. There was no local tenderness over the sacroiliac joints. He had hypothenar and thenar atrophy. He was in Tanner stage III of puberty. Pubic and axillary hair showed insufficient development and he had no facial hair.

Laboratory investigations showed normal results of complete blood counts. The fasting glucose level was 220mg/dl. The HbA1c level was 10% (<7%). Urinalysis revealed glucosuria (150mgr/dl) and macroalbuminurea (450mg/24 hours). The erythrocyte sedimentation rate was 25mm/h. C-reactive protein, rheumatoid factor, antinuclear antibodies were not detected. Immunoglobulins and complements were at normal levels.



Figure 1. The patient with Rosenbloom Syndrome.

Electroneuromyography revealed severe sensorimotor polyneuropathy. Respiratory function tests were performed in order to establish the vital capacity. Forced vital capacity was 2 l (60% of the predicted value). Ocular examination showed neovascular glaucoma in the right eye and diabetic proliferative vitreoretinopathy in the left eye.

Skeletal age was 15 according to a wrist roentgenogram. A lumbosacral roentgenogram revealed a loss of lumbar lordosis. Facet joints and sacroiliac joints showed no abnormality. Feet roentgenograms revealed osteoporosis, tarsal abduction deformity, cavus deformity, severe Achilles tendon calcifications and large calcaneal spurs.

His pain was resolved with antiinflammatory medication. Range of motion exercises and stretching was performed for one month. There was no improvement in any joint range at the end of the program.

LJM has recently been recognized as part of the clinical spectrum of IDDM by Rosenbloom (1). LJM is seen not only in IDDM but also in NIDDM (4). The joints most frequently affected in LJM are the proximal interphalangeal joints of the fingers, the wrists, elbows, spine, knees, ankles and the toes (5). Thick, tight, and waxy skin, most prominent over the dorsum of the hand



Figure 2. The prayer sign.

and the forearm was described in diabetic patients leading to restriction of joint movements (1, 2). The exact pathogenesis of LJM is unclear. Evidence suggests that the diabetic hyperglycemic state leads to an increase in nonenzymatic glycosylation causing increased cross-linking of collagen (2). The most commonly suggested explanation for LJM has been that impaired degradation of collagen leads to its accumulation (6). The degree of the cross-linked collagen correlates with the presence of retinopathy and nephropathy (2).

Rosenbloom et al. suggested that alterations of periarticular connective tissue are related to changes occurring in the microvasculature (3). Our patient with LJM had retinopathy, nephropathy and neuropathy as microvascular complications. Rosenbloom et al. found microvascular complications in 50% of patients with LJM (3). They suggest that after 16 years of diabetes, the risk for microvascular complications is 83% if LJM is present, but only 25% if LJM is absent. Starkman et al. found that LJM is associated with the duration of diabetes in IDDM patients of less than 40 years of age but did not find any relationship in NIDDM patients (7).

Our patient's pulmonary capacity was diminished. Shuyler et al. initially reported the diminished pulmonary capacity in diabetes (8). Barta subsequently described a single patient with limited joint mobility and limited pulmonary capacity (9). The pathologic findings in the thickened skin of subjects with LJM have been postulated to occur in the lungs as well (8).

Our patient also had diabetic hand syndrome superimposed on LJM. In diabetic hand syndrome there is pain, muscle atrophy, neuropathy and disability. None of them are seen in LJM (10). It is likely that people affected with LJM would be at higher risk of developing of other connective tissue proliferative problems and possible neuropathy (2).

Our patient had no improvement in joint range with physical therapy. However Rosenbloom reported the responsiveness of LJM to physical therapy, and Sherry et al. described an adolescent patient with LJM who had improvement in joint range with physical therapy (2,5). It was reported that for the prophylactic treatment of LJM in diabetic patients, range of motion exercises should be done routinely (11).

LJM is a risk factor for the development of microvascular complications in diabetic patients. Earlier detection of LJM may help us to identify patients with an increased risk of microvascular complications. Beginning the prophylactic range of motion exercises after the diagnosis of diabetes may help prevent future contractures and disability in diabetic patients.

Correspondence author:

Serpil SAVAŞ

PK 76

32000 Isparta-TURKEY

References

- Rosenbloom AL, Frias JL. Diabetes; short stature and joint stiffness - a new syndrome. Clin Res 92 A: 22,1974.
- Rosenbloom AL. Limitation of finger joint mobility in diabetes mellitus. J Diabet Complications Apr-Jun; 3 (2):77-87,1989.
- Rosenbloom AL, Silverstein JH, Lezotte DC, Richardson K, McCallum M. Limited joint mobility in childhood Diabetes mellitus indicates increased risk for microvascular disease. N Eng J Med 305:191-4,1981.
- Fernando DJ, Vernidharan J. Limited joint mobility in Sri Lankan patients with non-insulin-dependent diabetes. Br J Rheumatol Mar;36:3,374-6, 1997.
- Sherry DD, Rothstein RL, Petty RE. Joint contractures preceding insulin-dependent diabetes mellitus. Arth Rheum 25 (11):1362-64, 1982.
- Chang K, Uitto J, Rowold EA, Grant GA, Kilo C, Williamson JR. Increased collagen cross-linkages in experimental diabetes, reversal by α -aminopropionitrile and D-penicillamine. Diabetes 29:778-81,1980.
- Starkman HS, Gleason RE, Rand IL, Miller DE, Soeldner JS. Limited joint mobility (LJM) of the hand in patients with diabetes mellitus: relation to chronic complications. Ann Rheum Dis 45:130-5, 1986.
- Schuyler MR, Niewoehner E, Inkley SR, Kohn R. Abnormal lung elasticity in juvenile diabetes mellitus. Amer Rev Resp Dis 113:37-41, 1976.
- Barta L. Flexion contractures in a diabetic child (Rosenbloom Syndrome) Eur J Pediatr 135(1):101-2, 1980.
- Jung Y, Hohmann TC, Gerneth JA. Diabetic hand syndrome. Metabolism 20: 1008-14, 1974.
- Shinabarger NI. Limited joint mobility in adults with diabetes mellitus. Physical Therapy 67(2):215-8, 1987.