

1-1-1998

How Hyperosmolar Solutions (Glucose 30%) May Prevent Stress Induced Ulcers in Rats

Mustafa YANDI

Nazim AĞAOĞLU

Tevhide SEL

Nihat MANSUR

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

YANDI, Mustafa; AĞAOĞLU, Nazim; SEL, Tevhide; and MANSUR, Nihat (1998) "How Hyperosmolar Solutions (Glucose 30%) May Prevent Stress Induced Ulcers in Rats," *Turkish Journal of Medical Sciences*: Vol. 28: No. 1, Article 5. Available at: <https://journals.tubitak.gov.tr/medical/vol28/iss1/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Mustafa YANDI¹
Nazım AĞAOĞLU¹
Tevhide SEL²
Nihat MANSUR¹

How Hyperosmolar Solutions (Glucose 30%) May Prevent Stress Induced Ulcers in Rats*

Received: October 12, 1995

Abstract: It is accepted that administration of hyperosmolar solutions (glucose 30%) prevent stress-induced ulcers in the rats; but their mechanisms of action are not elucidated. In this study, we have investigated whether glucose is absorbed or not through the gastric wall. C14 labelled glucose solution of 30% was given to groups A (n=6), and B (n=8) and physiologic serum to group C(n=7). Restriction of movement with cold application was performed in group B and C. Ulcer indexes were 0.56 ± 0.12 in group A, 0.83 ± 0.27 in group B and 1.63 ± 0.43 in group C. Net absorption rates were 12.2 ± 4.8 %, 23.01 ± 8.3 % and 53.1 ± 4.65 % in

groups A, B, and C, respectively. However, radioactivity levels of the C14-labelled glucose in the gastric wall 13.27 ± 0.02 %, 21.32 ± 0.016 %, and 8.03 ± 0.005 % in groups A, B, and C, respectively. Differences between groups were found to be statistically significant in ulcer index, absorption rate, and amount of radioactivity in the stomach. These results show that glucose is absorbed through the gastric wall and that has a protective effect on stress-induced ulcers in rats.

Key Words: Hyperosmolar Glucose, Stomach, Stress Ulcers..

¹Department of Surgery, Faculty of Medicine, Karadeniz Technical University, Trabzon,

²Department of Biochemistry, Faculty of Veterinary Medicine, Ankara University, Ankara-Turkey

Introduction

It is well known that various nutritional supplements, particularly hyperosmolar glucose, prevent development of stress-induced ulcers in rats and that glucose treatment has been found to be more effective than antacids, H₂ receptor blockers and other medicines (1-4). There is clear evidence that stress ulcers formation (and subsequent bleeding) is caused by excessive gastric acid secretion (5), mucosal ischemia (6), mucosal metabolic energy deficit due to lack of nutrients (7), and possibly an increased permeability to hydrogen ions (8). Various explanations about the protective effect of hyperosmolar solutions exist. Some authors argue that these solutions prevent stress ulcers in rats due to increased endogen prostaglandins in their gastric mucosa (9). Others propose that hyperosmolar glucose supports energy metabolism by direct absorption from the gastric cell but this mechanism has not been proven yet (9,10), while still other authors suggest that glucose is not absorbed from the gastric mucosa (11, 12). We observed in our previous study that intragastric administration of 30% glucose and fructose solutions was absorbed and had a protective effect on stress ulcers as determined by macroscopic evaluations (13, 14). But it

was not clear if this absorption originated from gastric absorption of glucose or dilution of glucose in the gastric lumen. To overcome the dilutional factor of the glucose within the stomach, we planned this experimental study to accurately estimate the C14 glucose absorption rate from the gastric lumen directly.

Material and Methods

The experiment was performed on 22 Swiss-Albino type rats of 160-190 gr. All rats were conditioned for two months on standard rat chow and water. The rats were not given any food, only water, during the last 24 hours before the experiment. Their abdomens were opened by an upper median incision, under ether anaesthesia. Cardia and pylorus were ligated using silk sutures. The stomach was then washed with physiological serum and the contents were aspirated. The rats were divided into 3 groups:

A) Rats given 4 ml glucose solution of 30% + 0.6 ml C14 labelled glucose solution (n=6),

B) Rats given 4 ml glucose solution of 30% + 0.6 ml C14 labelled glucose solution + stress and exposed to cold (n=8),

C) Rats given 4 ml physiologic serum + 0.6 ml




Figure 1. Ulcer Indexes in the groups
* The differences between groups A and C ($p < .01$), B and C ($P < .05$) were significant, A and B ($p > .05$) was insignificant.




Figure 2. Net Absorption Rates of C14 Glucose in the groups
** The differences between all groups were significant ($p < .01$)

C14 labelled glucose solution + stress and exposed to cold (n=7).

After these solutions were given and irrigated to the stomach, 1ml of these solutions were aspirated for a background count of each group.

Preparation of the C14-labelled glucose solution was according to the following procedure: D glucose-UL 14C (Sigma G 4395, 6 mCi/mmol, 0.1 mCi/ml in aqueous sol.) was used for this experiment. 2 ml of D-Glucose-UL 14C were added to 18 ml physiologic serum. Then, 0.6 ml from this solution were used for each animal.

The rats in group A were placed in their cages and

the others (groups B and C) were kept under cold re-stained stress for four hours. At the end of the fourth hour, the rats were sacrificed by an intracardiac injection of sodium pentobarbiturate. The contents in their stomachs were aspirated, their volumes were determined and they were put under deep freeze. The stomach of each animal was removed en bloc, opened along great curvature and spread out on blocks of paraffin. They were washed with physiologic serum, dried by air and the lesions were examined under a dissecting microscope by a pathologist unaware of the treatment groups. Lesions were graded by the Scheric method (15-16). Ulcer indexes were found by dividing length by number of lesions.




Figure 3. Distribution of C14 Glucose in several tissues in rats

*** The differences between the stomach and the other tissues were significant ($p < .01$)

*** The differences between group B and the other groups were significant ($P < .05$)

Blood and tissue samples were taken from the right heart, gastric wall, ileum, liver, spleen; and kidney. Vesica urinaries were empty, so no urine was taken. Two gastric samples from antrum (non ulcer area) and fundus (ulcer area) were taken. Blood and gastric juices were centrifuged; tissue samples were frozen with acetone-solid CO_2 . Liquid and hard tissue samples were stored under deep freeze. All the samples were weighed with a precision balance.

Liquid and tissue samples were counted by beta liquid scintillator in the Department of Biochemistry, Faculty of Veterinary Medicine, Ankara University. Tissues were dissolved by sodium hydroxide (1 mol/L). Their alkalinity was neutralized by nitric acid (1 mol/L) and counted directly by liquid scintillator counter. Scintillation cocktail including triton X-100 was used for the liquid samples (17).

Calculation of dpm (disintegration per minute) of the samples was as follows: First, cpm (count rate per minute) results of all standard solutions and samples were converted to dpm per ml or mg values. Absorption per cent was calculated as follows (18):

Net dpm = Standard dpm/ml of group Samples dpm/ml

Absorption % for solution = $[\text{Net dpm/ml/standard dpm/ml}] * 100$

Absorption % for tissue = $[\text{Tissue samples dpm per mg/Standard dpm per mg}]$

Statistical analyses were performed using a Mann-Whitney U test.

Results

At the end of the experiment, twenty one rats were alive. One rat died due to ether anaesthesia. Ulcer indexes were the lowest for group A (administrated 30% glucose and unstressed) and the highest for group C (administrated physiologic serum and stressed). The differences in ulcer indexes between groups B and C ($p < .05$), and A and C ($p < .01$) were significant. But, no significant difference was found between groups A and B (Fig. 1).

C14-labelled glucose was found to be absorbed throughout the gastric mucosa in all three groups. The absorption rates of glucose in group C were highest and lowest in group A. The differences in absorption rates between all groups were statistically significant ($p < .01$) (Fig. 2).

The differences in C14-labelled glucose level of gastric samples (in both nonulcerative and ulcerative areas) between groups were not statistically significant ($p > .05$). So, stomach glucose levels were taken as the average levels for ulcerated and intact areas of the gastric samples. The mean distribution level of the gastric walls was significantly higher than those of other tissues; liver, ileum, spleen, and kidney ($p < .01$). The mean diffusion rate of group B was significantly different than those of other groups ($p < .05$). In addition, Figure 3 shows distribution level of C14 labelled glucose for ileum, liver, spleen and kidney. The distribution levels of glucose in the various other tissues were not significantly different from each other, and they are negligible.

Discussion

Tube feeding diminishes the incidence of stress bleeding in several types of critically ill patients (17). The physiopathology of stress ulceration is complex and multifactorial. The factors such as secretion of gastric acid (5), mucosal ischemia (6), deficiency of metabolic energy (7) and vasoactive substances as histamin and serotonin due to increasing of H^+ permeability (8) are the possible causes of the ulcers. So, the prophylaxis and treatment of stress ulcers are still not clear yet.

In the present experimental study, 30 percent glucose showed a protective effect from stress-induced ulcers in rats (Fig 1). But the effect of 30% glucose is by a mechanism other than neutralizing ions, since the acid buffering capacity of 30% glucose is equivalent to normal saline solution, a solution with no prophylactic effect in the rat restraint model (11). Intra-gastric administration of 30 percent glucose may have several physiologic effects relevant to its prophylaxis of stress ulceration: systemic absorption and utilization of glucose, elevation of endogen prostaglandins, administration of volume, direct supply of the glucose solution to the gastric mucosa (12).

It is important note that systemic absorption and utilization of glucose was impossible in our experiment because the pylorus and cardias of the rats were ligated.

Elevation of endogen prostaglandins in the gastric wall may have an effect on the prevention of stress ulcers. Since many prostaglandins possess potent vasoactive properties, enhancement of gastric mucosal blood flow by these agents may play an im-

portant role in mediating cytoprotection. A number of prostaglandins are known to exhibit potent vasodilatory actions and affect various vascular beds throughout the body, including those of the gastrointestinal tract. These vasoactive effects are especially evident during parenteral administration and often occur quickly after injection (9). Ephgrave et al. (10) concluded that endogen prostaglandins were useful in gastric mucosal defense but were not necessary for the protection afforded by 30% glucose. They also suggested that 30% glucose was absorbed directly through the gastric mucosa. But the same authors in another experimental study (12) argued that hyperosmolar glucose increased intragastric volume and intraluminal pH. They postulated that these factors may contribute to the protection of gastric mucosa from stress ulceration.

From an experimental study like ours, Mullane J.F. et al concluded that no substantial transport of intraluminal glucose from the stomach occurred (11). They argued that glucose was not directly absorbed through the gastric wall. But they did not directly estimate glucose absorption rate by glycerol or radioactive glucose, they measured only intragastric concentration of glucose so they have had used an empirical method.

In our previous investigation (13, 14) we did not estimate the glucose absorption rate by using glycerol or radioactive glucose. We observed that glucose was absorbed and that energy charges were elevated in the gastric mucosa. We performed this study to accurately estimate the absorption rates of glucose throughout gastric mucosa.

C14-glucose was absorbed through the gastric wall in all groups and the difference in absorption between the groups was statistically significant ($p < .05$, Fig 2).

While absorption in the nonstressed group with administration of 30% glucose was 12.23 ± 4.8 %, the application of stress increased absorption to 23.01 ± 8.3 %, and this was related to mucosal energy deficiency and defect in the mucosal barrier. Absorption rate in the stress applied group with physiological serum, was 53.19 ± 4.65 %. This was referred to the fact that C14-glucose concentration in this group was low as compared to the other groups, in contrast to the hunger of the gastric mucosa for energy. In addition, sodium concentration in the stress applied group with physiological serum (group C) was highest. So, sodium and glucose cotransport or symport

(Na⁺-dependent glucose transport, SG Cotransporter) may facilitate diffusion of the glucose molecules (19).

The difference in distribution of the C14 glucose between the stomach and other tissues was significant ($p < .01$, Fig. 3). The estimated the C14 radioactivity in the gastric wall was highest in group B and lowest in group C. Despite the fact that the absorption rate within the stomach was highest in group C (53.2%), level of the C14-labelled glucose in the gastric wall was the lowest in the C group (8.03 %). This can be explained by the fact that there was only C14 labelled

glucose in the lumen of the stomach in this group, so we postulated that there was a greater utilization of the absorbed glucose. This hypothesis could be supported by estimation of the radioactivity in the expired air, but we did not measure it. The distribution of C14 glucose in the ileum and the other tissues were much lower and not statistically significant (Fig.3). These results show that glucose is absorbed directly through the gastric wall, but not through the ileum and distal gastrointestinal tract, and that it protected rats from stress-induced ulcers.

References

- Lally KP, Andrassy RJ, Wilz WR, Hosbein OP.: Evaluation of various nutritional supplements in the prevention of stress-induced gastric ulcers in rat, *Surg Gyn&Obst* 158: 124-8, 1984.
- Moody FG, Hung LY, Simmons MA: Stress and acute gastric mucosal lesion, *Am J Dig Dis* 21: 148-54, 1976.
- Hase T, Moss BJ.: Microvascular changes of gastric mucosa in the development of stress ulcer in rats, *Gastroenterol* 65:224-34, 1973.
- Skillman JJ, Silen W.: Acute gastroduodenal "stress" ulceration: Barrier disruption of varied pathogenesis. *Gastroenterol* 59: 478-82, 1970.
- Menguy R, Masters YF.:Gastric mucosal energy metabolism and "stress ulceration", *Ann Surg* 180:538-48, 1974.
- Bounous G, Sutherland NG, Mc Ardle H, Gurd FN.: The prophylactic use of an elemental diet in experimental hemorrhagic and intestinal ischemia. *Ann Surg* 166:312-38, 1967.
- Menguy R, Respailletes L and Masters YF.: Mechanism of stress ulcer:Influence of hypovolemic shock on energy metabolism in the gastric mucosa, *Gastroenterol* 66: 46-55, 1974.
- Altamarino M.: Back diffusion of H⁺ during gastric secretion, *Amer J Physiol* 218:1-16, 1970.
- Miller TA.: Protective effects of prostaglandins against gastric mucosal damage: current knowledge and proposed mechanism "Editorial Review". 1983 *The Am Phys Soc: G601-G623*.
- Ephgrave KS, Horton JW, Burns DK. Hyperosmolar glucose prevents stress ulceration in the rat restraint model despite inhibition of endogenous prostaglandins, *Surg Gyn&Obst* 164: 9-16, 1987.
- Mulane FJ, Wilfong RG: Starvation, glucose, and stress ulcers in the rat, *Arch Surg* 109. 416-19, 1974.
- Ephgrave KS, Kleiman-Wexler RL, Adair CG. Enteral nutrients prevent stress ulceration and increase intragastric volume, *Crit Care Med* 18/6: 621-4, 1990.
- Yandi M. Protective effect of honey and 25% dextrose on stress-induced ulcers in rats. "Letters to the editor. *J Islam Acad Sci* 1: 171, 1988.
- Yandi M, Değer O, Turgutalp HN, Gacar N. The protective effects of glucose solutions on stress-induced ulcers in rats: supporting of mucosal energy metabolism, *Gastroenteroloji* 2/2: 193-197, 1991.
- Lockenvitz E, Schwillle PO, Hanish E, Engelhardt. Influence of exogenous glucagon on gastric secretion, mucosal blood flow and stress ulcers in the rat doseresponse results during resting conditions and immobilization stress, *Res Exp Med* 182: 245-53, 1983.
- Hanisch E, Schiwille PO, Engelhardt W. Effects of various vagotomies and sympathectomies on gastric secretory functions and related hormones in the non-stressed and stressed rat. *Scand J Gastroent* 19 suppl. 89: 99-104, 1984.
- Martin LF, Booth FVM, Reines D. Stress ulcers and organ failure in intubated patients in surgical intensive care units, *Ann Surg* 215/4: 332-7, 1992.
- Bergmeyer HU. *Methods of Enzymatic Analysis*, 3. Ed. VCH, Weinheim, Germany, 1986, 428-38.
- Ganong WF. *Review of Medical Physiology*, 16 ed. A Lange Medical Book Middle East Ed. 1993, 429-30.