

2023

Effects of progressive functional ankle exercises in spastic cerebral palsy, plantarflexors versus dorsiflexors: a randomized trial

MELEK VOLKAN YAZICI

GAMZE ÇOBANOĞLU

GÖKHAN YAZICI

BÜLENT ELBASAN

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

YAZICI, MELEK VOLKAN; ÇOBANOĞLU, GAMZE; YAZICI, GÖKHAN; and ELBASAN, BÜLENT (2023) "Effects of progressive functional ankle exercises in spastic cerebral palsy, plantarflexors versus dorsiflexors: a randomized trial," *Turkish Journal of Medical Sciences*: Vol. 53: No. 5, Article 17. <https://doi.org/10.55730/1300-0144.5682>

Available at: <https://journals.tubitak.gov.tr/medical/vol53/iss5/17>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Effects of progressive functional ankle exercises in spastic cerebral palsy, plantarflexors versus dorsiflexors: a randomized trial

Melek VOLKAN YAZICI^{1*}, Gamze ÇOBANOĞLU², Gökhan YAZICI², Bülent ELBASAN²

¹Department of Physical Therapy and Rehabilitation, Faculty of Health Sciences, Yüksek İhtisas University, Ankara, Türkiye

²Department of Physical Therapy and Rehabilitation, Faculty of Health Sciences, Gazi University, Ankara, Türkiye

Received: 02.02.2023 • Accepted/Published Online: 31.05.2023 • Final Version: 26.10.2023

Background/aim: Children with cerebral palsy (CP), even those who have very mild impairment, have lower muscle strength than their typically developing peers. The ankle dorsiflexors (DFs) and plantarflexors (PFs) of children with CP are especially weak. Weakness in the ankle muscles causes problems in functional skills, mobility, and balance in spastic CP (SCP). The aim of this study was to investigate the effects of progressive functional exercises (PFEs) on the DF, PF, or dorsi-plantar flexor (DPF) muscles in children with SCP, specifically, the functional mobility, balance, and maximum voluntary contraction (MVC), and compare the effects of strengthening these muscles individually or combined.

Materials and methods: This randomized trial was conducted between December 1st, 2018, and May 15th, 2019, at Gazi University, Department of Physiotherapy and Rehabilitation. Randomly assigned into groups were 27 independently ambulant patients with unilateral/bilateral SCP, where PFEs were applied to the DF, PF, or DPF muscles. Muscle tone, balance, and functional mobility were assessed. The MVC was evaluated by surface electromyography. PFEs were performed 4 times a week, for 6 weeks.

Results: The spasticity of the PF muscles decreased in all of the groups. PFE of the DF muscles led to an increase in ankle joint range of motion (ROM) and improved functional mobility ($p < 0.05$). PFE of the PF muscles resulted in improvements in balance and functional mobility ($p < 0.05$). PFE of the DPF muscles brought about improvements in balance but not in functional mobility ($p < 0.05$). No significant difference in the MVC was observed in any of the groups ($p > 0.05$).

Conclusion: Gains are obtained according to the function of a muscle group. By training the DF muscles, it is possible to improve function and ROM. Furthermore, training the PF muscles led to improvements in balance and functional mobility, indicating that it is possible to bring about positive changes in spastic muscles. This study showed that muscle groups must be exercised according to the intended goal.

Key words: Cerebral palsy, exercise, gastrocnemius muscle, anterior tibial muscle, posture balance, surface electromyography

1. Introduction

Spastic cerebral palsy (SCP) is the most common type of CP, which leads to motor deficits and disorders in the development of movement and posture. Motor deficits of SCP include negative phenomena such as weakness, fatigue, incoordination, and positive phenomena such as spasticity, clonus, rigidity, and spasms [1]. As a result of these deficits, children with SCP are likely to have developmental delays in gross motor function, standing, walking, and balance when compared to normally developing children [2].

Weakness is a prevailing impairment in individuals with CP and is caused by impaired neural ability and altered intrinsic capacity of the muscles [3]. Studies investigating the relationship between spasticity and strength in SCP have demonstrated that muscle weakness has a direct effect

on function and leads to more limitations in function than spasticity [4–6]. Muscle weakness in SCP was reported to be associated with deficiencies in walking, running and a decrease in functional scales, consequently, weakness causes a restriction in participation in daily life activities [5, 7, 8]. Children with CP, even those who have very mild impairment, are weaker than their typically developing peers [9, 10]. For example, compared to the strength of the muscles in children with typical development, the muscle strengths of the ankle dorsiflexors (DFs) and plantarflexors (PFs) of children with CP were about 50% and 35% compared to their typically developing peers, respectively [11]. The DFs were found to be far weaker compared to the PFs [3]. Previously, the muscles of children with CP were thought to be incapable of adapting; however, it is now known that muscles in CP are indeed plastic [12]. The

* Correspondence: melek_volkan89@hotmail.com

adaptations and improvements that can occur following strength or power training in the muscles of children with CP has been shown; however, the effect of progressive functional exercises (PFEs) is still unclear [7, 13, 14].

The foot and ankle joints are affected in almost all individuals with CP [10]. A significant decrease was demonstrated in tibialis anterior activation in children with CP when compared to their typically developing peers [15]. The PFs of the foot, which are the gastrocnemius and soleus muscles, resist and control the forward rotation of the tibia over the foot during the mid and terminal stance phases of gait [16]. The eccentric, antigravity activity of the soleus muscle is of primary importance due to modulating the site of application of the ground reaction force on the foot during weight bearing, thus allowing a biomechanical advantage in ambulation [17, 18].

It has been demonstrated that in patients with CP, exercises targeting the knee and hip muscles have beneficial effects on gait and function [19–22]. However, studies demonstrating the effects of exercises applied on ankle muscles on functional mobility are insufficient, and there are no studies examining the effect of PFE of these muscles on balance or muscle activation [23, 24]. PFE is a type of exercise with an overall goal of functional independence, which helps patients develop skills to perform activities of daily life easier [25]. PFEs consist of many daily activities involving the lower limbs, such as sit-to-stand and stair climbing [26]. It was reported that when exercises involve practicing functional closed-kinetic-chain exercises, gains in strength may lead to greater improvements in functional motor performance [26, 27]. Therefore, the primary aim of this study was to investigate the effects of PFE on muscle tone, balance, and functional mobility in patients with SCP. The second aim was to investigate the changes in the maximum voluntary contractions (MVC) of the lower extremity muscles following 6-weeks of PFE using surface electromyography (sEMG).

The primary hypothesis herein was that balance and functional mobility would improve following PFEs in patients with SCP.

2. Materials and methods

2.1. Study design

This study was planned as a randomized clinical trial. The randomization was performed by a physiotherapist blinded to the study using an internet-based method (randomizer.org). The study population comprised 27 participants who were randomized into 3 groups:

- Dorsiflexor (DF) group
- Plantar flexor (PF) group
- Dorsi-plantar flexor (DPF) group.

2.2. Participants

The participants included children aged 5–15 years with

unilateral or bilateral SCP. Inclusion criteria were being level I or II according to the Gross Motor Function Classification System (GMFCS) [28], and being able to understand and follow simple instructions. Children who had a condition which limited the ability of exercising, had received botulinum toxin injections in the past 6 months, and had undergone an orthopedic operation in the past year were excluded. All of the children and parents provided written informed consent. The study was approved by the ethical review board of Gazi University with approval date and number 26.11.2018-863. The study was conducted between December 1st, 2018, and May 15th, 2019, at Gazi University, Department of Physiotherapy and Rehabilitation, Ankara, Türkiye. The authors conformed to the ethical guidelines of the 1975 Declaration of Helsinki. The study was registered on clinicaltrials.gov with ID: NCT03901703.

2.3. Outcome measures

Each participant was assessed individually to avoid rivalry. It was ensured that the participants learned all of the tests correctly before they were performed to eliminate the learning effect. Time was provided between the assessments to ensure at the children had enough time to rest and reduce the effect of fatigue. The assessments were performed by the same therapist, in the same order at baseline, and following 6-weeks of intervention to avoid inter-rater bias.

The assessment of dynamic muscle length was performed using the Modified Tardieu Scale (MTS). This scale measures the point of resistance to a rapid velocity stretch and is performed twice; rapidly and slowly. A 'catch' resulting from the 'overactive stretch reflex' may be felt in the rapid range of motion (ROM) at a particular angle. This angle was defined as R1. The slow passive joint ROM was measured with goniometry and recorded in a standardized format. This gave an indication of muscle length at rest, or R2. Muscle testing for R1 and R2 was performed using the protocol and test positions stated by Tardieu et al. [29, 30]. A large difference between R1 and R2 implies that there is a large dynamic component due to spasticity, whereas a small difference between them means that there is predominantly fixed contracture in the muscle.

The Pediatric Balance Scale (PBS), a modification of the Berg Balance Scale, was developed as a balance measure for participants [31]. The scale consists of 14 items that are scored from 0 points (lowest function) to 4 points (highest function) with a maximum score of 56 points [32].

The Functional Reach Test was used to measure dynamic balance. The distance that an individual is able to reach forward from a starting standing position with a fixed base of support without loss of balance is measured using a measuring tape at the level of the acromion [33].

This test has shown validity and reliability in children with CP [34].

The Timed Up and Go Test (TUG) measures the time taken by an individual to stand up from a chair, walk a distance of 3 m, turn around, walk back to the chair, and sit down. This test is used in children with CP who are ambulatory to assess dynamic balance and functional mobility [35]. The child was seated with their feet flat on the floor with their hip and knees in 90° flexion. A marking tape was used to stick a mark on the floor at a distance of 3 m from the chair to indicate where the child must turn around. The timing of the test began upon movement to stand after the cue “ready, go” and concluded when the child sat down on the chair. The time needed to complete the test was recorded. The children performed 1 practice trial, followed by 3 test trials. The minimum time was recorded for each trial [36].

For the measurement of sEMG, an 8-channel noninvasive sEMG Noraxon MiniDTS (Noraxon USA Inc., Scottsdale, AZ) system was used. Disposable, self-adhesive Ag/AgCl electrodes (Noraxon Dual EMG Electrode) were used to record the EMG signals. Each electrode was placed parallel to the muscle fibers evaluated, according to the guidelines recommended by the sEMG for the noninvasive assessments of muscles (SENIAM) [37]. The MVCs of the tibialis anterior, gastrocnemius (medial and lateral heads), vastus medialis, and medial hamstrings were measured. The MVCs were recorded by enabling the child to maximally contract the muscle against resistance provided by the therapist. This was done to ensure that the muscle maintained maximum contraction while recording. All of the measurements were performed in 2 trials for each region and the average of these measurements was taken. The raw sEMG signals were passed through 20 Hz, infinite impulse response (IIR), Butterworth high-pass and 500 Hz, IIR, Butterworth low-pass motion artefact filters. Then, the root mean square (RMS) values were calculated from the raw sEMG data in successive time windows (0.1 s) in order to evaluate the signals.

2.4. Intervention

The PFE protocol was applied for 6 weeks [21]. Each week, 4 sessions were applied: 2 sessions, at 45 min/session with a physiotherapist, and also 2 sessions/week, for 30 min at home under the supervision of a parent/caregiver. All of the exercises were applied with 3 sets and 10 repetitions in each set. Continuity was checked and necessary adjustments were made by interviewing the parents of the participants weekly.

All of the children, regardless of their group, participated in a functional exercise program consisting of exercises targeting the abdominal and back muscles, hip extensors, abductors, and the quadriceps muscles. The loading of the exercises was increased by altering the ROM

of the joints, the difficulty of the exercise, or by changing the height/stability of the surfaces. The description, progression, and photographs of the exercise protocol can be seen in Appendix 1.

2.4.1. DF group exercises

In this group, the participants were given 2 exercises. The first was walking up on a ramp and then walking down on the ramp backwards. The second was dorsiflexion of the feet while sitting with the knees in increasing range of flexion. The difficulty of movement was adjusted by changing the degree of flexion in the knee while sitting.

2.4.2. PF group exercises

Participants in the PF group were given calf raises to train the PF muscles. The difficulty of the exercises was increased by changing the angle of plantar flexion. At the beginning, the movement was initiated from the position where the PF muscles were in their shortened position. At the end of the week 2, the exercise had proceeded to the position where the foot was flat and in contact with the ground. At the end of week 4, plantar flexion was initiated from the position where the muscles were elongated, and the foot was in DF.

2.4.3. DPF group exercises

The participants in this group performed the exercises given in both the DF and PF groups.

2.5. Statistical analysis

Statistical analysis was performed using IBM SPSS Statistics for Windows 21.0 (IBM Corp., Armonk, NY, USA). The variables were investigated using visual (histograms, probability plots) and analytical (Shapiro-Wilk test) methods to determine whether they were normally distributed. Nonnormally distributed data were expressed as the median and interquartile range (IQR). The Wilcoxon test was used to compare the changes in the groups. When the level of significance was different in all 3 groups, pre-post measurement differences were determined using the Kruskal-Wallis test. When the level of significance was different in 2 groups, pre-post measurement differences were determined using the Mann-Whitney U test. The level of statistical type-1 error was set as $p < 0.05$. Based on the results of a previous study, it was estimated that a sample size of 27 participants (9 per group) would have a power of 80% to detect statistically difference in balance analysis (the PBS) for a value of 0.05 [38].

3. Results

A total of 27 participants completed this study, with 9 participants in each group. The flow chart of participation in the study can be seen in Figure 1. Demographic information regarding the participants is shown in Table 1. The participants were included in the rehabilitation program for 4 sessions a week, for 6 weeks. A minimum of 22 and




Figure 1. Flow chart of the participation in the study.

Table 1. Demographic characteristics and functional status of the participants.

| | DF group median (IQR) n: 9 | PF group median (IQR) n: 9 | DPF group median (IQR) n: 9 | p-value |
|-------------|----------------------------------|----------------------------------|-----------------------------------|---------|
| Age (years) | 8 (6.5/12.5) | 8 (6.5/13) | 6 (5.5/11.5) | 0.599 |
| Height (cm) | 130 (115.5/156.5) | 136 (120/151) | 120 (116.5/154.5) | 0.759 |
| Weight (kg) | 23 (18.5/49) | 28 (23.5/33) | 20.8 (18.3/55.4) | 0.832 |
| | n (%) | n (%) | n (%) | |
| Sex | Male | 5 (55.6) | 4 (44.4) | 0.648 |
| | Female | 4 (44.4) | 5 (55.6) | |
| GMFCS | I | 6 (66.7) | 4 (44.4) | 0.565 |
| | II | 3 (33.3) | 5 (55.6) | |

DF: Dorsiflexor, PF: plantarflexor, DPF: dorsi-plantar flexor, IQR: inter quartile range, cm: centimeters, kg: kilograms, GMFCS: Gross Motor Function Classification System.

a maximum of 24 rehabilitation sessions were performed per participant. No adverse effect or event was seen in any of the groups. None of the participants reported discomfort or muscle soreness.

When the MTS results were examined, statistically significant differences were observed in the rapid (R1) assessments of the affected side PFs in all of the exercise groups,

and in the less affected side PFs of the DF group. Furthermore, it can be seen that there was statistically significant difference in the slow (R2) assessment of the affected side PFs in the DF group. These results can be seen in Table 2, ($p < 0.05$). When the level of effectiveness between the groups was examined, all three groups were shown to be equally effective on muscle tone (Table 2, $p > 0.05$).

Table 2. Comparison of the MTS measurements within and between the groups before and after PPE treatment.

| Plantarflexors | DF group n: 9 | | | | | PF group n: 9 | | | | | DPF group n: 9 | | | | | | |
|--------------------------|---------------------------|--------------------------|-----------------------------|--------------------------------|---------------------------|--------------------------|-----------------------------|--------------------------------|---------------------------|--------------------------|-----------------------------|--------------------------------|---------------------------|--------------------------|-----------------------------|--------------------------------|------------------------------|
| | Before median (IQR) | After median (IQR) | p-value within groups | Change (Δ) median IQR | Before median (IQR) | After median (IQR) | p-value within groups | Change (Δ) median IQR | Before median (IQR) | After median (IQR) | p-value within groups | Change (Δ) median IQR | Before median (IQR) | After median (IQR) | p-value within groups | Change (Δ) median IQR | p-value between groups |
| Affected side | Slow (R2) (°) | 10 (3.5/12.5) | 15 (10/20) | *0.015 | | 10 (7/15) | 10 (10/20) | 0.197 | | 10 (2.5/20) | 15 (9/20) | 0.104 | | | | | |
| | Rapid (R1) (°) | 0 (-1.5/7.5) | 12 (7.5/17.5) | *0.008 | 5 (0/6.5) | 10 (5/15) | *0.042 | 5 (0/15) | 0 (-7.5/7.5) | 10 (9/15) | *0.017 | 0 (0/0) | | | | | 0.140 |
| | Difference | 5 (0/10) | 0 (0/6.5) | 0.092 | | 6 (3/9.5) | 5 (0/7.5) | 0.260 | | 10 (5/15.5) | 5 (0/7.5) | *0.049 | | | | | |
| Less affected side | Slow (R2) (°) | 20 (11.5/20) | 20 (15/20) | 0.465 | | 15 (10/20) | 20 (13.5/20) | 0.465 | | 20 (17.5/20) | 20 (10/20) | 0.102 | | | | | |
| | Rapid (R1) (°) | 10 (10/20) | 20 (15/20) | *0.039 | | 10 (5/11) | 15 (11/20) | 0.092 | | 20 (15/20) | 20 (10/20) | 1.000 | | | | | |
| | Difference | 0 (0/7.5) | 0 (0/0) | 0.066 | | 5 (0/10) | 0 (0/0) | 0.206 | | 0 (0/2.5) | 0 (0/0) | 0.180 | | | | | |

(°): Degrees, R1: the first point of resistance felt by the therapist due to the catch resulting from the overactive stretch reflex; R2: the passive joint ROM. *Indicates that there was a statistically significant difference (*p < 0.05 within the groups after 6 weeks of PPE treatment [Wilcoxon signed-rank test] and changes between the groups [Kruskal-Wallis test]).

When the PBS results were examined no statistically significant difference was observed in the DF group (Table 3, $p > 0.05$), however, statistically significant differences were seen in the PF and DPF groups. When the level of effectiveness between PF and DPF groups was examined, it can be seen that there was no difference between the groups, as seen in Table 3, $p < 0.05$. According to our results, none of the children in the DF group developed at a level of clinical significance (minimal clinically important difference (MCID): 5.83 points) whilst two children in each of the PF and DF + PF groups, had shown development in balance at a clinically significant level. The MCID can be seen in Figure 2. When the functional reach tests were examined, there were statistically significant improvements only in the PF group as seen in Table 3, $p < 0.05$.

When the TUG test scores were compared, statistically significant changes were found in the DF and PF groups as seen in Table 3 ($p < 0.05$), however no statistically significant improvement was found in the DPF group (Table 3, $p > 0.05$). These differences appeared to be equally significant between the groups (Table 3, $p > 0.05$).

As a result of the sEMG analysis performed to evaluate MVC before and after treatment, significant changes were found in the affected Tibialis Anterior muscle and the less affected Gastrocnemius Medialis muscle in the PF group as seen in Table 4 ($p < 0.05$). No statistically significant difference was found in the results of the other groups.

5. Discussion

This study is the first study to investigate the effects of PFEs of the DF, PF and DPF muscles and interpret the effects of PFEs on these muscle groups separately.

The MTS assessments showed an increase in ROM (R2) in the DF group and an increase in the angle of muscle reaction (R1) in all of the groups. Accordingly, this indicated that there was a significant decrease in spasticity of the PF muscles in all of the groups. Achieving significance in all of the groups revealed that exercises targeting the ankle muscles may generally decrease spasticity. It is our belief that the DF exercises included in this study may have generated an active stretching effect on the PF muscles with every movement, resulting in an increase in the ankle DF joint ROM. The study conducted by dos Santos et al. on closed kinetic chain exercises showed that PFE effectively increase lower extremity muscle strength, thereby facilitating lower extremity contraction and allowing agonist and antagonist muscles to work effectively. As a result, they stated that this led to reduction of muscle tone in the lower extremities [39]. The exercises in the current study are believed to have generated similar results.

This study was the first to investigate the effects of PFEs on balance. It was seen that in groups including

plantar flexion exercises, namely the PF and DPF groups, significant changes in the total score of the PBS were evident. When the functional reach test scores were examined, it was seen that statistically significant differences were only established in the PF group. It can be concluded that when PFEs are performed, greater improvements can be achieved in balance. PFEs alone resulted in improvement in static balance. The fact that there was no change in balance as a result of PFEs on the DF group muscles suggests that DF muscles are not as active as PF muscles in balance strategies. Considering that the PF soleus muscle is an important antigravity muscle, the results found herein support these findings [40].

According to the TUG results, there was a significant increase in the functional mobility of the DF and PF groups following the PFEs. The reason for the increase seen in the DF group can be attributed to the development of a more effective gait with the improvement in the DF ROM, and the improvement in the PF group may have been due to the increased balance skills in the PF muscles. No improvement was noted in the DPF group. This finding shows that muscle-specific PFEs lead to better results and that it is necessary to focus directly on a muscle group in order to improve functional mobility. In a study by Schranz et al., 20 patients, aged 6–16 years, with unilateral and bilateral spastic CP were randomly assigned to either progressive resistance exercise or functional exercise groups [41]. According to the results of their study, the increase in strength reached statistical significance only in the functional exercise group. The exercises used in their study were similar to those used in the PF group herein. In addition to their study, significant improvements in balance and TUG were achieved in the current study. In our opinion, this further gain was due to the fact that focused was placed on the foot-ankle area, which is the most affected area in CP patients [42]. From this point of view, the improvements obtained in balance and tone of the PF muscles will contribute greatly to functional abilities.

Lorentzen et al. investigated the effectiveness of gait training performed on an inclined treadmill for 6 weeks in a study involving 32 adults with CP [43]. While the control group continued their routine exercises, the study group practiced walking on an inclined treadmill. As a result of the study, compared to the control group, it was found that there was a statistically significant improvement in gait speed and passive ankle joint stiffness in the experimental group. Walking on an inclined surface was one of the exercises given in the present study, targeting the DF muscles. In addition to these results, statistical improvements were also found in the TUG and muscle tone in the current study. This difference may have occurred due to the younger age of the participants. It is

Table 3. Comparison of the balance and functional test measurements within and between the groups before and after PFE treatment.

| Before median (IQR) | DF group n: 9 | | | | PF group n: 9 | | | | DPF group n: 9 | | | | | | | |
|---------------------|--------------------|-----------------------|-----------------------|---------------------|--------------------|-----------------------|-----------------------|---------------------|--------------------|-----------------------|-----------------------|---------------------|--------------------|-----------------------|-----------------------|------------------------|
| | After median (IQR) | p-value within groups | Change (Δ) median IQR | Before median (IQR) | After median (IQR) | p-value within groups | Change (Δ) median IQR | Before median (IQR) | After median (IQR) | p-value within groups | Change (Δ) median IQR | Before median (IQR) | After median (IQR) | p-value within groups | Change (Δ) median IQR | p-value between groups |
| Affected side | 28 (23.5/29.5) | 29 (24.5/32.5) | 0.307 | | 22 (12/28) | 29 (23/31) | *0.008 | | 24 (15.5/28) | 19 (16.5/32) | 0.398 | | 24 (15.5/28) | 19 (16.5/32) | | |
| Less affected side | 29 (23.5/33) | 30 (26.5/34.5) | 0.173 | | 23 (13/30) | 30 (25/31) | *0.011 | | 27 (17/35.5) | 25 (25/33) | 0.888 | | 27 (17/35.5) | 25 (25/33) | | |
| PBS | 53 (49/55) | 53 (50.5/56) | 0.344 | | 46 (44.5/54.5) | 51 (49/56) | *0.017 | | 49 (46/52.5) | 52 (50/5405) | *0.018 | | 49 (46/52.5) | 52 (50/5405) | | 2 (0.5/6) |
| TUG | 6.68 (5.28/7.85) | 5.68 (5.01/6.50) | *0.038 | -0.68 (-1.56/-0.08) | 7.45 (5.76/8.65) | 6.62 (5.53/7.83) | *0.042 | | 6.16 (6/8.06) | 6.22 (5.60/7.13) | 0.660 | | 6.16 (6/8.06) | 6.22 (5.60/7.13) | | |

PBS; Pediatric Balance Scale, TUG; Time Up and Go test. * Indicates that there was a statistically significant difference (*p < 0.05 within the groups after 6 weeks of PFE treatment [Wilcoxon signed-rank test] and changes between the groups [Mann-Whitney U test]).




Figure 2. MCID of the PBS results. PBS: Pediatric Balance Scale.

known that, as individuals with CP get older, the severity of the pathologies that occur in their joints increases, their soft tissues adapt to these changes over time and are less sensitive to the interventions. For this reason, it is our belief that an exercise such as walking upwards on a ramp or an inclined surface may have a positive effect on DF muscles when performed at a younger age since the joints may be more open to development with intervention, and this may have led to improvements in the functional tests. In the study by Hye-Jin Cho, 25 participants with CP were randomized into 2 groups, as a functionally progressive resistance exercise group and a control group [27]. The participants in the intervention group participated in functional progressive resistance exercises for 30 min/day, 3 times/week, for 6 weeks. At the end of their study, the intervention group showed improvements in muscle tone, dynamic balance, and functional ability. Thus, when the results of their study and those herein are considered, it can be seen that even 6 weeks of functional training can be beneficial in many aspects. In a study conducted by Ryan et al., children with SCP were included in resistance training of the ankle PFs for 10 weeks. As a result, the authors stated that resistance training did not improve muscle strength, activity, or participation and that adverse effects were seen in 90% of the participants [44]. In the present study, the ankle PFs were trained using functional exercises and body weight only. The loading of the exercises was achieved by

changing the foot position and as a result, improvements were seen in balance and functional mobility, a decrease in spasticity was achieved and furthermore, there were no adverse effects in any of the participants. A study by Surana et al. investigated the effects of functional lower extremity exercises in SCP and similar to the current study, they put forth promising results in gait capacity and performance [45]. Current guidelines on exercise interventions in CP also state that exercises must be functional and in a real-life context [46]. Therefore, in our opinion, PFEs may be more beneficial than resistance exercises in the ankle area.

In the current research, the MVC results showed that the muscles included in the sEMG exhibited different degrees of activity. While there were increases in the MVCs of some of the muscles, decreases were evident in others. In our opinion, the significant changes found in the affected tibialis anterior muscle and the less affected gastrocnemius medialis muscle in the PF group occurred coincidentally, and an interpretation is not possible due to many factors. In the literature, studies using sEMG have reported that reciprocal inhibition is insufficient in patients with CP and may cause problems during muscle relaxation, leading to errors in MVC. Furthermore, it was reported that obtaining isolated muscle contraction in patients with SCP is very difficult [47]. According to previous sEMG studies, there may be coactivations in the muscles, which may also lead to errors in the results [48, 49]. Rose et al. stated that type I muscle fibers are more common in the muscles of patients

Table 4. Comparison of the MVC of the muscle measurements with sEMG within the groups before and after PFE treatment.

| | DF group n: 9 | | | | PF group n: 9 | | | | DPF group n: 9 | | | |
|------------------------------|---------------------------|--------------------------|-----------------------------|---------------------------|--------------------------|-----------------------------|---------------------------|--------------------------|-----------------------------|---------------------------|--------------------------|-----------------------------|
| | Before median (IQR) | After median (IQR) | p-value within groups | Before median (IQR) | After median (IQR) | p-value within groups | Before median (IQR) | After median (IQR) | p-value within groups | Before median (IQR) | After median (IQR) | p-value within groups |
| Vastus medialis obliquus | Affected side | 9.2 (3.5/46.5) | 28.5 (7.1/57.8) | 0.374 | 49.4 (16.3/110.3) | 13.9 (6.1/47.8) | 0.314 | 57.2 (42.9/74.4) | 39.8 (20.4/61.8) | 0.214 | | |
| | Less affected side | 46.7 (13.7/61.3) | 27.3 (6.5/52.1) | 0.173 | 71.9 (4.5/106.2) | 10.4 (3.5/81.1) | 0.038 | 46.8 (7.9/61.8) | 41.6 (12.1/64.2) | 0.889 | | |
| Semitendinosus | Affected side | 21.9 (9.9/87) | 34 (12.1/71.8) | 0.859 | 10.9 (6.4/98.2) | 7.3 (5/28.8) | 0.260 | 38.3 (21.3/80) | 35.9 (13.4/87.9) | 0.515 | | |
| | Less affected side | 17.6 (11.7/71) | 28.8 (12.9/76.7) | 0.953 | 60.6 (5.9/144.5) | 12.5 (5.4/68.6) | 0.173 | 55.9 (16/144.3) | 59 (37.4/106.3) | 0.767 | | |
| Tibialis anterior | Affected side | 53.6 (14.8/123.8) | 15 (9.7/30) | 0.051 | 66.8 (49.3/85.7) | 30.5 (5.2/47) | *0.021 | 26.4 (10.1/87.8) | 40.4 (13.2/59.3) | 0.374 | | |
| | Less affected side | 83.6 (15.7/156) | 49.9 (11.7/160.5) | 0.678 | 66.8 (34.9/162) | 37.8 (20/77.9) | 0.051 | 78.2 (13.6/132.5) | 92.7 (12.6/145.3) | 0.678 | | |
| Gastrocnemius me- dialis | Affected side | 17.3 (8.2/40.3) | 16.8 (6.6/46.9) | 0.859 | 17.6 (4.3/73) | 13.5 (4.1/48.8) | 0.594 | 10.1 (3.8/38.7) | 6.8 (2.2/29.4) | 0.173 | | |
| | Less affected side | 15.9 (3.8/32.9) | 40.8 (8.2/65.2) | 0.173 | 79.6 (20.8/105) | 25.2 (9.8/39.1) | *0.021 | 15 (6.5/73.5) | 21.3 (5.8/71.7) | 0.767 | | |
| Gastrocnemius late- ralis | Affected side | 14.4 (5.1/33.7) | 30.3 (10.5/43.1) | 0.374 | 7.9 (3.2/47.3) | 10 (4.7/33.3) | 0.767 | 16.8 (6.9/25.1) | 22.7 (4.4/41.3) | 0.374 | | |
| | Less affected side | 26.7 (7.4/65.2) | 36.7 (18.7/53) | 0.441 | 51.9 (10.2/103.7) | 40.9 (8.9/56.9) | 0.110 | 27.1 (5.8/36) | 14.1 (6.2/28) | 0.441 | | |

*Indicates that there was a statistically significant difference (*p < 0.05 within the groups after 6 weeks of PFE treatment [Wilcoxon signed-rank test]).

with SCP. High muscle fiber firing cannot be achieved in MVC measurements, and therefore, may also lead to errors in measurements [49]. In addition to these factors, we believe that the pathological changes occurring in the muscles in patients with SCP may have caused a shift in the reference points, which are important for accuracy in sEMG measurements. Superficial electrodes are placed on the motor points of the relevant muscles. It was reported by SENIAM that the most accurate measurements can be made from these points [37]. However, these reference points are determined for healthy individuals who show typical development. It is known that pathological changes that directly affect muscle fibers in SCP lead to elongation in the sarcomere length of the muscles and changes in the fiber type [50]. Furthermore, it was reported that motor points in the muscles of patients with CP may be at different points in the muscle compared to their healthy peers [51]. These factors may have led to an increase in the signals received from neighboring regions during measurements, which is known as *physiological crosstalk*. It is our belief that there may not have been a significant difference in muscle activation due to these factors.

Even though the sEMG results in the current study showed that there was no statistically interpretable difference in the MVC, it can be seen that PFEs led to positive effects on function, muscle tone, and balance according to the muscle group focused on. Boyd et al. reported that in order to achieve functional development in patients with CP, exercises must consist of functional movements and should be practiced in a context-specific manner [46]. The present research supports these findings. Therefore, we believe that this study will guide physiotherapists working in the rehabilitation of patients with CP on how to achieve beneficial results with functional exercise.

This study had some limitations. Although there was an improvement in function, the effect of the PFE on strength was not measured. Such effects could have been reported objectively with the use of a reliable method to measure muscle strength. Furthermore, a larger sample

size may have yielded more power in detecting statistically significant relationships. In future studies, the effectiveness of PFEs on strength should be investigated with a longer exercise duration, with objective measures for strength, and a larger sample size.

The results herein revealed that functional improvement is exercise-specific. Gains are obtained according to the function of the muscle group. It was seen that the PFEs of the PF muscles of patients with CP led to an improvement in balance, whereas the PFEs of the DF muscles led to an increase in the ROM of the ankle joint. When PFEs were performed on both the DF and PF muscles, the effects of both exercise groups appeared in combination; however, the results were not as effective as exercising the PF muscles individually.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest

The authors have no competing interests to declare.

Informed consent

Due to the nature of this research, participants of this study did not agree for their data to be shared publicly, so supporting data are not available.

Acknowledgment

The authors would like to thank Prof. Dr. Nevin Atalay Güzel for her assistance in the study.

Ethical approval

The study was approved by the ethical review board of Gazi University with approval date and number of 26.11.2018-863. The authors conformed to the ethical guidelines of the 1975 Declaration of Helsinki. All of the children and parents provided their written informed consent.

References

1. Jan MM. Cerebral palsy: comprehensive review and update. *Annals of Saudi Medicine* 2006; 26 (2): 123-132. <https://doi.org/10.5144/0256-4947.2006.123>
2. Wimalasundera N, Stevenson VL. Cerebral palsy. *Practical Neurology* 2016; 16 (3): 184-194. <https://doi.org/10.1136/practneurol-2015-001184>
3. Hanssen B, Peeters N, Vandekerckhove I, De Beukelaer N, Bar-On L et al. The contribution of decreased muscle size to muscle weakness in children with spastic cerebral palsy. *Frontiers in Neurology* 2021; 26 (12): 692582. <https://doi.org/10.3389/fneur.2021.692582>
4. Mockford M, Caulton JM. The pathophysiological basis of weakness in children with cerebral palsy. *Pediatric Physical Therapy* 2010; 22 (2): 222-233. <https://doi.org/10.1097/PEP.0b013e3181dbaf96>
5. Givon U. Muscle weakness in cerebral palsy. *Acta Orthopaedica et Traumatologica Turcica* 2009; 43 (2): 87-93. <https://doi.org/10.3944/AOTT.2009.087>
6. Elder GC, Kirk J, Stewart G, Cook K, Weir D et al. Contributing factors to muscle weakness in children with cerebral palsy. *Developmental Medicine and Child Neurology* 2003; 45 (8): 542-550. <https://doi.org/10.1017/s0012162203000999>

7. Kara OK, Gursen C, Cetin SY, Tascioglu EN, Muftuoglu S et al. The effects of power exercises on body structure and function, activity and participation in children with cerebral palsy: an ICF-based systematic review. *Disability and Rehabilitation* 2022; 1-14. <https://doi.org/10.1080/09638288.2022.2138575>
8. Michelsen JS, Lund MC, Alkjær T, Finni T, Nielsen JB et al. Wearable electromyography recordings during daily life activities in children with cerebral palsy. *Developmental Medicine and Child Neurology* 2020; 62 (6): 714-722. <https://doi.org/10.1111/dmcn.14466>
9. Damiano DL, Vaughan CL, Abel ME. Muscle response to heavy resistance exercise in children with spastic cerebral palsy. *Developmental Medicine and Child Neurology* 1995; 37 (8): 731-739. <https://doi.org/10.1111/j.1469-8749.1995.tb15019.x>
10. Fowler EG, Staudt LA, Greenberg MB. Lower-extremity selective voluntary motor control in patients with spastic cerebral palsy: Increased distal motor impairment. *Developmental Medicine and Child Neurology* 2010; 52 (3): 264-269. <https://doi.org/10.1111/j.1469-8749.2009.03586.x>
11. Burtner P, Qualls C, Woollacott M. Muscle activation characteristics of stance balance control in children with spastic cerebral palsy. *Gait & Posture* 1998; 8 (3): 163-174. [https://doi.org/10.1016/s0966-6362\(98\)00032-0](https://doi.org/10.1016/s0966-6362(98)00032-0)
12. Moreau NG, Lieber RL. Effects of voluntary exercise on muscle structure and function in cerebral palsy. *Developmental Medicine and Child Neurology* 2022; 64 (6): 700-708. <https://doi.org/10.1111/dmcn.15173>
13. Kara OK, Livanelioglu A, Yardımcı BN, Soylu AR. The effects of functional progressive strength and power training in children with unilateral cerebral palsy. *Pediatric Physical Therapy* 2019; 31 (3): 286-295. <https://doi.org/10.1097/PEP.0000000000000628>
14. Merino-Andres J, Garcia de Mateos-Lopez A, Damiano DL, Sanchez-Sierra A. Effect of muscle strength training in children and adolescents with spastic cerebral palsy: A systematic review and meta-analysis. *Clinical Rehabilitation* 2022; 36 (1): 4-14. <https://doi.org/10.1177/026921552111040199>
15. Bland DC, Prosser LA, Bellini LA, Alter KE, Damiano DL. Tibialis anterior architecture, strength, and gait in individuals with cerebral palsy. *Muscle & Nerve* 2011; 44 (4): 509-517. <https://doi.org/10.1177/026921552111040199>
16. Di Giulio I, Maganaris CN, Baltzopoulos V, Loram ID. The proprioceptive and agonist roles of gastrocnemius, soleus and tibialis anterior muscles in maintaining human upright posture. *The Journal of Physiology* 2009; 587 (10): 2399-2416. <https://doi.org/10.1113/jphysiol.2009.168690>
17. Howard JJ, Herzog W. Skeletal muscle in cerebral palsy: from belly to myofibril. *Frontiers in Neurology* 2021; 12: 620852. <https://doi.org/10.3389/fneur.2021.620852>
18. Handsfield GG, Williams S, Khuu S, Lichtwark G, Stott NS. Muscle architecture, growth, and biological Remodelling in cerebral palsy: a narrative review. *BMC Musculoskeletal Disorders* 2022; 23 (1): 233. <https://doi.org/10.1186/s12891-022-05110-5>
19. Wiley ME, Damiano DL. Lower-extremity strength profiles in spastic cerebral palsy. *Developmental Medicine and Child Neurology* 1998; 40 (2): 100-107. <https://doi.org/10.1111/j.1469-8749.1998.tb15369.x>
20. Damiano DL, Kelly LE, Vaughn CL. Effects of quadriceps femoris muscle strengthening on crouch gait in children with spastic diplegia. *Physical Therapy* 1995; 75 (8): 658-667. <https://doi.org/10.1093/ptj/75.8.658>
21. Dodd KJ, Taylor NF, Graham HK. A randomized clinical trial of strength training in young people with cerebral palsy. *Developmental Medicine and Child Neurology* 2003; 45 (10): 652-657. <https://doi.org/10.1017/s0012162203001221>
22. Heo SG, Lee HS, Park SW. Effects of strengthening exercise on gait ability and GMFM in cerebral palsy: A systematic review and meta-analysis. *Journal of the Korean Society of Physical Medicine* 2018; 13 (3): 39-47. <https://doi.org/10.13066/kspm.2018.13.3.39>
23. Engsborg JR, Ross SA, Collins DR. Increasing ankle strength to improve gait and function in children with cerebral palsy: A pilot study. *Pediatric Physical Therapy* 2006; 18 (4): 266-275. <https://doi.org/10.1097/01.pcp.0000233023.33383.2b>
24. Jung JW, Her JG, Ko J. Effect of strength training of ankle plantarflexors on selective voluntary motor control, gait parameters, and gross motor function of children with cerebral palsy. *Journal of Physical Therapy Science* 2013; 25 (10): 1259-1263. <https://doi.org/10.1589/jpts.25.1259>
25. O'Sullivan SB, Schmitz TJ. *Physical Rehabilitation*. 5th ed. Philadelphia, PA, USA: F.A. Davis Company; 2006.
26. Blundell S, Shepherd R, Dean C, Adams R, Cahill B. Functional strength training in cerebral palsy: A pilot study of a group circuit training class for children aged 4–8 years. *Clinical Rehabilitation* 2003; 17 (1): 48-57. <https://doi.org/10.1191/0269215503cr584oa>
27. Cho HJ, Lee BH. Effect of functional progressive resistance exercise on lower extremity structure, muscle tone, dynamic balance and functional ability in children with spastic cerebral palsy. *Children* 2020; 7 (8): 85. <https://doi.org/10.3390/children7080085>
28. Palisano R, Rosenbaum P, Walter S, Russell D, Wood E et al. Gross motor function classification system for cerebral palsy. *Developmental Medicine and Child Neurology* 1997; 39 (4): 214-223. <https://doi.org/10.1111/j.1469-8749.1997.tb07414.x>
29. Gracies JM, Burke K, Clegg NJ, Browne R, Rushing C et al. Reliability of the Tardieu scale for assessing spasticity in children with cerebral palsy. *Archives of Physical Medicine and Rehabilitation* 2010; 91 (3): 421-428. <https://doi.org/10.1016/j.apmr.2009.11.017>
30. Tardieu G. A la recherche d'une technique de mesure de la spasticite (in French). *Revista de Neurologia* 1954; 91: 143-144.
31. Berg KO, Maki BE, Williams JI, Holliday PJ, Wood-Dauphinee SL. Clinical and laboratory measures of postural balance in an elderly population. *Archives of Physical Medicine and Rehabilitation* 1992; 73 (11): 1073-1080.

32. Franjoine MR, Gunther JS, Taylor MJ. Pediatric balance scale: A modified version of the berg balance scale for the school-age child with mild to moderate motor impairment. *Pediatric Physical Therapy* 2003; 15 (2): 114-128. <https://doi.org/10.1097/01.PEP.0000068117.48023.18>
33. Duncan PW, Weiner DK, Chandler J, Studenski S. Functional reach: A new clinical measure of balance. *Journal of Gerontology* 1990; 45 (6): 192-197. <https://doi.org/10.1093/geronj/45.6.m192>
34. Bartlett D, Birmingham T. Validity and reliability of a pediatric reach test. *Pediatric Physical Therapy* 2003; 15 (2): 84-92. <https://doi.org/10.1097/01.PEP.0000067885.63909.5C>
35. Chrysagis N, Skordilis EK, Koutsouki D. Validity and clinical utility of functional assessments in children with cerebral palsy. *Archives of Physical Medicine and Rehabilitation* 2014; 95 (2): 369-374. <https://doi.org/10.1016/j.apmr.2013.10.025>
36. Carey H, Martin K, Combs-Miller S, Heathcock JC. Reliability and responsiveness of the timed up and go test in children with cerebral palsy. *Pediatric Physical Therapy* 2016; 28 (4): 401-408. <https://doi.org/10.1097/PEP.0000000000000301>
37. Stegeman D, Hermens H. Standards for surface electromyography: The European project Surface EMG for non-invasive assessment of muscles (SENIAM). Enschede Roessingh Research and Development 2007: 108-112.
38. Cho C, Hwang W, Hwang S, Chung Y. Treadmill training with virtual reality improves gait, balance, and muscle strength in children with cerebral palsy. *Tohoku Journal of Experimental Medicine* 2016; 238 (3): 213-218. <https://doi.org/10.1620/tjem.238.213>
39. Dos Santos AN, Pavao SL, Rocha NACF. Sit-to-stand movement in children with cerebral palsy: A critical review. *Research in Developmental Disabilities* 2011; 32 (6): 2243-2252. <https://doi.org/10.1016/j.ridd.2011.05.001>
40. Sahrman AS, Stott NS, Besier TF, Fernandez JW, Handsfield GG. Soleus muscle weakness in cerebral palsy: Muscle architecture revealed with diffusion tensor imaging. *PloS One* 2019; 14 (2): e0205944. <https://doi.org/10.1371/journal.pone.0205944>
41. Schranz C, Kruse A, Belohlavek T, Steinwender G, Tilp M et al. Does home-based progressive resistance or high-intensity circuit training improve strength, function, activity or participation in children with cerebral palsy? *Archives of Physical Medicine and Rehabilitation* 2018; 99 (12): 2457-2464. <https://doi.org/10.1016/j.apmr.2018.06.010>
42. Agarwal A, Verma I. Cerebral palsy in children: An overview. *Journal of Clinical Orthopaedics and Trauma* 2012; 3 (2): 77-81. <https://doi.org/10.1016/j.jcot.2012.09.001>
43. Lorentzen J, Kirk H, Fernandez-Lago H, Frisk R, Scharff Nielsen N et al. Treadmill training with an incline reduces ankle joint stiffness and improves active range of movement during gait in adults with cerebral palsy. *Disability and Rehabilitation* 2017; 39 (10): 987-993. <https://doi.org/10.1080/09638288.2016.1174745>
44. Ryan JM, Lavelle G, Theis N, Noorkoiv M, Kilbride C et al. Progressive resistance training for adolescents with cerebral palsy: The STAR randomized controlled trial. *Developmental Medicine and Child Neurology* 2020; 62 (11): 1283-1293. <https://doi.org/10.1111/dmcn.14601>
45. Surana BK, Ferre CL, Dew AP, Brandao M, Gordon AM et al. Effectiveness of lower-extremity functional training (LIFT) in young children with unilateral spastic cerebral palsy: A randomized controlled trial. *Neurorehabilitation and Neural Repair* 2019; 33 (10): 862-872. <https://doi.org/10.1177/1545968319868719>
46. Jackman M, Sakzewski L, Morgan C, Boyd RN, Brennan SE et al. Interventions to improve physical function for children and young people with cerebral palsy: International clinical practice guideline. *Developmental Medicine and Child Neurology* 2022; 64 (5): 536-549. <https://doi.org/10.1111/dmcn.15055>
47. Tedroff K, Knutson LM, Soderberg GL. Synergistic muscle activation during maximum voluntary contractions in children with and without spastic cerebral palsy. *Developmental Medicine and Child Neurology* 2006; 48 (10): 789-796. <https://doi.org/10.1017/S0012162206001721>
48. Stackhouse SK, Binder-Macleod SA, Lee SC. Voluntary muscle activation, contractile properties, and fatigability in children with and without cerebral palsy. *Muscle & Nerve* 2005; 31 (5): 594-601. <https://doi.org/10.1002/mus.20302>
49. Rose J, McGill KC. Neuromuscular activation and motor-unit firing characteristics in cerebral palsy. *Developmental Medicine and Child Neurology* 2005; 47 (5): 329-336. <https://doi.org/10.1017/s0012162205000629>
50. Smith LR, Lee KS, Ward SR, Chambers HG, Lieber RL. Hamstring contractures in children with spastic cerebral palsy result from a stiffer extracellular matrix and increased in vivo sarcomere length. *The Journal of Physiology* 2011; 589 (10): 2625-2639. <https://doi.org/10.1113/jphysiol.2010.203364>
51. Rose J, McGill KC. The motor unit in cerebral palsy. *Developmental Medicine and Child Neurology* 1998; 40 (4): 270-277. [https://doi.org/10.1016/S1836-9553\(12\)70111-X](https://doi.org/10.1016/S1836-9553(12)70111-X)

Appendix: The Progressive functional exercise protocol

1. Stepping up on a 20 cm step



1. Stepping up sideways on a 20 cm step

