

Hülya PINARCIK<sup>1</sup>  
Zekeriya AKTÜRK<sup>2</sup>  
Fulya AKIN<sup>3</sup>  
Mehmet AKIN<sup>4</sup>  
Gülperi PINARCIK<sup>4</sup>

## Can Cystatin C be used as a Marker of Microalbuminuria?

Received: March 26, 2001

Departments of <sup>1</sup>Biochemistry, <sup>2</sup>Family Medicine, and <sup>3</sup>Internal Medicine, Haydarpaşa State Hospital, <sup>4</sup>Department of Pediatrics, Zeynep Kamil Hospital, Istanbul - TURKEY

Cystatin C is now widely used for the assessment of glomerular filtration rate in clinical practice (1,2). As microalbuminuria is the first sign of renal involvement in diabetes, the correlation of cystatin C with microalbuminuria is an important factor in deciding on the usefulness of this test for diabetic patients. At the same time, microalbuminuria is an early sign that appears before glomerular filtration rate (GFR) deteriorates (3).

In order to evaluate the correlation between serum cystatin C and microalbuminuria we measured the serum cystatin C and microalbuminuria levels of 64 type II diabetic patients. Cystatin C levels were elevated (greater than 1.7 mg/l) in 25 (92.59%) of the 27 microalbuminuric patients (Mean  $\pm$  SD = 2.29  $\pm$  0.43 mg/l) but in only 5 (13.51%) of the 37 non-microalbuminuric patients (Mean  $\pm$  SD 1.04  $\pm$  0.47 mg/l) ( $t = 10.886$ ;  $p < 0.0001$ ). The sensitivity and specificity of cystatin C in detecting microalbuminuria was calculated to

be 92.59% and 83.33% respectively. These results can be interpreted in favor of using cystatin C as a marker of microalbuminuria. Nevertheless, some factors, such as the glycemic control status of the patients, concomitant diseases, and drug therapy, may have affected the microalbuminuria as well. Unfortunately, it was not possible to eliminate these factors in this study. Hence, these findings need to be verified in a more standardized setting with respect to physical activity level, diabetic control, concomitant illnesses, and drug therapy in order to reveal microalbuminuria due to solely diabetic nephropathy.

**Correspondence author:**

Zekeriya AKTÜRK  
Trakya University Medical Faculty  
Department of Family Medicine  
22030 Edirne - TURKEY

### References

1. Newman DJ, Thakkar H, Edwards RG, Wilkie M, White T, Grubb AO, Price CP. Serum cystatin C measured by automated immunoassay: a more sensitive marker of changes in GFR than serum creatinine. *Kidney Int.* 47 (1): 312-316, 1995.
2. Nilsson-Ehle H, Grubb A. New markers for the determination of GFR: iohexol clearance and cystatin C serum concentration. *Kidney Int Suppl* 47: 17-24, 1994.
3. Chiarelli F, Verrotti A, Mohn A, Morgese G. The importance of microalbuminuria as an indicator of incipient diabetic nephropathy: therapeutic implications. *Ann Med* 29 (5): 439-445, 1997.