

Suspension of *Arthroderma* and *Trichophyton* species in RPMI-1640 medium provided long-term viability at room temperature

Çağrı ERGİN^{1*}, Macit İLKİT²

¹Department of Medical Microbiology, Faculty of Medicine, Pamukkale University, Denizli, Turkey

²Division of Mycology, Department of Microbiology, Faculty of Medicine, Çukurova University, Adana, Turkey

Received: 16.06.2014 • Accepted/Published Online: 23.10.2014 • Printed: 30.06.2015

To the Editor:

Research on dermatophytic fungi requires taxonomically well-defined strains. Furthermore, the long-term preservation of certain human pathogenic dermatophytes is crucial for mycological investigations. Subculturing is time-consuming, prone to contamination, and impractical, especially in routine practice. Although strain preservation under mineral oil, silica gel, soil, sand, or sterile water has been successful, periodic subculturing of these preserved strains is a major problem in laboratories with limited funds. Therefore, modified protocols of basic lyophilization and cryopreservation are the recommended storage methods in these settings (1).

In June 2008, we lyophilized standard dermatophyte species in sterile water in clear glass vials (AIM SV-15B) for vibrational spectroscopy analysis (2). The strains used were obtained from the reference collection of the CBS-KNAW Fungal Biodiversity Centre (CBS; Utrecht, the Netherlands); the strains along with their reference numbers are listed in the Table. Following the study period, 18 dermatophyte strains were resuspended with RPMI-1640 (with glutamine and without sodium bicarbonate, Sigma R6504). All suspensions were stored in a laboratory at room temperature, avoiding direct sunlight.

In March 2014, after about 6 years in storage, 15 (83.3%) of the 18 *Trichophyton* species were successfully subcultured onto Sabouraud glucose agar (HiMedia, India; without cycloheximide and chloramphenicol) (Table). Neither bacterial nor mold contamination was observed.

In clinical mycology laboratories, RPMI-1640 medium is routinely used in antifungal susceptibility testing (3). However, the incidental and unexpected findings of this study indicate that RPMI-1640 medium may also be suitable for strain preservation of *Trichophyton* species and its teleomorphs, i.e. *Arthroderma* spp. Moreover, because of its low cost, this method would be useful for most

laboratories, in particular for analyses that require long-term strain preservation, such as antifungal susceptibility testing, strain quality control evaluations, or molecular investigations.

Table. Viability testing results for *Arthroderma* and *Trichophyton* species stored in RPMI-1640 medium.

Species	CBS No.	Result
<i>Arthroderma</i> strains		
<i>A. simiii</i> (MT-)	417.65	+
<i>A. simiii</i> (MT+)	448.65	+
<i>A. vanbreuseghemii</i>	428.63	+
<i>T. mentagrophytes</i> complex		
<i>T. asteroides</i>	424.63	+
<i>T. erinacei</i>	344.79	+
<i>T. erinacei</i>	511.73	+
<i>T. erinacei</i>	677.86	+
<i>T. mentagrophytes</i>	110.65	+
<i>T. rubrum</i> complex		
<i>T. fluviomuniense</i>	592.68	+
<i>T. kuryangei</i>	422.67	-
<i>T. kuryangei</i>	518.63	+
<i>T. megninii</i>	389.58	+
<i>T. rubrum</i>	302.60	+
<i>T. rubrum</i>	392.58	-
<i>T. raubitschekii</i>	202.88	-
<i>T. raubitschekii</i>	287.86	+
<i>T. raubitschekii</i>	100084	+
<i>T. raubitschekii</i>	102856	+

MT, Mating type; +, growth; -, no growth.

* Correspondence: cagri@pau.edu.tr

References

1. Homolka L. Methods of cryopreservation in fungi. In: Gupta VK, Tuohy MG, Aychamy M, Turner KM, O'Donovan A, editors. *Laboratory Protocols in Fungal Biology: Current Methods in Fungal Biology*. Heidelberg, Germany: Springer; 2013. pp. 9–16.
2. Ergin Ç, İlkit M, Gök Y, Özel MZ, Çon AH, Kabay N, Söyleyici S, Döğen A. Fourier transform infrared spectral evaluation for the differentiation of clinically relevant *Trichophyton* species. *J Microbiol Methods* 2013; 93: 218–223.
3. Singh J, Zaman M, Gupta AK. Evaluation of microdilution and disk diffusion methods for antifungal susceptibility testing of dermatophytes. *Med Mycol* 2007; 45: 595–602.